Old Herborn University Seminar Monograph

32. Ageing and the microbiome

EDITORS:

PETER J. HEIDT JOHN BIENENSTOCK THOMAS C.G. BOSSCH MICHAEL ZASLOFF VOLKER RUSCH



Old Herborn University Seminar Monograph 32

ISBN 3-923022-44-1 ISSN 1431-6579

COPYRIGHT © 2018 BY THE OLD HERBORN UNIVERSITY FOUNDATION. THIS PUBLICATION OR PARTS OF IT MAY BE REPRODUCED OR TRANSMITTED IN ANY FORM UNDER THE RESTRICTION THAT THE SOURCE SHOULD BE MENTIONED. THIS PUBLICATION CAN ALSO BE FOUND ON, AND DOWNLOADED FROM: http://www.old-herborn-university.de/

EDITORS:

Prof. Dr. Peter J. Heidt Department of Pediatrics Leiden University Medical Center Albinusdreef 2 2333 ZA - Leiden The Netherlands

Prof. John Bienenstock, M.D. The Mcmaster Brain-Body Institute St. Joseph's Healthcare Hamilton 50 Charlton Avenue East Hamilton, Ontario L8N 4A6 Canada

Prof. Dr. Dr. h.c. Thomas C.G. Bosch Zoological Institute Am Botanischen Garten 1-9 D-24118 Kiel Germany Prof. Michael Zasloff, M.D., Ph.D. MedStar Georgetown Transplant Institute Georgetown University Hospital 3800 Reservoir Road, NW Washington, DC 20007 USA

Dr. rer. nat. Volker Rusch Old Herborn University Foundation Postfach 1765 D-35727 Herborn-Dill Germany



Old Herborn University

Old Herborn University Foundation Postfach 1765 D-35727 Herborn-Dill Germany Telephone: +49 - 2772 - 921100 Telefax: +49 - 2772 - 921101

Participating authors		
I.	INTRODUCTORY LECTURE: SOME INSIGHTS ABOUT AGEING FROM A SHARK AND AN OYSTER (<i>Michael Zasloff</i>) Introduction to the seminar Literature	1 1 2
Π.	PHYSIOLOGY OF HUMAN AGEING (<i>Robert J. Pignolo</i>) Summary	5 5 6 6 7 8 8 9 9 10 11 12 12 13 14 15 16 16 16 17 18 19
III.	AGEING AND THE MICROBIOME: AN AVENUE FOR INVENTION (<i>Brian K. Kennedy</i>) Summary Introduction Microbiome alterations during ageing Ageing interventions and the microbiome	25 25 25 26 27

	Microbiome interventions and ageing Conclusion Literature	28 28 29
IV.	HOST-MICROBIOTA INTERACTIONS DURING AGEING (Dario Riccardo Valenzano)	33
	Summary	33
	Ageing and homeostasis	33
	Gut microbiota contribute to host physiology	34
	Gut microbiota and immune function: a reciprocal influence	34
	Gut microbiota in ageing and disease	35
	Gut microbiota play a causal role in modulating host ageing and	
	life span	36
	Turquoise killifish as a model to study vertebrate immunosenescence Literature	37 38
V.	CONTRASTING CONSERVED AND TISSUE-SPECIFIC RESPONSES TO AGEING IN VERTEBRATES (Daniela Esser, Lena Best, and Christoph Kaleta)	43
	Summary	43
	Introduction	43
	Material and methods	46
	Results	47
	Discussion	51
	Acknowledgements	53
	Literature	54
VI.	AGEING AND THE MICROBIOME – LESSONS FROM NON-SENESCENT MODELS	
	(Benedikt M. Mortzfeld and Thomas C.G. Bosch)	59
	Summary	59
	The signature of ageing	59
	Microbes matter during the ageing process Non-senescent model organisms to study the interactions between	60
	FoxO and microbes	61
	Ageing antagonist FoxO controls both stem cells and microbes	63
	Conclusions	64
	Acknowledgements	65
	Literature	65

VII.	AGEING AND GUT TO BRAIN SIGNALLING (Wolfgang Kunze)	71
	Introduction	71
	Ageing general	71
	Microbiome	71
	Inflammation	72
	Gut motility	73
	Myenteric plexus	74
	Extrinsic afferents	75
	Intramural gatekeeper	76
	Literature	78
VIII.	STROKE AND MICROBIOTA – CURRENT RESEARCH AND CONCEPTS	
	(Joseph Anrater)	81
	Summary	81
	Introduction	81
	Effects of microbiota on immune cells involved in stroke	0.0
	pathophysiology	82
	Altered intestinal immune status and its effects on ischaemic stroke	83
	Pole of intestinal microbiota in post stroke infections	04
	Stroke microbiome and ageing	85
	Conclusions	85
	Literature	86
IX.	THE MICROBIOME, KMMUNE DEVELOPMENT, CHRONIC INFLAMMATION, AND CANCER	
	(Taylor N. Tibbs, Lacey R. Lopez, and Janelle C. Arthur)	91
	Introduction	91
	The immune system and microbiome in ageing	92
	Microbiome and cancer – a disease of ageing	96
	Literature	100
X.	AGEING AND THE MICROBIOME: SUMMARY OF THE 32 ND OLD HERBORN UNIVERSITY SEMINAR AND THE DISCUSSIONS	
	(Michael Zasloff)	107

Joseph Anrather, VMD, Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, 407 East 61st Street RR409, New York, NY10065, USA. (E-mail: joa2006@med.cornell.edu).

Janelle C. Arthur, PhD, Department of Microbiology and Immunology, Center for Gastrointestinal Biology and Disease, and Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, 125 Mason Farm Road, Chapel Hill, NC 27599, USA. (E-mail: janelle arthur@med.unc.edu).

Prof. Dr. Dr. h.c. Thomas C.G. Bosch, Zoological Institute, Christian-Albrechts-University Kiel, Am Botanischen Garten 1-9, D-24118 Kiel, Germany. (E-mail: tbosch@zoologie.uni-kiel.de).

Prof. Dr. Christoph Kaleta, Research Group Medical Systems Biology, Institute for Experimental Medicine, University of Kiel, Michaelisstraße 5, Kiel, Germany. (E-mail: c.kaleta@iem.uni-kiel.de).

Prof. Brian Kennedy, PhD, Departments of Biochemistry and Physiology, Yong Loo Lin School of Medicine, National University of Singapore, MD7, 8 Medical Drive, #02-11, Singapore 117597, Singapore. (E-mail: bkennedy@nus.edu.sg).

Wolfgang Kunze, PhD, The McMaster Brain-Body Institute & Department of Psychiatry, Juravinsky Innovation Tower, Room T-3306, St. Joseph's Healthcare, 50 Charlton Avenue East, Hamilton, Ontarion L8N 4A6, Canada. (E-mail: kunzew@mcmaster.ca).

Prof. Robert J. Pignolo, MD, PhD, Robert and Arlene Kogod Professor of Geriatric Medicine, Department of Medicine, Division of Geriatric Medicine and Gerontology, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA.

(E-mail: pignolo.robert@mayo.edu).

Dario Riccardo Valenzano, PhD, Max Planck Institute for Biology of Ageing, Joseph-Stelzmann-Straße 9b, 50931 Cologne, Germany. (E-mail: dario.valenzano@age.mpg.de).

Prof. Michael Zasloff, MD, PhD, Medstar Georgetown Transplant Institute, MedStar Georgetown University Hospital, 3800 Reservoir Road, NW, Washington, DC 20007, USA. (E-mail: maz5@georgetown.edu).

INTRODUCTORY LECTURE: SOME INSIGHTS ABOUT AGEING FROM A SHARK AND AN OYSTER

MICHAEL ZASLOFF

MedStar Georgetown Transplant Institute, Georgetown University School of Medicine, Washington, DC 20007, USA

The subject of the 32nd Old Herborn Seminar is on the impact of ageing on the microbiome. In this brief introductory lecture I would like to explore the subject of ageing. What underlies the ageing process? In general, ageing is a process that occurs as the regenerative potential of the organism diminishes. As a consequence, the functions of vital organs gradually fail. Why does this happen? Is there an ageing "clock" in the human body? Are we composed of cells that are genetically programmed to count the hours until their senescence is reached? Or is the rate of ageing controlled by a "master clock"? Can we slow down the ageing clock?

Perhaps if we examine some longlived animals we might discern some hints about the ageing mechanism. One of the oldest known animals is a clam collected off the coast of Iceland (Scourse et al., 2006). Based on the "rings" of the shell of this animal, it was estimated to be about 500 years old, and called the "Ming Clam" to highlight that it began its life during the Chinese Ming dynasty. Other clams caught in that area were similar in age, suggesting that the Ming was not an unusual outlier. Clearly these individuals survived predation. As extraordinary they survived infection and malignancy, despite (or because of?) the absence of an adaptive immune system. This animal teaches us that a robust adaptive immune system is not necessarily required for longevity.

The Greenland shark was recently

discovered to be the oldest living vertebrate, with the maximum age estimated to be 600 yrs. (Nielsen et al., 2016). The ages of the animals were determined by radiocarbon dating of the lens of the eye, which is formed during embryogenesis. Prior studies had shown that the Greenland shark reached sexual maturity at about 200 years. The ageing clock of this animal appears to run about 10-fold more slowly than the human's. This animal teaches us that tissue renewal and organ regeneration, including reproductive capacity, can be sustained in a vertebrate for hundreds of years.

In 1961 Hayflick and Moorehead proposed that an animal's longevity had its basis at the cellular level (Hayflick and Moorehead, 1961). They demonstrated that the fibroblasts from short lived animals would undergo fewer divisions in cell culture before reaching senescence compared with the cells of longer-lived animals. It was later suggested that the number of cellular divisions a cell was capable of sustaining was dependent upon the integrity of the chromosomal telomeres, and the expression of telomerase. With each successive division, the telomeres of the stem cells shortened, until chromosomal replication was compromised.

Subsequent studies failed to validate the relationship between the longevity of a species and the number of divisions to senescence. A rather telling study asked whether the fibroblasts grown from a biopsy taken from a 90+ year old human would undergo senescence after fewer divisions in vitro than those from younger individuals (Maier et al., 2007). The results demonstrated that the fibroblasts from the nonagenarians behaved very much like young cells with respect to the maximal number of replications. These observations suggest that the replicative capacity of a stem cell is influenced by factors contributed by the *in vivo* setting. What anti-senescence factors are present in vivo, but absent in vitro? Might a gradual decrease in "anti-senescence "hormones" be responsible for ageing?

Caloric restriction has been shown to increase longevity in flies, worms, fish and mice (Balasubramanian et al., 2017), so better understanding of the physiological consequences of caloric restriction could provide some insight into the processes influencing longevity: IGF-1, insulin, and growth hormone concentrations fall in serum (Bartke Westbrook, 2012; Cady and and Sadagurski, 2017); plasma ketone bodies increase, providing the brain with a highly efficient source of energy, reducing its dependence on glucose; and most curious, reduced inflammation within the hypothalamus (Cady and Sadagurski, 2017).

A recent study suggests that

progressive hypothalamic inflammation might be a key factor in systemic ageing (Zhang et al., 2013). In this study, a lentivirus expressing GFP driven by an NF- κ B response element was injected to either the hypothalamus or the cortex of mice. As the animals aged, the intensity of GFP+ cells increased in the hypothalamus, but not in the cortex, demonstrating that the hypothalamus experiences an inflammatory milieu in contrast to the cortex. The hypothalamus has a central role in homeostatic regulation of metabolism, vital functions (temperature, blood pressure, etc), and growth, so inflammation could impact negatively on function. Indeed, in this study numerous hypothalamic peptides were examined to determine which, if any, changed in the older animals, and GnRH, which decreased, was identified as a candidate. Surprisingly, treatment of the older animals with parenteral GnRH stimulated neurogenesis in the brain. Most surprising GnRH treatment increased muscle strength, skin thickness, and cognitive functions, in a sense, restoring a more youthful phenotype.

So what controls ageing? Does inflammation within the hypothalamus drive generalized ageing of our bodies? Is it all that simple?

LITERATURE

- Balasubramanian, P., Howell, P.R., and Anderson, R.M.: Aging and Caloric Restriction Research: A Biological Perspective With Translational Potential. EBioMedicine 21: 37-44 (2017)
- Bartke, A. and Westbrook, R.: Metabolic characteristics of long-lived mice. Front. Genet. 3, 288 (2012).
- Cady, G. and Sadagurski, M.: Targeting neuroinflammation - a potential for antiaging interventions. Aging (Albany NY) 9,

1951-1952 (2017).

- Hayflick, L. and Moorhead, P.S.: The serial cultivation of human diploid cell strains. Exp. Cell Res. 25, 585-621 (1961).
- Maier, A.B., le Cessie, S., de Koning-Treurniet, C., Blom, J., Westendorp, R.G., and van Heemst, D.: Persistence of high-replicative capacity in cultured fibroblasts from nonagenarians. Aging Cell 6, 27-33 (2007).
- Nielsen, J, Hedeholm, R.B., Heinemeier, J., Bushnell, P.G., Christiansen, J.S., Olsen, J.,

Ramsey, C.B., Brill, R.W., Simon, M., Steffensen, K.F., and Steeffensen, J.E: Eye lens radiocarbon reveals centuries of longevity in the Greenland shark (Somniosus microcephalus). Science 353, 702-704 (2016).

Scourse, J., Richardson, C, Forsythe, G., Harris, I., Heinemeier, J., Fraser, N., Briffa, K., and Jones, P.: First cross-matched floating chronology from the marine fossil record: Data from growth lines of the longlived bivalve mollusc Arctica islandica. The Holocene 16, 967-974 (2006).

Zhang, G., Li, J., Purkayastha, S., Tang, Y., Zhang, H., Yin, Y., Li, B., Liu, G., and Cai, D.: Hypothalamic programming of systemic ageing involving IKK-β, NF-κB and GnRH. Nature 497, 211-216 (2013).

PHYSIOLOGY OF HUMAN AGEING

ROBERT J. PIGNOLO

Department of Medicine, Division of Geriatric Medicine and Gerontology, Mayo Clinic College of Medicine, Rochester, MN, USA

SUMMARY

The study of human ageing physiology is limited by confounders such as concomitant disease, as well as the role of environmental factors and interactions with the physical world. The objective of this chapter is to review human physiological ageing through primarily a systems-based approach, with consideration for the geriatric syndromes (multifactorial pathophysiology), resiliency, and common ageing mechanisms that may underlie declines in multiple physiologic systems. Each system has its own trajectory for age-related dysfunction with accompanying system-specific declines. There are a growing number of examples of factors that regulate the resilient responses to stressors and mitigate potential adverse physiological consequences. Common mechanisms that contribute to human age-related physiology include chronic, low-grade, non-microbial inflammation, cellular senescence, accumulation of damaged macromolecules (DNA, proteins, carbohydrates, lipids), and stem and progenitor cell dysfunction. Ageing physiology in humans is heterogeneous and almost always interdependent with other systems and extrinsic influences.

INTRODUCTION

The study of human ageing physiology is confounded by chronic diseases not easily distinguished from primary ageing processes and for which ageing is the major risk factor. Nevertheless, and to the extent possible, ageing physiology will be reviewed as a systemsbased approach, highlighting the major or key ageing features seen in healthy individuals. For many system-based ageing changes, the aetiologies are multifactorial or syndromic, and the geriatric syndromes such as pressure ulcers, incontinence, falls, dementia, and osteoporosis are good examples of the clinical consequences due predominately to ageing physiology in skin, the genitourinary system, muscle, central nervous system, and bone, respectively,

but also secondary to contributions from physiologic ageing in other systems and extrinsic factors (e.g., polypharmacy, amount of daily activity, etc.).

Human ageing physiology is heterogeneous among individuals, lending itself to the descriptors of "normal" and "successful" ageing and supporting the concept of resiliency in the case of the latter. Resilience to physiologic ageing has been demonstrated across several systems, although its description in humans, outside of psychological and behavioural studies, is relatively limited (*Hadley* et al., 2017). Common mechanisms contribute to ageing physiology in animal models, and control of these mechanisms likely mediate resistance effects. In humans, early evidence suggests that the same common ageing mechanisms are at play (*Kirkland*, 2016).

LIMITATIONS TO STUDYING HUMAN AGEING PHYSIOLOGY

Declines with age in physiological systems are commonly due to the effects of disease superimposed on fundamental ageing processes. A major challenge therefore is to distinguish the effects of ageing from disease. In humans, this can only be approximated by studying healthy individuals, tissue donors, those who experience non-disease related sudden death, or probably less helpful, autopsied individuals where the circumstances of death can be obtained. With this caveat, the physiology of system-specific human ageing is often considered to reflect primary process(es).

Superimposed on primary ageing processes, are environmental factors and interactions with the physical world that may regulate the degree and pace of these processes. For example, barrier facilities for animal models limit understanding of ageing processes influenced by the microbiome and immunological responses to environmental pathogens, which are applicable to animals in the wild and to humans. Behavioural factors, which rely on interactions with individuals, are also relevant since purposeful living, social networks and support systems influence health outcomes and mortality risk (Boyle et al., 2010a,b; Holt-Lunstad et al., 2010; Park et al., 2014).

Death is a poor endpoint to study ageing. Mortality is tightly linked to disease state and accidental death where ageing physiology is obscured by disease pathology and events unrelated to ageing, respectively. Furthermore, in the wild most animals will never achieve longevity close to their maximum life span because of death due to accidental reasons and to predation.

Finally, a specific physiological system may be dependent on multi-system interactions, and so dysfunction in one system may be related to dysfunctional processes in another or multiple systems. This relationship is true in disease states such as cardiopulmonary, cardiorenal, and hepatorenal syndromes, where dysfunction due to disease in one system contributes to the progression of disease in another system. The geriatric syndromes represent the equivalent dysfunctional outcomes due to ageing in one system being influenced by ageing in other systems and/or multiple age-related aetiologies that may be intrinsic or extrinsic to the individual. Therefore, examination of age-related changes in a single physiological system in isolation is confounded by the contributions of ageing in interacting systems.

DISRUPTION OF PHYSIOLOGIC RHYTHMS WITH AGEING

With ageing, disruption of circadian patterns cause phase advances (1-2 hours earlier) such as in the 24-hour body temperature trough and sleep onset. Suprachiasmatic nucleus neuronal loss in the hypothalamus results in attenuated pulsatile secretions and decreases in gonadotropins, growth hormone (GH), thyrotropin, melatonin, and adrenocorticotropic hormone (ACTH) (*Veldhuis*, 1997). There is also a delayed reset to a new photoperiod with ageing (*Hofman* and *Swaab*, 2006).

AGE-RELATED CHANGES IN THE GASTROINTESTINAL SYSTEM

Oropharynx

With ageing, there is a decrease in acinar cells and saliva production that is accompanied by thinning of the oral mucosa (Smith et al., 2013). These changes contribute to dry mouth complaints, recession of gums over time, and the increased risk of root carries. Loss of teeth promotes less effective mastication. Abnormal transfer of the food bolus to the pharynx (due to decreased oesophageal muscle compliance) is a major cause of inadequate nutritional intake in edentate individuals and places them at increased aspiration risk (Fulp et al., 1990; Frederick et al., 1996).

Oesophagus

There are four pathophysiological processes that occur in the oesophagus during ageing, including hypertrophy, nerve loss, decreased contraction/tone, and decreased sensation. Skeletal muscle hypertrophy of the upper one-third of the oesophagus carries no functional consequences (*Hall* et al., 2005). There is loss of myenteric ganglion cells which cause a decline in the amplitude of oesophageal contractions during peristalsis, but this does not substantially cause impairment of food movement (Hall et al., 2005). However, major deficits in secondary oesophageal contractions and a decrease in lower oesophageal sphincter tone causes increased gastric acid exposure (Kekki et al., 1982; Feldman et al., 1996; Hurwitz et al., 1997; Haruma et al., 2000,). Impaired sensation of distension at the distal oesophagus contributes to loss of symptomatology from injury (Lasch et al., 1997).

Stomach

Acidification of gastric contents with ageing is intact in the basal unstimulated state; however, there is an increased prevalence of *H. pylori* infection (*Marshall*, 1994; *Haruma* et al., 2000). Higher rates of gastritis, decreased production of prostaglandins, bicarbonate, and non-parietal fluid in the setting of decreased gastric emptying and microcirculation lead to impaired mucosal defences (*Guslandi* et al., 1999; *Tarnawski* et al., 2014). There is also a greater susceptibility to injury, with impaired healing and reduced efficacy of ulcer-healing drugs (*Tarnawski* et al., 2014).

Small intestine

Moderate villus atrophy and decreased absorption of micronutrients (e.g., calcium, folic acid, vitamin B12) with ageing are accompanied by an increase in bacterial overgrowth (*Parlesak* et al., 2003; *Salles*, 2007). Loss of innervation, especially of sensory and myenteric nerves, result in a higher incidence of painless ulcers (*Hilton* et al., 2001).

Large intestine

Mucosal atrophy, atrophy of the muscularis externus and altered coordination of contraction contribute to decreased colonic motility and the much higher rates of constipation with ageing (Bitar and Patil, 2004). Loss of myenteric plexus and intrinsic sensory neurons and concomitant lower anal sphincter tone lead to faecal incontinence (*Wade* and *Hornby*, 2005). Hypertrophy of mucularis mucosa with atrophy of the muscularis externus predispose to diverticuli (Comparato et al., 2007; Commane et al., 2009). Barrier function of colonic epithelium is compromised with ageing (Tran and Greenwood-Van Meerveld, 2013).

Hepatobiliary system

Liver mass and perfusion decline with ageing (*Schmucker*, 2005). There is

impairment in synthetic function, with a slight decrease in albumen levels, lower LDL and LDL receptor levels, lower cytochrome P450 content and lower vitamin K-dependent clotting factors (*Sotaniemi* et al., 1997; *Fu* and *Nair*, 1998; *Schmucker*, 2005). The regenerative response to injury decreases and the bile lithogenic index increases with ageing (*Valdivieso* et al., 1978; Schmucker, 2005).

Exocrine pancreas

There are minor atrophic fibrotic changes in the ageing pancreas with an increased number of cysts and side branching of ducts (*Bulow* et al., 2014). Although there is a decline in stimulated pancreatic flow, this does not impact exocrine function.

CARDIOVASCULAR AGEING

Heart weight increases as a function of ageing, with enlargement of the left atrium and enlargement and hypertrophy of the left ventricle (*Kitzman* et al., 1988). A decrease in cardiomyocyte number is accompanied by an increase in cardiomyocyte size (*Olivetti* et al., 1991). The aortic valve and mitral annulus thickens with development of calcific deposits. Mitral annular calcification predisposes to conduction abnormalities. Increased atrial and ventricular premature beats become more common with age (*Fleg* and *Kennedy*, 1992).

Due mostly through compensatory mechanisms, ejection fraction, stroke volume, and cardiac output do not change substantially with ageing (*Ferrari* et al., 2003). There is an increase in early diastolic filling, and an increase in end-diastolic filling due to a compensatory greater contribution from atrial systole (marked by a normal S4 in older individuals). There is a relative loss of chronotropic and inotropic responsiveness to β -adrenergic stimuli/catecholamines as well as inotropic response to digitalis glycosides. Peak cardiac output to maximal exercise is decreased.

In the vasculature, arterial wall thickness (intima-media) increases with advancing age, as does pulse wave velocity and total peripheral resistance (*Mitchell*, 2008). Endothelial nitric oxide release and β -adrenergic-mediated vasodilation decreases (*Mitchell*, 2008).

AGEING OF THE RESPIRATORY SYSTEM

Three hallmark features describe ageing in the respiratory system: change in lung volumes, increased alveolar-arterial oxygen gradient, and ventilationperfusion mismatch (*Chan* and *Welsh*, 1998). Due to decreased elastic recoil, increased chest wall rigidity, and a loss of force-generating capacity of respiratory muscles, forced vital capacity, forced expiratory volume at 1 second, and vital capacity decrease and functional residual capacity increases (*Janssens*, 2005). Diminished lung diffusion capacity results in an increase in the alveolar-arterial oxygen gradient, measured as the diffusion capacity of carbon monoxide which drops by 2-3 ml/min/mmHg per decade (*Stam* et al., 1994). Secondary to decreased elastic recoil, the intrapleural pressure becomes less negative causing areas of the lung base to close. With this closure, there is a redistribution of inspired air to underperfused apical areas. The resultant increase in physiological dead space and underventilation of dependent lung areas is responsible for ventilation-perfusion mismatches typical of the ageing lung (Stam et al., 1994).

AGEING OF THE IMMUNE SYSTEM

Ageing of the immune system is reflected by diminished immune responsiveness, so-called "immunosenescence", altered immune system physiology, and impaired immune regulation. Due to immunosenescence (Agarwal and Busse, 2010), there is a decreased response to new antigens/elevated susceptibility to infection and cancer, decreased vaccine efficiency (e.g., influenza), reactivation of latent infections (e.g., herpes simplex virus, tuberculosis), and perhaps compromised immune surveillance.

Important changes in immune system physiology with ageing include decreased production and maturation of B- and T-cells (e.g., marrow decline, thymic involution), inversion in proportional representation of memory vs naïve cells (T memory cells increase and T naïve cells decrease with age), decreased isotype switching and affinity maturation of B cells, accumulation of CD28-negative T-cells, and impaired formation of the T-cell – Antigen Presenting Cell "immune synapse" (*Panda* et al., 2009).

Altered immunoregulation can be seen as an increase in autoimmune syndromes (systemic lupus erythematosus, rheumatoid arthritis, and systemic sclerosis, others), oligoclonal expansion of T- and B-cells (decrease in diversity with age), monoclonal gammopathies, and increased inflammation (*Ramos-Casals* et al., 2004; *Arai* et al., 2015; *Ray* and *Yung*, 2018).

CHANGES IN THE AGEING KIDNEY

Kidney weight and volume remain unchanged with ageing in individuals who experienced sudden death and in healthy individuals, respectively (*Glassock* and *Rule*, 2012). Nephrosclerosis increases progressively with age (Glassock and Rule, 2012). A sclerosis score is the total number of histological abnormalities (global glomerular sclerosis, any tubular atrophy, interstitial fibrosis > 5%, any arteriosclerosis). Among living kidney donors with less than 10% glomerulosclerosis, glomerular density declines with age but the size of glomeruli and tubules increase; in those with greater than 10% glomerulosclerosis, glomerular density increases with age with a rise in the number of small sclerotic glomeruli and tubular atrophy. The number of functional glomeruli decreases with age

in both live kidney donors and in individuals at autopsy. There is compensation by functional nephrons in older healthy individuals. Glomerular filtration rate (GFR) is mostly reported to decline with age, but is variable and may be unchanged in as much as onethird in healthy cohorts (Lindeman et al., 1985; *Esposito* et al., 2007). Estimated GFR tends to vary based on the formula employed, and can be underestimated in healthy older adults when age is used as a surrogate for muscle mass. Changes in renal function with age may in part be driven by a vascular process whereby vasoconstrictive responses to angiotensin are intact but the vasodilatory responses to acetylcholine or to an acute sodium load are impaired (Hollenberg et al., 1974).

Parameter	Change with age	Response to challenges with age
Red cell life span	=	
Iron turnover	=	
Blood volume	=	
Bone marrow mass/cellularity	\downarrow	
Bone marrow fat	1	
Donor suitability for haematpoietic cell transplantation	\downarrow	
Tolerance of chemotherapy	\downarrow	\downarrow
Compenstory response to: Phlebotomy Hypoxia		↓ ↓
Colony size of stimulated erythroid progenitor cells		Ļ
Production of BM stimulatory hormones (e.g. SCF, GM-CSF, IL-	.3)* ↓	Ļ
Total WBC	=	
Clonal expansion of specific white blood cells	Ť	
Platelets (total number)	=	
Platelet responsiveness	↑	
Bleeding time	\downarrow	\downarrow
Procoagulant state (↑fibrinogen, ↑Factors V, VII, VIII, IX, ↑kininogen, ↑prekalikrein	Ţ	ſ
Fibrin degradation products (D-dimension	\$) ↑	
Plasminogen activator inhibitor-1 (major inhibitor of fibrinolysis)	Ť	
Deep vein thrombosis	Ť	\uparrow

Table 1: Changes in the haematopoietic system with age

 \uparrow : Increase; \downarrow : Decrease; =: Unchanged.

*SCF: Stem cell factor; GM-CGF: Granulocyte-Macrophage-Colony-stimulating factor; IL-3: Interleukin-3.

AGEING IN THE HAEMATOPOIETIC SYSTEM

Changes in the haematopoietic system with age are shown in Table 1. Major changes outside of the immune system (see above) include decreased bone marrow, marrow cellularity, and compensatory responses to phlebotomy and hypoxia as well as increased components related to the procoagulant state

Hormone	Change with ageing
Growth hormone	N in males; ↓ in females
Insulin-like growth factor I	\downarrow
ACTH	Ν
Cortisol	Ν
DHEA	\downarrow
Renin	Ν
Aldosterone	\downarrow
TSH	N or ↑
Τ4	Ν
Τ3	\downarrow
РТН	\uparrow
Calcitonin	\downarrow
1,25 (OH)2 D	\downarrow
LH	\uparrow or N in males; \uparrow in females
FSH	\uparrow or N in males; \uparrow in females
Testosterone	\downarrow or N in males; \uparrow in females
Free testosterone	\downarrow
Atrial natriuretic factor	\uparrow
Insulin	1
Glucagon	\downarrow

Table 2: Endocrine changes with ageing

↑: Increase; ↓: Decrease; N: Normal.

DHEA: dehydroepiandrosterone; TSH: thyroid stimulating hormone; T: thyroxine;

PTH: parathyroid hormone; LH: luteinizing hormone; FSH: follicle stimulating hormone.

(*Lipschitz* et al., 1984; *Pinto* et al., 2003; *Franchini*, 2006). In a young adult, haematopoietic marrow (red) resides in skull, vertebra, flat bones, and proximal femoral and humeral meta-

physes. Fatty marrow (yellow) predominates in remainder of skeleton, and gradually replaces red marrow with age (*Custer* and *Ahlfeldt*, 1932; *Kricun*, 1985).

ENDOCRINE CHANGES WITH AGEING

Table 2 summarizes the endocrine changes with ageing (*Lamberts* et al., 1997). In healthy individuals, levels of some hormones are not altered with age (e.g., thyroid hormone). In older individuals without disease, the homeostatic setpoint for a given hormone may be the same as in youth, at expense of alteration in another (e.g., "normal" testosterone with elevated LH), different from youth (e.g., DHEA, IGF-1 levels decline with age) or the same as in youth, except with a stressor (e.g., cortisol response).

There is an increase in frequency of endocrine disease with ageing (e.g., hypothyroidism, post-menopausal osteoporosis) and often older individuals have different presentations of endocrine disease (e.g., hypothyroidism, hyperthyroidism) (*Gambert* and *Escher*, 1988). Different presentations



Figure 1: Changes in adipose tissue with ageing.

of non-endocrine disease are also common when coexistent with endocrine disease (e.g., angina in the setting of hyperthyroidism). In addition, there are alterations in hormones secondary to medications frequently prescribed in the elderly (e.g., hyperprolactinemia from phenothiazines).

Although not as abrupt as the de-

testosterone ("andropause"), DHEA ("adrenopause"), and GH/IGF-I ("somatopause"). These changes, mediated by declines in LH/FSH, ACTH, and GH, respectively, are controlled at the level of the hypothalamus-pituitary complex (*Jones* and *Boelaert*, 2015).

cline in oestrogen levels during men-

opause, there are inexorable declines in

AGE-RELATED CHANGES IN ADIPOSE TISSUE

Two major changes in adipose tissue that occur with ageing are the re-distribution of fat and the decrease in lean body mass (Figure 1). Adipose tissue redistributes from subcutaneous to visceral depots (*Kuk* et al., 2009). This redistribution is accompanied by an increase in circulating fatty acids and inflammatory cytokines as well as insulin resistance and impaired glucose metabolism (*Stout* et al., 2017). When older individuals are compared to younger ones, even after controlling for waist circumference, visceral fat increases and subcutaneous fat decreases (*Kuk* et al., 2009).

MUSCLE AGEING

The hallmark of muscle aging, sarcopenia, is operationally defined by thresholds of muscle loss, decreased muscle strength, and lower physical performance (*Locquet* et al., 2018). Risk of sarcopenia is very well predicted by the combination of age, grip strength, and calf circumference (*Locquet* et al., 2018). Loss of muscle mass as a proportion of body weight is almost universal with ageing and muscle loss in the lower extremities is greater than



Figure 2: Contributing factors to muscle ageing (left) and consequences of sarcopenia (right).

that in the upper extremities. The loss of muscle mass is attributed to the decrease in motor unit number and size, lower physical activity common in older individuals, elevated inflammatory cytokines, diminished testosterone, oestrogen, and GH/IGF-I levels, elevations in myostatin and activin A, as well as nutritional deficits and genetic predisposition (Figure 2) (*Narici* and *Maffulli*, 2010). Histomorphologically, aged muscle shows increased fatty infiltration, fibrosis, decreased fast twitch > slow twitch fibers, smaller myofibril size, and fewer myofibrils (*Faulkner* et al., 2007).

Functional and clinical consequences of muscle ageing include decreased strength, increased fatigability, shorter time to peak tension and time to relaxation, decreased regenerative capacity, greater insulin resistance. lower volume of distribution for watersoluble drugs and decreased levels of glycolytic enzymes (Narici and Maf*fulli*, 2010). Increased mortality is associated with sarcopenia (Reinders et al., 2015).

SKIN AGEING

Primary changes that occur with skin ageing include epidermal thinning, decreased vascularity, and the loss of rete pegs which project downward from the epidermis between the dermal papillae (*Montagna* and *Carlisle*, 1990). The elastin network of the skin is diminished, there is a loss of subcutaneous fat, and epidermal turnover is slowed (*Uitto*, 2008). Histologically, there is a loss of melanocytes, Langerhans' cells, sweat and sebaceous glands, as well as Meissner's and Pacinian corpuscles (*Yaar* and *Gilchrest*, 2001). Microarchitectural changes lead to poor nutrient transfer, a loss of protective lipids, and decreased area for heat transfer. Bleeding can more easily occur into dermal-epidermal space. Clinically, the ageing skin demonstrates an



Figure 3: Ageing in the central nervous system by major anatomic, pathophysiologic, as well as cognitive and behavioural changes.

increased fragility to shear stress, slower wound healing, xerosis, reduced barrier function, compromised thermoregulation, wrinkling and sagging, decreased sensory perception, and purpura.

CENTRAL NERVOUS SYSTEM (CNS) AGEING

Key features of ageing in the CNS are outlined in Figure 3. Anatomically, the brain volume decreases, with predominant losses in the frontal and temporal lobes and with greater loss of white versus grey matter. Cerebral blood flow declines. Apoptotic neuronal loss becomes prominent in the cerebellum and cerebral cortex (*Dorszewska*, 2013). There is a reduction of the dendritic tree, loss of neuronal processes, and diminution of synapses (*van der Zee*, 2015).

Despite compensatory repatterning due to neuronal loss, the rate of neuronal loss exceeds that of neuronal formation. Pathological findings such as increased neurofibrillary tangles and senile plaques become more evident, although are not present to the extent that would be found in Alzheimer's disease (Fjell et al., 2014). There is reduced acetylcholine availability due to loss of cholinergic and muscarinic neurons, as well as lower synthesis and release of acetylcholine. Diminished levels of dopamine are found in the substantia striatum and nigra (Klostermann et al., 2012).

Cognitively, episodic and working memory declines (*Wilson* et al., 2002). There is a loss of specific skills and

knowledge with reduced processing speed and language ability. Executive function declines, as doses attention span, problem solving, as well as processing and learning new information. Procedural, primary, and semantic memories are unchanged with age. Recognition of familiar objects and faces and visual perception remain unaffected in healthy older adults.

SENSORY CHANGES WITH AGE

Vision

Periorbital tissues atrophy and lacrimal gland function, tear production, and goblet cell function are reduced with ageing (*Van Haeringen*, 1997). Despite decreased tear production, watery eyes are a common phenomenon since tissue atrophy leads to less effective drainage.

With ageing, the lens increases in thickness and hardness and the ciliary muscle decreases in strength; together, presbyopia or farsightedness results (Strenk et al., 2005). The lens also vellows and becomes more opaque with age, lessening colour discrimination. Increased light scatter due to lens alterations also reduces contrast. The resting pupil size and responsiveness decreases. In older individuals, the cornea yellows, undergoes focal and generalized thickening, and is more likely to have Hasall-Henle warts and especially arcus senilis (Salvi et al., 2006). There is a decrease in the production and drainage of the aqueous humour in the ageing eye, reduced speed and accuracy of pursuit and saccadic eye movements, and accumulation of scleral calcifications (Cogan's plaques) (Salvi et al., 2006). There is a loss of retinal pigment epithelial cells which contributes to the decreases in number, size, and function of rods in the retina (*Liem* et al., 1991). Although the number of cones in the fovea is unchanged with age, their function is reduced. Vitreous floaters due to collagen condensations increase with age. Blood flow to the visual cortex is decreased.

Hearing

In the ageing ear, there is a loss of hair cells in the organ of Corti, decreased innervation of cochlear and auditory centres in the brain, stiffness and calcification of the basilar membrane of the sensory apparatus, thickening of the stria vascularis capillaries (source of endolymph), and degeneration of the spiral ligament (Howarth and Shone, 2006). Hearing loss due to these changes are usually of high frequency sound (presbycusis) and result in reduced speech discrimination, poor localization of the sound source, and decreased ability to discriminate between target sound and background noise (Howarth and Shone, 2006).

Taste and smell

There is a reduction in papillae on tongue and decreased taste sensitivity with ageing. Uneven gustatory deficiencies across the tongue exist, but the loss of taste in older individuals is due mostly to an impaired sense of olfaction (*Boyce* and *Shone*, 2006).

Detection thresholds for smelling are altered with age owing in part to a decrease in the number of sensing neurons (*Boyce* and *Shone*, 2006). This results in an underappreciated reduction in appetite, the extremes of which can be best appreciated in the observation that people with anosmia forget to eat.

AGEING OF THE GENITOURINARY SYSTEM

Bladder

Decreased innervation of the detrusor muscle and CNS changes lead to diminished detrusor contractility, lower maximum bladder capacity and flow rate, reduced ability to withhold voiding, and an increased post-void residual volume (Elbadawi et al., 1998). The constellation of these age-related alterations greatly increases the risk for urinary incontinence. The decline in oestrogen in women leads to a shorter urethral length, a reduced maximum urethral closure pressure, and less effective urethral barrier function (Capobianco et al., 2012); together, these factors predispose women to an increased risk of urinary tract infection.

Male reproductive system

Age-related neurologic, vascular, and endocrine changes in men lead to the greater stimulation required for erection, decreased spontaneous erections, diminished firmness of erections, lower force of ejaculation, smaller ejaculate volumes, and increased refractory times between erections (*Seftel*, 2005). Despite these changes, reports of lower sexual activity in older men are variable (*Lindau* and *Gavrilova*, 2010). However, an objective decline in male reproductive ability with advanced age is likely due to lower Leydig cell number, degeneration of the seminiferous tubules leading to diminished sperm production, an increase in sperm chromosomal abnormalities, reduced sperm motility, and a reduced ability of sperm to fertilize eggs (*Harris* et al., 2011). Enlargement of the prostate gland with ageing is common.

Female reproductive system

The loss of oocytes and ovarian dysfunction in older women is primarily responsible for the decline in oestrogen during the perimenopausal period and the reduction in implantation efficiency, decreased likelihood for pregnancy, lower number of live births, and poor success of in vitro fertilization (Tarlatzis and Zepiridis, 2003). Oestrogen deficiency causes diminished vaginal elasticity, reduced clitoral engorgement after stimulation, increased vaginal dryness and atrophy, as well as decreased cervicovaginal secretions (Kingsberg, 2002). Oestrogen deficiency is responsible for an increase in vaginal pH, which predisposes to colonization by enteric flora.

SYNDROMIC APPROACH

For many if not all system-specific ageing changes, multiple aetiologies are at play, including the influence of ageing changes in other physiologic systems. While characterizing ageing in a single system *in isolation* serves to simplify the task, it also minimizes the true nature of its complex regulation. The geriatric syndromes serve as examples of the complexity involved to fully explain common age-related changes. Geriatric syndromes have been operationally defined as multi-factorial, having multi-system involvement, and sharing risk factors such as older age, functional impairment, and impaired mobility (*Inouye* et al., 2007). Common geriatric syndromes include pressure ulcers, incontinence, falls, delirium, dementia, and osteoporosis.

Osteoporosis is the hallmark of bone ageing (*Chandra* et al., 2018) and will be used here as an illustration of the syndromic approach to describing physiologic ageing in the skeleton (Figure 4). Bone loss is ultimately the



Figure 4: Osteoporosis as an example of a geriatric syndrome with multifactorial aetiologies for bone ageing.

result of a remodelling imbalance, where an uncoupling of bone formation to bone resorption causes a net loss of bone tissue. There are at least three major contributions to age-related bone loss: secondary hyperparathyroidism due to vitamin D deficiency and decreased renal calcium absorption, impaired bone formation and osteocyte function and, in both women and men, lower oestrogen levels.

Cell senescence in the bone micro-environment, declines in IGF-I levels, and decreased physical activity and response to mechanical loading contribute to poor bone formation. In the setting of unimpaired osteoclast function, there is a relative increase in bone resorption. With the decline in oestrogen, two key phenomena occur which contributes to bone loss; there is an increase in the activation frequency of osteoclasts (i.e., initiation of local resorption events) and there are alterations in local growth factors and cytokines favouring an inflammatory milieu with inhibitory effects on bone formation. There are also random remodelling errors that may increase with ageing.

RESILIENCE TO PHYSIOLOGIC AGEING

It is well established that psychosocial factors influence resilience to agerelated social and behavioural stressors, but there is a limited understanding of human resilience in responses to physiologic or pathologic stressors (*Hadley* et al., 2017). It is clear, however, that low levels of resilience confer vulnerability to stressors (Figure 5). Specific physiologic resilient responses differ depending on both the stressor being exerted and the clinical or physiologic property to be maintained or restored.

There are growing examples of



Figure 5: Relationships among stressors, aging, and resilience to adverse physiologic consequences.

resilient responses that are mediated by regulatory factors, which mitigate potential adverse consequences. Falls are common stressors in older individuals and can result in fractures, especially of the hip or wrist. Bone mineral density remains a very strong predictor of hip fracture up to 25 years after the measurement and preservation of femoral neck bone mass, even in individuals greater than age 75, does occur and minimizes the likelihood of fracture (*Black* et al., 2018).

Neuropathologic changes of Alzheimer disease (AD) uncommonly exist in isolation in the brains of older individuals, but are more often variably combined, or comorbid, with other

lesions that underlie the dementia syndrome including vascular brain injury (VBI), Lewy body disease (LBD), and hippocampal sclerosis (HS). The frequency and severity of these neuropathologic changes in cognitively intact older individuals varies, with some showing a lesion burden considered sufficient evidence for dementia, a situation commonly referred to as apparent cognitive resilience. Here resilience to neuropathological stressors appears to be dependent on the presence of HS only, since cognitively intact individuals can have evidence for high lesion burden consistent with AD, VBI, and/or LBD at brain autopsy (Latimer et al., 2017).

COMMON MECHANISTIC APPROACH

There are common mechanisms thought to underlie physiologic ageing

based on evidence in lower animal models which are now being confirmed

in non-human primates and in humans. These include chronic, low-grade, nonmicrobial inflammation, cellular senesaccumulation cence, of damaged macromolecules (DNA, proteins, carbohydrates, lipids), as well as stem and progenitor cell dysfunction (Kirkland, 2016).

In animal studies, targeting senescent cells using genetic or pharmacological approaches delays, prevents, or alleviates multiple age-related phenotypes associated with chronic diseases,

geriatric syndromes, and loss of physiological resilience. In preclinical studies, reduction of senescent cell bursuccessfully ameliorated den has frailty, cardiac dysfunction, vascular hyporeactivity and calcification, diabetes mellitus, liver steatosis, osteoporosis, vertebral disk degeneration, pulmonary fibrosis, and radiation-induced damage. Senolytic agents are being tested in proof-of-concept clinical trials based on these findings (*Kirkland* et al., 2017).

ACKNOWLEDGEMENTS

RJP is supported by the Robert and Arlene Professorship in Geriatric Medicine at the Mayo Clinic, Rochester, MN, USA.

LITERATURE

- Agarwal, S. and Busse, P.J.: Innate and adaptive immunosenescence. Ann. Allergy Asthma Immunol. 104, 183-190; quiz 190-182 (2010).
- Arai, Y., Martin-Ruiz, C.M., Takayama, M., Abe, Y., Takebayashi, T., Koyasu, S., Suematsu, M., Hirose, N., and von Zglinicki, T.: Inflammation, But Not Telomere Length, Predicts Successful Ageing at Extreme Old Age: A Longitudinal Study of Semi-supercentenarians. EBioMedicine 2, 1549-1558 (2015).
- Bitar, K.N. and Patil, S.B.: Aging and gastrointestinal smooth muscle. Mech. Ageing Dev. 125, 907-910 (2004).
- Black, D.M., Cauley, J.A., Wagman, R., Ensrud, K., Fink, H.A., Hillier, T.A., Lui, L.Y., Cummings, S.R., Schousboe, J.T., and Napoli, N.: The Ability of a Single BMD and Fracture History Assessment to Predict Fracture Over 25 Years in Postmenopausal Women: The Study of Osteoporotic Fractures. J. Bone Miner. Re.s 33, 389-395 (2018).
- Boyce, J.M. and Shone, G.R.: Effects of ageing on smell and taste. Postgrad. Med. J. 82,

239-241 (2006).

- Boyle, P.A., Buchman, A.S., Barnes, L.L., and Bennett, D.A.: Effect of a purpose in life on risk of incident Alzheimer disease and mild cognitive impairment in community-dwelling older persons. Arch. Gen. Psychiatry 67, 304-310 (2010a).
- Boyle, P.A., Buchman, A.S., and Bennett, D.A.: Purpose in life is associated with a reduced risk of incident disability among community-dwelling older persons. Am. J. Geriatr. Psychiatry 18, 1093-1102 (2010b).
- Bulow, R., Simon, P., Thiel, R., Thamm, P., Messner, P., Lerch, M.M., Mayerle, J., Volzke, H., Hosten, N., and Kuhn, J.P.: Anatomic variants of the pancreatic duct and their clinical relevance: an MR-guided study in the general population. Eur. Radiol. 24, 3142-3149 (2014).
- Capobianco, G., Donolo, E., Borghero, G., Dessole, F., Cherchi, P.L., and Dessole, S.: Effects of intravaginal estriol and pelvic floor rehabilitation on urogenital aging in postmenopausal women. Arch. Gynecol. Obstet. 285, 397-403 (2012).
- Chan, E.D. and Welsh, C.H.: Geriatric respira-

tory medicine. Chest 114, 1704-1733 (1998).

- Chandra, A., Rosenzweig, A., and Pignolo, R.J.: Osteobiology of Aging. In: Fractures in the Elderly (Eds.: Pignolo, R.J., Forciea, M.A., and Johnson, J.C.). 2nd Edition. Humana-Springer, New York, 3-37 (2018).
- Commane, D.M., Arasaradnam, R.P., Mills, S., Mathers, J.C., and Bradburn, M.: Diet, ageing and genetic factors in the pathogenesis of diverticular disease. World J. Gastroenterol. 15, 2479-2488 (2009).
- Comparato, G., Pilotto, A., Franze, A., Franceschi, M., and Di Mario, F.: Diverticular disease in the elderly. Dig. Dis. 25, 151-159 (2007).
- Custer, R.P. and Ahlfeldt, F.E.: Studies on the structure and function of bone marrow. Part II. J. Lab. Clin. Med. 17, 960-962 (1932).
- Dorszewska, J.: Cell biology of normal brain aging: synaptic plasticity-cell death. Aging Clin. Exp. Res. 25, 25-34 (2013).
- Elbadawi, A., Diokno, A.C., and Millard, R.J.: The aging bladder: morphology and urodynamics. World J. Urol. 16, Suppl. 1, S10-34 (1998).
- Esposito, C., Plati, A., Mazzullo, T., Fasoli, G., De Mauri, A., Grosjean, F., Mangione, F., Castoldi, F., Serpieri, N., Cornacchia, F., and Dal Canton, A.: Renal function and functional reserve in healthy elderly individuals. J. Nephrol. 20, 617-625 (2007).
- Faulkner, J.A., Larkin, L.M., Claflin, D.R., and Brooks, S.V.: Age-related changes in the structure and function of skeletal muscles. Clin. Exp. Pharmacol. Physiol. 34, 1091-1096 (2007).
- Feldman, M., Cryer, B., McArthur, K.E., Huet, B.A., and Lee, E.: Effects of aging and gastritis on gastric acid and pepsin secretion in humans: a prospective study. Gastroenterology 110, 1043-1052 (1996).
- Ferrari, A.U., Radaelli, A., and Centola, M.: Invited review: aging and the cardiovascular system. J. Appl. Physiol. (1985) 95, 2591-2597 (2003).
- Fjell, A.M., McEvoy, L., Holland, D., Dale, A.M., Walhovd, K.B., and Alzheimer's Disease Neuroimaging Initiative: What is nor-

mal in normal aging? Effects of aging, amyloid and Alzheimer's disease on the cerebral cortex and the hippocampus. Prog. Neurobiol. 117, 20-40 (2014).

- Fleg, J.L. and Kennedy, H.L.: Long-term prognostic significance of ambulatory electrocardiographic findings in apparently healthy subjects greater than or equal to 60 years of age. Am. J. Cardiol. 70, 748-751 (1992).
- Franchini, M.: Hemostasis and aging. Crit. Rev. Oncol. Hematol. 60, 144-151 (2006).
- Frederick, M.G., Ott, D.J., Grishaw, E.K., Gelfand, D.W., and Chen, M.Y.: Functional abnormalities of the pharynx: a prospective analysis of radiographic abnormalities relative to age and symptoms. AJR Am. J. Roentgenol. 166, 353-357 (1996).
- Fu, A. and Nair, K.S.: Age effect on fibrinogen and albumin synthesis in humans. Am. J. Physiol. 275, E1023-1030 (1998).
- Fulp, S.R., Dalton, C.B., Castell, J.A., and Castell, D.O.: Aging-related alterations in human upper esophageal sphincter function. Am. J. Gastroenterol. 85, 1569-1572 (1990).
- Gambert, S.R., and Escher, J.E.: Atypical presentation of endocrine disorders in the elderly. Geriatrics 43, 69-71, 76-78 (1988).
- Glassock, R.J., and Rule, A.D.: The implications of anatomical and functional changes of the aging kidney: with an emphasis on the glomeruli. Kidney Int. 82, 270-277 (2012).
- Guslandi, M., Pellegrini, A., and Sorghi, M.: Gastric mucosal defences in the elderly. Gerontology 45, 206-208 (1999).
- Hadley, E.C., Kuchel, G.A., Newman, A.B., Workshop, S., and Participants. Report: NIA Workshop on Measures of Physiologic Resiliencies in Human Aging. J. Gerontol. A Biol. Sci. Med. Sci. 72, 980-990 (2017).
- Hall, K.E., Proctor, D.D., Fisher, L., and Rose, S.: American gastroenterological association future trends committee report: effects of aging of the population on gastroenterology practice, education, and research. Gastroenterology 129, 1305-1338 (2005).
- Harris, I.D., Fronczak, C., Roth, L., and Meacham, R.B.: Fertility and the aging

male. Rev. Urol. 13, e184-190 (2011).

- Haruma, K., Kamada, T., Kawaguchi, H., Okamoto, S., Yoshihara, M., Sumii, K., Inoue, M., Kishimoto, S., Kajiyama, G., and Miyoshi, A.: Effect of age and Helicobacter pylori infection on gastric acid secretion. J. Gastroenterol. Hepatol. 15, 277-283 (2000).
- Hilton, D., Iman, N., Burke, G.J., Moore, A., O'Mara, G., Signorini, D., Lyons, D., Banerjee, A.K., and Clinch, D.: Absence of abdominal pain in older persons with endoscopic ulcers: a prospective study. Am. J. Gastroenterol. 96, 380-384 (2001).
- Hofman, M.A., and Swaab, D.F.: Living by the clock: the circadian pacemaker in older people. Ageing Res. Rev. 5, 33-51 (2006).
- Hollenberg, N.K., Adams, D.F., Solomon, H.S., Rashid, A., Abrams, H.L., and Merrill, J.P.: Senescence and the renal vasculature in normal man. Circ. Res. 34, 309-316 (1974).
- Holt-Lunstad, J., Smith, T.B., and Layton, J.B.: Social relationships and mortality risk: a meta-analytic review. PLoS Med 7, e1000316 (2010).
- Howarth, A. and Shone, G.R.: Ageing and the auditory system. Postgrad. Med. J. 82, 166-171 (2006).
- Hurwitz, A., Brady, D.A., Schaal, S.E., Samloff, I.M., Dedon, J., and Ruhl, C.E.: Gastric acidity in older adults. JAMA 278, 659-662 (1997).
- Inouye, S.K., Studenski, S., Tinetti, M.E., and Kuchel, G.A.: Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. J. Am. Geriatr. Soc. 55, 780-791 (2007).
- Janssens, J.P.: Aging of the respiratory system: impact on pulmonary function tests and adaptation to exertion. Clin. Chest Med. 26, 469-484 (2005).
- Jones, C.M., and Boelaert, K.: The Endocrinology of Ageing: A Mini-Review. Gerontology 61, 291-300 (2015).
- Kekki, M., Samloff, I.M., Ihamaki, T., Varis, K., and Siurala, M.: Age- and sex-related behaviour of gastric acid secretion at the population level. Scand. J. Gastroenterol. 17, 737-743 (1982).

- Kingsberg, S.A.: The impact of aging on sexual function in women and their partners. Arch. Sex. Behav. 31, 431-437 (2002).
- Kirkland, J.L.: Translating the Science of Aging into Therapeutic Interventions. Cold Spring Harb. Perspect. Med. 6, a025908 (2016).
- Kirkland, J.L., Tchkonia, T., Zhu, Y., Niedernhofer, L.J., and Robbins, P.D.: The Clinical Potential of Senolytic Drugs. J. Am. Geriatr. Soc. 65, 2297-2301 (2017).
- Kitzman, D.W., Scholz, D.G., Hagen, P.T., Ilstrup, D.M., and Edwards, W.D.: Agerelated changes in normal human hearts during the first 10 decades of life. Part II (Maturity): A quantitative anatomic study of 765 specimens from subjects 20 to 99 years old. Mayo Clin. Proc. 63, 137-146 (1988).
- Klostermann, E.C., Braskie, M.N., Landau, S.M., O'Neil, J.P., and Jagust, W.J.: Dopamine and frontostriatal networks in cognitive aging. Neurobiol. Aging 33, 623.e15-623.e24 (2012).
- Kricun, M.E.: Red-yellow marrow conversion: its effect on the location of some solitary bone lesions. Skeletal Radiol. 14, 10-19 (1985).
- Kuk, J.L., Saunders, T.J., Davidson, L.E., and Ross, R.: Age-related changes in total and regional fat distribution. Ageing Res. Rev. 8, 339-348 (2009).
- Lamberts, S.W., van den Beld, A.W., and van der Lely, A.J.: The endocrinology of aging. Science 278, 419-424 (1997).
- Lasch, H., Castell, D.O., and Castell, J.A.: Evidence for diminished visceral pain with aging: studies using graded intraesophageal balloon distension. Am. J. Physiol. 272, G1-3 (1997).
- Latimer, C.S., Keene, C.D., Flanagan, M.E., Hemmy, L.S., Lim, K.O., White, L.R., Montine, K.S., and Montine, T.J.: Resistance to Alzheimer Disease Neuropathologic Changes and Apparent Cognitive Resilience in the Nun and Honolulu-Asia Aging Studies. J. Neuropathol. Exp. Neurol. 76, 458-466 (2017).
- Liem, A.T., Keunen, J.E., van Norren, D., and

van de Kraats, J.: Rod densitometry in the aging human eye. Invest. Ophthalmol. Vis. Sci. 32, 2676-2682 (1991).

- Lindau, S.T. and Gavrilova, N.: Sex, health, and years of sexually active life gained due to good health: evidence from two US population based cross sectional surveys of ageing. BMJ 340, c810 (2010).
- Lindeman, R.D., Tobin, J., and Shock, N.W.: Longitudinal studies on the rate of decline in renal function with age. J. Am. Geriatr. Soc. 33, 278-285 (1985).
- Lipschitz, D.A., Udupa, K.B., Milton, K.Y., and Thompson, C.O.: Effect of age on hematopoiesis in man. Blood 63, 502-509 (1984).
- Locquet, M., Beaudart, C., Reginster, J.Y., Petermans, J., and Bruyere, O.: Comparison of the performance of five screening methods for sarcopenia. Clin. Epidemiol. 10, 71-82 (2018).
- Marshall, B. J.: Helicobacter pylori. Am. J. Gastroenterol. 89, S116-128 (1994).
- Mitchell, G.F.: Effects of central arterial aging on the structure and function of the peripheral vasculature: implications for end-organ damage. J. Appl. Physiol. (1985) 105, 1652-1660 (2008).
- Montagna, W. and Carlisle, K.: Structural changes in ageing skin. Br. J. Dermatol. 122, Suppl. 35, 61-70 (1990).
- Narici, M.V. and Maffulli, N.: Sarcopenia: characteristics, mechanisms and functional significance. Br. Med. Bull. 95, 139-159 (2010).
- Olivetti, G., Melissari, M., Capasso, J.M., and Anversa, P.: Cardiomyopathy of the aging human heart. Myocyte loss and reactive cellular hypertrophy. Circ. Res. 68, 1560-1568 (1991).
- Panda, A., Arjona, A., Sapey, E., Bai, F., Fikrig, E., Montgomery, R.R., Lord, J.M., and Shaw, A.C.: Human innate immunosenescence: causes and consequences for immunity in old age. Trends Immunol. 30, 325-333 (2009).
- Park, D.C., Lodi-Smith, J., Drew, L., Haber, S., Hebrank, A., Bischof, G.N., and Aamodt, W.: The impact of sustained engagement on

cognitive function in older adults: the Synapse Project. Psychol. Sci. 25, 103-112 (2014).

- Parlesak, A., Klein, B., Schecher, K., Bode, J.C., and Bode, C.: Prevalence of small bowel bacterial overgrowth and its association with nutrition intake in nonhospitalized older adults. J. Am. Geriatr. Soc. 51, 768-773 (2003).
- Pinto, A., De Filippi, R., Frigeri, F., Corazzelli, G., and Normanno, N.: Aging and the hemopoietic system. Crit. Rev. Oncol. Hematol. 48, S3-S12 (2003).
- Ramos-Casals, M., Brito-Zeron, P., Lopez-Soto, A., and Font, J.: Systemic autoimmune diseases in elderly patients: atypical presentation and association with neoplasia. Autoimmun. Rev. 3, 376-382 (2004).
- Ray, D. and Yung, R.: Immune senescence, epigenetics and autoimmunity. Clin. Immunol., In press (2018).
- Reinders, I., Murphy, R.A., Koster, A., Brouwer, I.A., Visser, M., Garcia, M.E., Launer, L.J., Siggeirsdottir, K., Eiriksdottir, G., Jonsson, P.V., Gudnason, V., and Harris, T.B.: Muscle Quality and Muscle Fat Infiltration in Relation to Incident Mobility Disability and Gait Speed Decline: the Age, Gene/Environment Susceptibility-Reykjavik Study. J. Gerontol. A Biol. Sci. Med. Sci. 70, 1030-1036 (2015).
- Salles, N.: Basic mechanisms of the aging gastrointestinal tract. Dig. Dis. 25, 112-117 (2007).
- Salvi, S.M., Akhtar, S., and Currie, Z.: Ageing changes in the eye. Postgrad. Med. J. 82, 581-587 (2006).
- Schmucker, D.L.: Age-related changes in liver structure and function: Implications for disease? Exp. Gerontol. 40, 650-659 (2005).
- Seftel, A.D.: From aspiration to achievement: assessment and noninvasive treatment of erectile dysfunction in aging men. J. Am. Geriatr. Soc. 53, 119-130 (2005).
- Smith, C.H., Boland, B., Daureeawoo, Y., Donaldson, E., Small, K., and Tuomainen, J.: Effect of aging on stimulated salivary flow in adults. J. Am. Geriatr. Soc. 61, 805-808 (2013).

- Sotaniemi, E.A., Arranto, A.J., Pelkonen, O., and Pasanen, M.: Age and cytochrome P450-linked drug metabolism in humans: an analysis of 226 subjects with equal histopathologic conditions. Clin. Pharmacol. Ther. 61, 331-339 (1997).
- Stam, H., Hrachovina, V., Stijnen, T., and Versprille, A.: Diffusing capacity dependent on lung volume and age in normal subjects. J. Appl. Physiol. (1985) 76, 2356-2363 (1994).
- Stout, M.B., Justice, J.N., Nicklas, B.J., and Kirkland, J.L.: Physiological Aging: Links Among Adipose Tissue Dysfunction, Diabetes, and Frailty. Physiology (Bethesda) 32, 9-19 (2017).
- Strenk, S.A., Strenk, L.M., and Koretz, J.F.: The mechanism of presbyopia. Prog. Retin. Eye Res. 24, 379-393 (2005).
- Tarlatzis, B.C. and Zepiridis, L.: Perimenopausal conception. Ann. NY Acad. Sci. 997, 93-104 (2003).
- Tarnawski, A.S., Ahluwalia, A., and Jones, M.K.: Increased susceptibility of aging gastric mucosa to injury: the mechanisms and clinical implications. World J. Gastroenterol. 20, 4467-4482 (2014).
- Tran, L. and Greenwood-Van Meerveld, B.: Age-associated remodeling of the intestinal epithelial barrier. J. Gerontol. A Biol. Sci. Med. Sci. 68, 1045-1056 (2013).
- Uitto, J.: The role of elastin and collagen in cutaneous aging: intrinsic aging versus photoexposure. J. Drugs Dermatol. 7, s12-

16 (2008).

- Valdivieso, V., Palma, R., Wunkhaus, R., Antezana, C., Severin, C., and Contreras, A.: Effect of aging on biliary lipid composition and bile acid metabolism in normal Chilean women. Gastroenterology 74, 871-874 (1978).
- van der Zee, E.A.: Synapses, spines and kinases in mammalian learning and memory, and the impact of aging. Neurosci. Biobehav. Rev. 50, 77-85 (2015).
- Van Haeringen, N.J.: Aging and the lacrimal system. Br. J. Ophthalmol. 81, 824-826 (1997).
- Veldhuis, J.D.: Altered pulsatile and coordinate secretion of pituitary hormones in aging: evidence of feedback disruption. Aging (Milano) 9, 19-20 (1997).
- Wade, P.R., and Hornby, P.J.: Age-related neurodegenerative changes and how they affect the gut. Sci. Aging Knowledge Environ. 2005, pe8 (2005).
- Wilson, R.S., Beckett, L.A., Barnes, L.L., Schneider, J.A., Bach, J., Evans, D.A., and Bennett, D.A.: Individual differences in rates of change in cognitive abilities of older persons. Psychol. Aging 17, 179-193 (2002).
- Yaar, M. and Gilchrest, B.A.: Skin aging: postulated mechanisms and consequent changes in structure and function. Clin. Geriatr. Med. 17, 617-630 (2001).

AGEING AND THE MICROBIOME: AN AVENUE FOR INTERVENTION

BRIAN K. KENNEDY

Departments of Biochemistry and Physiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore; Centre for Healthy Ageing, National University Health System, Singapore, Singapore Institute for Clinical Sciences, Singapore; Buck Institute for Research on Aging, Novato, CA, USA

SUMMARY

Progress in ageing research has been impressive in recent years, leading to an increased understanding of the biological determinants of ageing and the identification of a number of interventions that might slow ageing in humans. Given that ageing is the largest risk factor for the chronic diseases that are increasing in prevalence globally, this raises the possibility that through the use of these interventions it will be feasible to slow ageing and prevent the onset of many diseases simultaneously. However, ageing is not generally considered as a disease and any intervention to slow ageing will necessitate a high safety profile to be used in healthy individuals. For this reason, it is important to consider non-pharmaceutical interventions, and modification of the microbiome is a promising avenue for exploration. Here I discuss how manipulation of the microbiome may impact ageing, highlighting data from animal models and human studies.

INTRODUCTION

Demographics are rapidly changing, leading to an increasing percentage of the population over the age of 65. In some countries that percentage will approach 40% in the near future. With ageing comes functional decline and disease onset. Most individuals over the age of 65 have one or more chronic diseases (e.g. neurologic, cardiovascular and metabolic syndromes, as well as cancers). While each of these diseases have identified risk factors, they pale in comparison to the risks associated with ageing itself (Kennedy et al., 2014). Targeting risks factors has proven effective in disease prevention; lowering cholesterol and glucose reduces onset and progression of cardiovascular and

metabolic disease, respectively. What would happen if the biggest risk factor, ageing, was targeted?

Efforts to understand the molecular determinants of ageing have been underway for nearly a century and great progress has been made in the last few decades, largely through the use of non-vertebrate models such as yeast, worms and flies. Prior to these studies, many researchers proposed that ageing, due to its complexity, was highly difficult to modulate. Working with these organisms, however, researchers identified hundreds of genes that impact ageing and proposed several mechanisms to explain their action (*Longo* et al., 2012; *Uno* and *Nishida*, 2016; *Piper*

and *Partridge*, 2018). A surprising finding was that ageing pathways and mechanisms are likely conserved between these organisms, even though they are often quite divergent from an evolutionary perspective (*Smith* et al., 2008). Further extending this observation, more recent reports indicate that many of these same genes impact mammalian, and even possibly human, ageing (*Tian* et al., 2017).

Calorie (or dietary) restriction was shown to slow ageing in mice and rats over eight decades ago, and remains a hot topic for research today (*Balasubramanian* et al., 2017). The pathways modulated by nutrient limitation, including insulin/IGF and mTOR signalling, also modulate ageing and recent evidence has suggested that reducing mTOR activity may impact aspects of human ageing (*Mannick* et al., 2014). Interestingly, these interventions have been shown to alter the microbiome as well, raising the question as to whether their effects are modulated through the gut microbiota.

Here links between the microbiome and ageing are discussed using three short perspectives, with a focus on the gut. First, I detail changes in the gut microbiome that occur with human ageing, findings that might point to targeted modulation. Second, I outline the links between known interventions affecting ageing and alterations in the microbiome. Finally, I discuss the prospects of using targeted interventions to modulate microbiome composition as a means of slowing ageing. The intersection of two major fields of study, microbiome and ageing, offers great promise for novel discoveries and more importantly an avenue toward extending human healthspan, the disease free and highly functional period of life.

MICROBIOME ALTERATIONS DURING AGEING

Perhaps not surprisingly, there are extensive microbiome changes during the ageing process (Kundu et al., 2017). Diversity between individuals also increases with ageing (Claesson et al., 2011). One surprising finding was that the microbiome of residential dwelling elders was significantly different from those in long-term residential care (*Claesson* et al., 2011). Community dwelling elders have a more diverse microbiome, correlated with a diet higher in fibre and reduced levels of age-associated inflammatory factors, such as IL-6, TNF- α and C-reactive protein. Metagenomic analysis further indicated higher metabolism of butyrate and short-chain fatty acids in community dwellers, who were also healthier by several parameters. Microbiome changes are also associated with several diseases of ageing, including neurodegenerative conditions (*Kundu* et al., 2017). Finally, a recent study suggests that semi-supercentenarians (105-109 years) have a microbiome associated with health promotion, for instance enriched for taxa such as *Bifidobacterium*, *Christensenellaceae*, and *Akkermansia* (*Biagi* et al., 2010). None of these studies demonstrate a causal role for the microbiome in mediating ageing, they certainly raise that possibility. How easy it will be to modify the microbiome of elders, however, remains unclear.

Microbiome-related changes are also evident in ageing mice, with notable similarities and differences. One genus that declines in both human and mice is *Akkermansia*, which is associated with protection from a range of diseases (*Langille* et al., 2014). Germ-free mice are also a major tool for research and a recent study suggests that these mice have enhanced lifespan (Thevaranjan et al., 2017), similar in some respects to that of C. elegans exposed to dead bacteria as a food source (Thevaranjan et al., 2017) or adult worms on plates of deprived bacteria entirely (Kaeberlein et al., 2006). Germ-free mice also experienced lower inflammation, which was increased when these mice were co-housed with old conventional animals (*Thevaranjan* et al., 2017). Numerous other changes have been detected, although some discrepancies exist between different mouse studies. By continuing to exploit the mouse model to understand the relationship between microbiome composition, ageing, and disease, it is almost certain that insights will be gained regarding human ageing and interventional strategies developed.

AGEING INTERVENTIONS AND THE MICROBIOME

A major step forward in ageing research has been the identification of dietary and small molecule-based interventions that slow ageing and extend healthspan in animal models. Among the most prominent of these are calorie restriction, metformin and rapamycin, all of which have been linked to changes in the microbiome. An advantage of the animal models is that it is easier to study the relationship between interventions that extend lifespan and microbiome changes. For instance, calorie restriction as expected is associated with significant changes in the microbiome, including an increase in bacterial species associated longer lifespan. Interestingly, both calorie restriction and intermittent fasting, which mimics the effects of calorie restriction, have been reported to extend lifespan in flies and this is associated with a reduced gut bacterial load (*Regan* et al., 2016; Catterson et al., 2018). The effect is more prominent in females, which are more prone to gut deterioration with age. Further analysis of the effects of calorie restriction and fasting will be important as these interventions enhance longevity and promote healthspan across a wide range of species.

Metformin is a widely used drug used to treat hyperglycaemia in the context of a range of metabolic

conditions and has also been reported to extend lifespan in several animal models (Barzilai et al., 2016). Intriguingly, retrospective studies in humans suggest that diabetic patients taking metformin have a lower-thanexpected mortality rate and moreover metformin may be protective for a range of other chronic conditions (Bannister et al., 2014). These findings together have suggested that metformin may deserve even more widespread use to extend human healthspan. A surprising finding regarding metformin's ageing effects came from C. elegans, where it was shown that lifespan extension by the drug required live bacteria as a food source (*Cabreiro* et al., 2013; Heintz and Mair, 2014). Axenic worms or those grown on dead bacteria do not respond to the drug. The mode of action was reported to be an inhibition of folate metabolism in *E. coli*, leading to reduced methionine. This is consistent with lifespan extension by methionine reduction in a variety of animal models, including worms. Several studies suggest that metformin affects the mammalian microbiome (Forslund et al., 2015; Wu et al., 2017; Bauer et al., 2018), but this relates to lifespan extension in mammals remains unknown.

Inhibition of the mTOR pathway is associated with lifespan and healthspan

extension in a range of animal models (*Kennedy* and *Lamming*, 2016), and preliminary data using rapalogs [a class of very specific mTOR inhibitors in which rapamycin is the founder (*Lamming* et al., 2013)] suggests similar effects may be possible in humans (*Mannick* et al., 2014, 2018). Acute rapamycin treatment in old mice impacts the faecal microbiome, with a notable increase in segmented filamentous bacteria. An earlier study in middle-aged mice reported modest effects of rapamycin on gut bacterial composition

(*Hurez* et al., 2015). Whether these changes are important for the longevity effects of rapamycin remains unknown. A number of other drugs have been reported to extend lifespan in mice and they may have affected microbiome composition as well. For instance, acarbose, an α -glucosidase inhibitor that extends lifespan in male mice (*Strong* et al., 2016), alters microbiome and in turn its efficacy as an antidiabetic treatment may be dependent on microbiome composition (*Su* et al., 2015; *Gu* et al., 2017).

MICROBIOME INTERVENTIONS AND AGEING

Drugs are promising interventions for targeting ageing, but as stated this approach is fraught with regulatory hurdles. By contrast, microbiomemediated interventions are likely less subject to regulatory concerns but research must be performed to determine to optimize the efficacy of any such approach (Kundu et al., 2017). Faecal microbiota transplants (FMTs) are perhaps the most promising to date, although the long-term effects of this approach remain controversial. At first blush, it would seem that to delay ageing, any intervention must be durable, or administered relatively frequently. Other approaches, including postbiotics and dietary approaches to modify bacterial populations, are probably even more primitive to date.

In particular, metagenomic studies that identify metabolite deficiencies in the gut of ageing people and corresponding changes in systemic metabolism offer great promise as it may be possible to correct these deficiencies with supplements or other approaches. For instance, healthier community dwelling elders were found to have enriched bacterial species for short chain fatty acid production, which may promote more youthful metabolism (Claesson et al., 2012). It should be noted that molecular changes associated with ageing are not always detrimental, and in some cases might be compensatory. A good example is testosterone, which declines with ageing in the male population. Testosterone supplementation to restore the hormone to youthful levels is not necessarily beneficial, and in fact is associated with increased risk of disease in some contexts (Sansone et al., 2017). Together, these observations call for more intensive studies in animal models and humans to (1) identify candidate microbiome-associated interventions that might extend healthspan and (2) test them effectively.

CONCLUSION

The possibility to slow human ageing and extend healthspan is a relatively recent occurrence, as this has only been relatively recently achieved in animal
models. The potential benefits of preventing multiple chronic diseases simultaneously and improving function later in age demand that this approach be explored. Meanwhile, the importance of gut bacterial species in human health and disease has emerged almost concurrently. To what extent do our residents age us, or keep us young? Merging these two concepts, ageing and microbiome research, offers great promise and requires intense investigation. Possibly we can live longer and healthier by training our residents to behave, or perhaps helping them train us to adopt healthier lifestyles?

LITERATURE

- Balasubramanian, P., Howell, P.R., and Anderson, R.M.:Aging and Caloric Restriction Research: A Biological Perspective With Translational Potential. EBioMedicine 21, 37-44 (2017).
- Bannister, C.A., Holden, S.E., Jenkins-Jones, S., Morgan, C.L., Halcox, J.P., Schernthaner, G., Mukherjee, J., and Currie, C.J.: Can people with type 2 diabetes live longer than those without? A comparison of mortality in people initiated with metformin or sulphonylurea monotherapy and matched, non-diabetic controls. Diabetes Obes. Metab. 16, 1165-1173 (2014).
- Barzilai, N., Crandall, J.P., Kritchevsky, S.B., and Espeland, M.A.: Metformin as a Tool to Target Aging. Cell Metab. 23, 1060-1065 (2016).
- Bauer, P.V., Duca, F.A., Waise, T.M.Z., Rasmussen, B.A., Abraham, M.A., Dranse, H.J., Puri, A., O'Brien, C.A., and Lam, T.K.T.: Metformin Alters Upper Small Intestinal Microbiota that Impact a Glucose-SGLT1-Sensing Glucoregulatory Pathway. Cell Metab. 27, 101-117 (2018).
- Biagi, E., Nylund, L., Candela, M., Ostan, R., Bucci, L., Pini, E., Nikkïla, J., Monti, D., Satokari, R., Franceschi, C., Brigidi, P., and De Vos, W.: Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians. PLoS One 5, e10667 (2010).
- Cabreiro, F., Au, C., Leung, K.Y., Vergara-Irigaray, N., Cochemé, H.M., Noori, T., Weinkove, D., Schuster, E., Greene, N.D., and Gems, D.: Metformin retards aging in C. elegans by altering microbial folate and

methionine metabolism. Cell 153, 228-239 (2013).

- Catterson, J.H., Khericha, M., Dyson, M.C., Vincent, A.J., Callard, R., Haveron, S.M., Rajasingam, A., Ahmad, M., and Partridge, L.: Short-Term, Intermittent Fasting Induces Long-Lasting Gut Health and TOR-Independent Lifespan Extension. Curr. Biol. 28, 1714-1724 (2018).
- Claesson, M.J., Cusack, S., O'Sullivan, O., Greene-Diniz, R., de Weerd, H., Flannery, E., Marchesi, J.R., Falush, D., Dinan, T., Fitzgerald, G., Stanton, C., van Sinderen, D., O'Connor, M., Harnedy, N., O'Connor, K., Henry, C., O'Mahony, D., Fitzgerald, A.P., Shanahan, F., Twomey, C., Hill, C., Ross, R.P., and O'Toole, P.W.: Composition, variability, and temporal stability of the intestinal microbiota of the elderly. Proc. Natl. Acad. Sci. USA 108, Suppl. 1, 4586-4591 (2011).
- Claesson, M.J., Jeffery, I.B., Conde, S., Power, S.E., O'Connor, E.M., Cusack, S., Harris, H.M., Coakley, M., Lakshminarayanan, B., O'Sullivan, O., Fitzgerald, G.F., Deane, J., O'Connor, M., Harnedy, N., O'Connor, K., O'Mahony, D., van Sinderen, D., Wallace, M., Brennan, L., Stanton, C., Marchesi, J.R., Fitzgerald, A.P., Shanahan, F., Hill, C., Ross, R.P., and O'Toole, P.W.: Gut microbiota composition correlates with diet and health in the elderly. Nature 488, 178-184 (2012).
- Forslund, K., Hildebrand, F., Nielsen, T., Falony, G., Le Chatelier, E., Sunagawa, S., Prifti, E., Vieira-Silva, S., Gudmundsdottir, V., Pedersen, H.K., Arumugam, M.,

Kristiansen, K., Voigt, A.Y., Vestergaard, H., Hercog, R., Costea, P., Kultima, J.R., Li, J., Jørgensen, T., Levenez, F., Dore, J.; MetaHIT consortium, Nielsen, H.B., Brunak, S., Raes, J., Hansen, T., Wang, J., Ehrlich, S.D., Bork, P., and Pedersen, O.: Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. Nature 528, 262-266 (2015).

- Gu, Y., Wang, X., Li, J., Zhang, Y., Zhong, H., Liu, R., Zhang, D., Feng, Q., Xie, X., Hong, J., Ren, H., Liu, W., Ma, J., Su, Q., Zhang, H., Yang, J., Wang, X., Zhao, X., Gu, W., Bi, Y., Peng, Y., Xu, X., Xia, H., Li, F., Xu, X., Yang, H., Xu, G., Madsen, L., Kristiansen, K., Ning, G., and Wang W.: Analyses of gut microbiota and plasma bile acids enable stratification of patients for antidiabetic treatment. Nat. Commun. 8,1785 (2017).
- Heintz, C. and Mair, W.: You are what you host: microbiome modulation of the aging process. Cell 156, 408-411 (2014).
- Hurez, V., Dao, V., Liu, A., Pandeswara, S., Gelfond, J., Sun, L., Bergman, M., Orihuela, C.J., Galvan, V., Padrón, Á., Drerup, J., Liu, Y., Hasty, P., Sharp, Z.D., and Curiel, T.J.: Chronic mTOR inhibition in mice with rapamycin alters T, B, myeloid, and innate lymphoid cells and gut flora and prolongs life of immune-deficient mice. Aging Cell. 14, 945-956 (2015).
- Kaeberlein, T.L., Smith, E.D., Tsuchiya, M., Welton, K.L., Thomas, J.H., Fields, S., Kennedy, B.K., and Kaeberlein, M.: Lifespan extension in Caenorhabditis elegans by complete removal of food. Aging Cell 5, 487-494 (2006).
- Kennedy, B.K., Berger, S.L., Brunet, A., Campisi, J., Cuervo, A.M., Epel, E.S., Franceschi, C., Lithgow, G.J., Morimoto, R.I., Pessin, J.E., Rando, T.A., Richardson, A., Schadt, E.E., Wyss-Coray, T., and Sierra, F.: Geroscience: linking aging to chronic disease. Cell 159, 709-713 (2014).
- Kennedy, B.K. and Lamming, D.W.: The Mechanistic Target of Rapamycin: The Grand ConducTOR of Metabolism and Aging. Cell Metab. 23, 990-1003 (2016).

Kundu, P., Blacher, E., Elinav, E., and

Pettersson S.: Our Gut Microbiome: The Evolving Inner Self. Cell 171, 1481-1493 (2017).

- Lamming, D.W., Ye, L., Sabatini, D.M., and Baur, J.A.: Rapalogs and mTOR inhibitors as anti-aging therapeutics. J. Clin. Invest. 123, 980-989 (2013).
- Langille, M.G., Meehan, C.J., Koenig, J.E., Dhanani, A.S., Rose, R.A., Howlett, S.E., and Beiko, R.G.: Microbial shifts in the aging mouse gut. Microbiome 2, 50 (2014).
- Longo, V.D., Shadel, G.S., Kaeberlein, M., and Kennedy, B.: Replicative and chronological aging in Saccharomyces cerevisiae. Cell Metab. 16, 18-31 (2012).
- Mannick, J.B., Del Giudice, G., Lattanzi, M., Valiante, N.M., Praestgaard, J., Huang, B., Lonetto, M.A., Maecker, H.T., Kovarik, J., Carson, S., Glass, D.J., and Klickstein, L.B.: mTOR inhibition improves immune function in the elderly. Sci. Transl. Med. 6, 268ra179 (2014).
- Mannick, J.B., Morris, M., Hockey, H.P., Roma, G., Beibel, M., Kulmatycki, K., Watkins, M., Shavlakadze, T., Zhou, W., Quinn, D., Glass, D.J., and Klickstein, L.B.: TORC1 inhibition enhances immune function and reduces infections in the elderly. Sci. Transl. Med. 10, doi: 10.1126/ scitranslmed.aaq1564 (2018).
- Piper, M.D.W. and Partridge, L.: Drosophila as a model for ageing. Biochim. Biophys. Acta Mol. Basis Dis. 1864, 2707-2717 (2018).
- Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N., Partridge, L.: Sex difference in pathology of the ageing gut mediates the greater response of female lifespan to dietary restriction. Elife 5, e10956 (2016).
- Sansone, A., Sansone, M., Lenzi, A., and Romanelli, F.: Testosterone Replacement Therapy: The Emperor's New Clothes. Rejuvenation Res. 20, 9-14 (2017).
- Smith, E.D., Tsuchiya, M., Fox, L.A., Dang, N., Hu, D., Kerr, E.O., Johnston, E.D., Tchao, B.N., Pak, D.N., Welton, K.L., Promislow, D.E., Thomas, J.H., Kaeberlein, M., and Kennedy, B.K.: Quantitative evidence for conserved longevity pathways

between divergent eukaryotic species. Genome Res. 18, 564-570 (2008).

- Strong, R., Miller, R.A., Antebi, A., Astle, C.M., Bogue, M., Denzel, M.S., Fernandez, E., Flurkey, K., Hamilton, K.L., Lamming, D.W., Javors, M.A., de Magalhães, J.P., Martinez, P.A., McCord, J.M., Miller, B.F., Müller, M., Nelson, J.F., Ndukum, J., Rainger, G.E., Richardson, A., Sabatini, D.M., Salmon, A.B., Simpkins, J.W., Steegenga, W.T., Nadon, N.L., and Harrison, D.E.: Longer lifespan in male mice treated with a weakly estrogenic agonist, an antioxidant, an alpha-glucosidase inhibitor or a Nrf2inducer. Aging Cell 15, 872-884 (2016).
- Su, B., Liu, H., Li, J., Sunli, Y., Liu, B., Liu, D., Zhang, P., and Meng, X.: Acarbose treatment affects the serum levels of inflammatory cytokines and the gut content of bifidobacteria in Chinese patients with type 2 diabetes mellitus. J. Diabetes 7, 729-739 (2015).
- Thevaranjan, N., Puchta, A., Schulz, C., Naidoo, A., Szamosi, J.C., Verschoor, C.P., Loukov, D., Schenck, L.P., Jury, J., Foley,

K.P., Schertzer, J.D., Larché, M.J., Davidson, D.J., Verdú, E.F., Surette, M.G., and Bowdish, D.M.E.: Age-Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage Dysfunction. Cell Host Microbe 21, 455-466 (2017).

- Tian, X., Seluanov, A., and Gorbunova, V.: Molecular Mechanisms Determining Lifespan in Short- and Long-Lived Species. Trends Endocrinol. Metab. 28, 722-734 (2017).
- Uno, M. and Nishida, E.: Lifespan-regulating genes in C. elegans. NPJ Aging Mech. Dis. 2, 16010 (2016).
- Wu, H., Esteve, E., Tremaroli, V., Khan, M.T., Caesar, R., Mannerås-Holm, L., Ståhlman, M., Olsson, L.M., Serino, M., Planas-Fèlix, M., Xifra, G., Mercader, J.M., Torrents, D., Burcelin, R., Ricart, W., Perkins, R., Fernàndez-Real, J.M., and Bäckhed, F.: Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. Nat. Med. 23, 850-858 (2017).

HOST-MICROBIOTA INTERACTIONS DURING AGEING

DARIO RICCARDO VALENZANO

Max Planck Institute for Biology of Ageing, Cologne, Germany

SUMMARY

Host-associated microbial communities take part to several fundamental aspects of host biology, including development, nutrient absorption, immunity and pathogenesis. Throughout host life, microbes associated with external body surfaces vary in composition and function and can shift from commensal to pathogenic. However, whether these microbial communities can directly influence host ageing has remained elusive for a long time. Recent work in laboratory model organisms has revealed for the first time that commensal microbes can actively modulate host ageing and that microbial communities associated with young hosts can have a health-promoting action in middle-age individuals that leads to significant life span extension. These findings suggest new opportunities to identify systemic ageing-modulating mechanisms via the gut microbiota and will help design innovative anti-ageing interventions that could impact human health and ageing-associated diseases.

AGEING AND HOMEOSTASIS

Ageing is associated with pervasive functional decline and overall decrease in physiological fitness that lead to a progressive increase in the risk of disease and death. During ageing, homeostatic processes fail at multiple levels of biological complexity. DNA repair mechanisms become less efficient (Vaidya et al., 2014) and protein homeostasis, which ensures proteome stability, functionally declines at older age (Morimoto and Cuervo, 2009). Membrane and organelle composition, as well as cell physiology, undergoes dramatic changes during ageing. Ageing is typically associated with in-creased risk of uncontrolled cellular proliferation - leading to cancer (*Finkel* et al., 2007) – and to protective proliferation arrest - called cellular senescence (*Childs* et al., 2015). If on the one hand cellular senescence acutely reduces the risk of cancer, on the other

hand it leads to inflammation and higher risk for disease and frailty (*Tchkonia* et al., 2013). Changes in the extracellular matrix composition, which is importantly involved in wound healing, stem cell functionality and longevity in experimental animal models (*Ewald* et al., 2015), additionally contribute to the age-dependent decline in organ and organism performance.

A combination of genetic and nongenetic factors affects the rate at which different organisms age and tune life expectancy in different species. Research done in model organisms has revealed key conserved molecular pathways, including the insulin-IGF1, the mTOR and AMPK pathways, which regulate ageing and life span across several species, from yeast to mammals (*Kenyon* et al., 1993; *Kapahi* et al., 2010; *Lapierre* and *Hansen*, 2012; *Weir* et al., 2017). Environmental manipulations, including temperature changes, different diets and various forms of stress, also significantly impact ageing in several model organisms (*Liu* and *Walford*, 1966; *Weindruch* et al., 1986; *Libert* et al., 2007; *Ermolaeva* et al., 2013). Overall, age-

ing and life span are complex physiological outcomes of the geneticallyencoded strategies that organisms evolved to maintain homeostasis throughout their lifetime in response to the continuous exposure to internal and external stimuli.

GUT MICROBIOTA CONTRIBUTE TO HOST PHYSIOLOGY

Complex microbial communities covering host surfaces occupy the interface between organisms and the external environment, from roots and leaves in plants to skin and mucosal surface including lungs and gut in animals. These microbial communities participate in a wide range of key biological processes, including nutrition, development (Sommer and Backhed, 2013; Hill et al., 2016), essential vitamin synthesis, metabolism (Nicholson et al., 2012), immune modulation (Geva-Zatorsky et al., 2017), defence against pathogens and disease. Host-associated microbiota constitute highly diverse and dynamic communities, able to modulate and buffer host physiology by promptly reacting to internal (e.g. hormonal and metabolic) and external (e.g. temperature, toxins, pathogens) stimuli. A large amount of circulating host metabolites in the host serum are synthesized by gut microbes (Wikoff et al., 2009), further supporting that host-associated microbiota are not passive passengers on host surfaces, but rather an integrative physiological partner that contributes to host homeostasis. Due to their essential contribution to virtually all host biological functions, it is reasonable to hypothesise that commensal microbes could be essential players during host ageing and could be targeted to design interventions aimed at modulating ageing and age-associated conditions.

GUT MICROBIOTA AND IMMUNE FUNCTION: A RECIPROCAL INFLUENCE

Mounting evidence indicates that the gut microbiota play a prominent role in shaping the immune system during development. In humans, new-borns acquire their initial resident gut microbiota from several sources, most prominently from the birth canal and from maternal skin during breast-feeding. These early-acquired bacterial consortia shape the structure of mature gut microbiota (*Dominguez-Bello* et al., 2010) and play an essential role for immunological development (*Round* and *Mazmanian*, 2009). Caesarean sections and formula feeding dramatically affect

the gut microbial composition, and consequently impact proper immune system development, with potentially long-lasting effects for individual health (Bokulich et al., 2016). Gut bacteria interact with cells of both the innate and adaptive immune system in the gut mucosa and different metabolites synthesized by gut bacteria can induce pro or anti-inflammatory host responses (Arpaia and Rudensky, 2014). Epitopes derived from gut bacteria, both commensal and pathogenic, significantly shape the evolution of lymphocyte antibody and receptor repertoires (*Zhao* and *Elson*, 2018). Consistently, specific-pathogen-free (SFP) raised mice have a much higher immunoglobulin (IgA) diversity compared to mono-colonized mice (*Lindner* et al., 2015).

While bacteria shape immune cells development and function, immune cells, in turn, play a key role in determining and controlling the composition and abundance of gut microbial communities. Pharmacologically immunosuppressed rats undergo a loss of commensal gut microbe diversity, which are generally associated with a healthy status (*Bhat* et al., 2017). Specific antibodies, i.e. IgAs, synthesized by intestinal plasma cells, are majorly secreted in the intestinal lumen and bind to a broad subset of bacteria to help maintain a healthy barrier function (*Bunker* et al., 2017). Under normal physiological conditions, IgAs target bacterial taxa that otherwise would contribute to pathogenesis (*Planer* et al., 2016). Conversely, specific pathogens, including influenza virus, succeed in infecting the host by precisely degrading human IgAs.

While bacteria influence immune system evolution, on the other hand the immune system acts as a selective force to shape the composition and function of gut microbes.

GUT MICROBIOTA IN AGEING AND DISEASE

Diet and lifestyle importantly affect differences in microbiota composition among individuals (Rothschild et al., 2018). The composition of host-associated gut microbiota undergoes dramatic changes in humans after birth, becomes stable during adulthood in non-pathological conditions, varies during pregnancy and then goes through drastic changes during ageing (Kostic et al., 2013). Although stable in composition during adulthood, microbiota has rapid functional metabolic oscillations during the day (Thaiss et al., 2016). Across many organisms, including laboratory flies, mice and humans, ageing is characterized by dramatic changes in the composition of the commensal gut microbiota, which could lead to dysbiosis and ultimately host demise (Claesson et al., 2012; Guo et al., 2014; Clark et al., 2015). While gut microbiota associated with healthy hosts are typically characterized by large bacterial taxonomic diversity, frailty and ageing are associated with loss of diversity and expansion of more pathogenic bacterial species. Studies across different human age cohorts have shown that large changes in the abundance of subdominant bacterial taxa in the gut are a hallmark of ageing. Moreover, exceptionally long-lived individuals, including supercentenarians, are characterized by the persistence of bacterial taxa associated with health (Biagi et al., 2016). While diversity-associated microbial taxa often decline during age, specific bacterial taxa, such as Clostridiales, are associated with malnutrition and increased frailty (O'Toole and Jeffery, 2015). In flies, reducing gut microbial dysbiosis by improving immune homeostasis promotes longer life span (Guo et al., 2015). In humans, after antibiotic treatment, pathogenic bacterial species, such as *Clostridium* difficile and Enterococcus faecalis, can restructure the gut microbial composition and cause severe chronic conditions that pose a major threat for public health (Backhed et al., 2012; Milani et al., 2016). In humans, faecal material transfer from healthy donors is successfully used in the clinic to resolve acute *Clostridium difficile* infections (*Lee* et al., 2016). Remarkably, transplanting microbes from obese individuals into



Figure 1: Ageing in the turquoise killifish. Reaching maturity as soon as 3-4 weeks post hatching, killifish display several ageing-related phenotypes by 16 weeks.

germ-free-raised mice leads to dramatic effects, including elevated adiposity and changes in fatty acid and amino acid metabolism, associated with systemic health (*Ridaura* et al., 2013). Overall, microbiota composition and function dramatically change during ageing and disease and manipulating the composition of the gut microbial communities via microbiota transplants has the opportunity to be a novel powerful intervention to impact the ageing process and improve systemic health.

GUT MICROBIOTA PLAY A CAUSAL ROLE IN MODULATING HOST AGEING AND LIFE SPAN

Despite evidence that the gut microbiota could dramatically modulate host metabolism and physiology, until recently it was not clear whether gut microbes could causally modulate host ageing and longevity. Recent work done in nematode worms, flies, fish and mice has shown that gut microbes can beneficially influence host ageing and life span, proving that specific components of this microbial consortium can *de facto* improve overall physiological fitness of the host (*Seidel* and *Valenzano*, 2018).

Work done in nematodes (*Caeno-rhabditis elegans*) has shown that worms feeding on different bacterial species and on varieties of *E. coli* strains different from the standard laboratory OP50 *E. coli* strain (*Girard* et al., 2007) live longer than standard-fed worms (*Sanchez-Blanco* et al.,

2016; Han et al., 2017). A recent study in a short-lived fish has further shown in vertebrates that the gut microbiota is causally involved in modulating host ageing and life span, and that youngassociated gut microbiota, transplanted to middle-age individuals, could lead to life span and health benefits (Smith et al., 2017). This study used the naturally short-lived turquoise killifish (Nothobranchius furzeri), which is the shortest-lived vertebrate raised in a laboratory setting (Figure 1), which includes captive strains with a median life span of about 4 months (Cellerino et al. 2016, Valenzano et al., 2017). For comparison, laboratory mice live 2.5-3 years and zebrafish can live until five years (*Kim* et al., 2016). Despite their short life span, turquoise killifish also display a wide range of age-related changes, including an increased occurrence of cancer, reduced regenerative capacity, increased cellular senescence, neurodegeneration and cognitive decline, making it a powerful new vertebrate model system to study ageing and age-related diseases (*Valenzano* et al., 2017). Turquoise killifish have complex gut microbiota, both in the wild and in captivity, similar in taxonomic diversity to mammals. During ageing, the overall microbial diversity of the resident gut microbiota decreases, while potentially pathogenic *Proteobacteria* become more prevalent, possibly contributing to host demise (*Smith* et al., 2017). After acute re-colonisation of the intestine of middle age individuals with gut microbiota from young donors, fish lived significantly longer, remained more active at old age, and kept highly diverse microbiota. Whether this gut microbial transfer influences immune function and whether its effect is sufficient to improve host health and therefore delay the ageing process is still an open question.

TURQUOISE KILLIFISH AS A MODEL TO STUDY VERTEBRATE IMMUNOSENESCENCE

Unlike invertebrate model organisms such as laboratory flies (Drosophila) and nematode worms (Caenorhabditis elegans), fish are equipped with an adaptive immune system consisting of both T and B-lymphocytes. B-lymphocytes undergo somatic recombination at the IgH (Immunoglobulin heavy chain) locus and generate an extremely diverse (>10⁹) antibody repertoire, which enables to build complex immune responses towards extraneous agents, such as bacteria, viruses, etc. B-cells that bind antigens from microbes and viruses undergo amplification, affinity maturation and unfold a targeted immune response against them in a highly regulated process. Antibody repertoire diversity can be used as a proxy of the function of the whole B-lymphocyte compartment, with high diversity being better than low diversity, and antibody repertoire variation throughout ageing can be adopted as a biomarker of overall health status. Although it is known that upon ageing the B-cell repertoire declines (Martin et al., 2015), it is not clear whether this decline correlates with the decline in the gut microbial diversity. Currently, it is not clear what

are the implications of the age-dependent decrease in immune function (e.g. decreased antibody repertoire diversity) for the changes in microbial diversity and function that occur during ageing. Age-dependent immune dysfunction could lead to proliferation of pathogenic bacteria that are already present in the young gut microbiota, eliciting bacterial community dynamics that favour more proliferative and pathogenic bacterial taxa over commensal and slowreplicating taxa. Alternatively, specific bacterial strains – within a specific subset of bacterial species - could evolve within the gut to escape immune attacks, independently of immune functional decline associated with ageing. Bacterial evolution towards increased pathogenicity could, in turn, lead to host damage and systemic functional decline. As a third alternative, agedependent host immune dysfunction and bacterial evolution could occur simultaneously, leading to increased pathogenicity of gut bacteria and agerelated diseases. The characterization of the sequence of the IgH locus in short-lived vertebrate species, such as the turquoise killifish, will enable to

study in detail whether the immunoglobulin diversity and abundance change during ageing and whether such changes are influenced by anti-ageing interventions, such as transplants of young-associated gut microbiota. Additionally, applying high-throughput approaches to study the immunoglobulin repertoire, such as the recently developed IgSeq method (*Weinstein* et al., 2009), will enable to functionally compare immunoglobulin diversity in the gut with commensal and pathogenic microbial diversity at different ages. High-resolution characterisation of

system and microbiome immune changes during host life will lead to understand whether immune senescence precedes the loss of microbial diversity or whether age-dependent loss of microbial diversity in the gut anticipates immune senescence. The use of a short-lived and experimentally tractable vertebrate model organism, such as the turquoise killifish, will hence be instrumental to answer key questions more rapidly than in longer-lived vertebrate model organisms, such as zebrafish and mice.

LITERATURE

- Arpaia, N. and Rudensky, A.Y.: Microbial metabolites control gut inflammatory responses. Proc. Natl. Acad. Sci. USA 111, 2058-2059 (2014).
- Backhed, F., Fraser, C.M., Ringel, Y., Sanders, M.E., Sartor, R.B., Sherman, P.M., Versalovic, J., Young, V., and Finlay, B.B.: Defining a healthy human gut microbiome: current concepts, future directions, and clinical applications. Cell Host Microbe 12, 611-622 (2012).
- Bhat, M., Pasini, E., Copeland, J., Angeli, M., Husain, S., Kumar, D., Renner, E., Teterina, A., Allard, J., Guttman, D.S., and Humar, A.: Impact of Immunosuppression on the Metagenomic Composition of the Intestinal Microbiome: a Systems Biology Approach to Post-Transplant Diabetes. Sci. Rep. 7, 10277 (2017).
- Biagi, E., Franceschi, C., Rampelli, S., Severgnini, M., Ostan, R., Turroni, S., Consolandi, C., Quercia, S., Scurti, M., Monti, D., Capri, M., Brigidi, P., and Candela, M.: Gut Microbiota and Extreme Longevity. Curr. Biol. 26, 1480-1485 (2016).
- Bokulich, N.A., Chung, J., Battaglia, T., Henderson, N., Jay, M., Li, H., Lieber, A., Wu, F., Perez-Perez, G.I., Chen, Y., Schweizer, W., Zheng, X., Contreras, M.,

Dominguez-Bello, M.G., and Blaser, M.J.: Antibiotics, birth mode, and diet shape microbiome maturation during early life. Sci. Transl. Med. 8, 343ra82 (2016).

- Bunker, J.J., Erickson, S.A., Flynn, T.M., Henry, C., Koval, J.C., Meisel, M., Jabri, B., Antonopoulos, D.A., Wilson, P.C., and Bendelac, A.: Natural polyreactive IgA antibodies coat the intestinal microbiota. Science 358, eaan6619 (2017).
- Childs, B.G., Durik, M., Baker, D.J., and van Deursen, J.M.: Cellular senescence in aging and age-related disease: from mechanisms to therapy. Nat. Med. 21, 1424-1435 (2015).
- Claesson, M.J., Jeffery, I.B., Conde, S., Power, S.E., O'Connor, E.M., Cusack, S., Harris, H.M., Coakley, M., Lakshminarayanan, B., O'Sullivan, O., Fitzgerald, G.F., Deane, J., O'Connor, M.J., Harnedy, N., O'Connor, K.A., O'Mahony, D., van Sinderen, D., Wallace, M., Brennan, L., Stanton, C., Marchesi, J.R., Fitzgerald, A.P., Shanahan, F., Hill, C.R., Ross, R.P., and O'Toole, P.W.: Gut microbiota composition correlates with diet and health in the elderly. Nature 488, 178-184 (2012).
- Clark, R.I., Salazar, A., Yamada, R., Fitz-Gibbon, S., Morselli, M., Alcaraz, J., Rana, A., Rera, M., Pellegrini, M., Ja, W.W., and

Walker, D.W.: Distinct Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal Function and Drive Mortality. Cell Rep. 12, 1656-1667 (2015).

- Ermolaeva, M.A., Segref, A., Dakhovnik, A., Ou, H.L., Schneider, J.I., Utermohlen, O., Hoppe, T., and Schumacher, B.: DNA damage in germ cells induces an innate immune response that triggers systemic stress resistance. Nature 501, 416-420 (2013).
- Ewald, C.Y., Landis, J.N., Porter Abate, J., Murphy, C.T., and Blackwell, T.K.: Dauerindependent insulin/IGF-1-signalling implicates collagen remodelling in longevity. Nature 519, 97-101 (2015).
- Finkel, T., Serrano, M., and Blasco, M.A.: The common biology of cancer and ageing. Nature 448, 767-774 (2007).
- Geva-Zatorsky, N., Sefik, E., Kua, L., Pasman, L., Tan, T.G., Ortiz-Lopez, A., Yanortsang, T.B., Yang, L., Jupp, R., Mathis, D., Benoist, C., and Kasper, D.L.: Mining the Human Gut Microbiota for Immunomodulatory Organisms. Cell 168, 928-943 (2017)..
- Girard, L.R., Fiedler, T.J., Harris, T.W., Carvalho, F., Antoshechkin, I., Han, M., Sternberg, P.W., Stein, L.D., and Chalfie, M.: WormBook: the online review of Caenorhabditis elegans biology. Nucleic Acids Res. 35, D472-475 (2007).
- Guo, L., Karpac, J., Tran, S.L., and Jasper, H.: PGRP-SC2 promotes gut immune homeostasis to limit commensal dysbiosis and extend lifespan. Cell 156, 109-122 (2014).
- Han, B., Sivaramakrishnan, P., Lin, C.J., Neve, I.A.A., He, J., Tay, L.W.R., Sowa, J.N., Sizovs, A., Du, G., Wang, J., Herman, C., and Wang, M.C.: Microbial Genetic Composition Tunes Host Longevity. Cell 169, 1249-1262 (2017).
- Hill, J.H., Franzosa, E.A., Huttenhower, C., and Guillemin, K.: A conserved bacterial protein induces pancreatic beta cell expansion during zebrafish development. Elife 5, e20145 (2016).
- Kapahi, P., Chen, D., Rogers, A.N., Katewa, S.D., Li, P.W., Thomas, E.L., and Kockel, L.: With TOR, less is more: a key role for

the conserved nutrient-sensing TOR pathway in aging. Cell Metab. 11, 453-465 (2010).

- Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtiang, R.: A C-Elegans Mutant That Lives Twice as Long as Wild-Type. Nature 366, 461-464 (1993).
- Kim, Y., Nam, H.G., and Valenzano, D.R.: The short-lived African turquoise killifish: an emerging experimental model for ageing. Dis. Model Mech. 9, 115-129 (2016).
- Kostic, A.D., Howitt, M.R., and Garrett, W.S.: Exploring host-microbiota interactions in animal models and humans. Genes Dev. 27, 701-718 (2013)..
- Lapierre, L.R. and Hansen, M.: Lessons from C. elegans: signaling pathways for longevity. Trends Endocrinol. Metab. 23, 637-644 (2012).
- Lee, C.H., Steiner, T., Petrof, E.O., Smieja, M., Roscoe, D., Nematallah, A., Weese, J.S., Collins, S., Moayyedi, P., Crowther, M., Ropeleski, M.J., Jayaratne, P., Higgins, D., Li, Y., Rau, N.V., and Kim, P.T.: Frozen vs Fresh Fecal Microbiota Transplantation and Clinical Resolution of Diarrhea in Patients With Recurrent Clostridium difficile Infection: A Randomized Clinical Trial. JAMA 315, 142-149 (2016).
- Libert, S., Zwiener, J., Chu, X., Vanvoorhies, W., Roman, G., and Pletcher, S.D.: Regulation of Drosophila life span by olfaction and food-derived odors. Science 315, 1133-1137 (2007).
- Lindner, C., Thomsen, I., Wahl, B., Ugur, M., Sethi, M.K., Friedrichsen, M., Smoczek, A., Ott, S., Baumann, U., Suerbaum, S., Schreiber, S., Bleich, A., Gaboriau-Routhiau, V., Cerf-Bensussan, N., Hazanov, H., Mehr, R., Boysen, P., Rosenstiel, P., and Pabst, O.: Diversification of memory B cells drives the continuous adaptation of secretory antibodies to gut microbiota. Nat. Immunol. 16, 880-888 (2015).
- Liu, K.L. and Walford, R.L.: Increased growth and life-span with lowered ambient temperature in the annual fish, *Cynolebia adloffi*. Nature 212, 1277-1278 (1966).
- Martin, V., Bryan Wu, Y.C., Kipling, D., and

Dunn-Walters, D.: Ageing of the B-cell repertoire. Philos. Trans. R. Soc. Lond. B Biol. Sci. 370, 20140237 (2015).

- Milani, C., Ticinesi, A., Gerritsen, J., Nouvenne, A., Lugli, G.A., Mancabelli, L., Turroni, F., Duranti, S., Mangifesta, M., Viappiani, A., Ferrario, C., Maggio, M., Lauretani, F., De Vos, W., van Sinderen, D., Meschi, T., and Venturab, M.: Gut microbiota composition and Clostridium difficile infection in hospitalized elderly individuals: a metagenomic study. Sci. Rep. 6, 25945 (2016).
- Morimoto, R.I. and Cuervo, A.M.: Protein homeostasis and aging: taking care of proteins from the cradle to the grave. J. Gerontol. A Biol. Sci. Med. Sci. 64, 167-170 (2009).
- Nicholson, J.K., Holmes, E., Kinross, J., Burcelin, R., Gibson, G., Jia, W., and Pettersson, S.: Host-gut microbiota metabolic interactions. Science 336, 1262-1267 (2012).
- O'Toole, P.W. and Jeffery, I.B.: Gut microbiota and aging. Science 350, 1214-1215 (2015).
- Ridaura, V.K., Faith, J.J., Rey, F.E., Cheng, J., Duncan, A.E., Kau, A.L., Griffin, N.W., Lombard, V., Henrissat, B., Bain, J.R., Muehlbauer, M.J., Ilkayeva, O., Semenkovich, C.F., Funai, K., Hayashi, D.K., Lyle, B.J., Martini, M.C., Ursell, L.K., Clemente, J.C., Van Treuren, W., Walters, W.A., Knight, R., Newgard, C.B., Heath, A.C., and Gordon, J.I.: Gut microbiota from twins discordant for obesity modulate metabolism in mice. Science 341, 1241214 (2013)..
- Rothschild, D., Weissbrod, O., Barkan, E., Kurilshikov, A., Korem, T., Zeevi, D., Costea, P.I., Godneva, A., Kalka, I.N., Bar, N., Shilo, S., Lador, D., Vila, A.V., Zmora, N., Pevsner-Fischer, M., Israeli, D., Kosower, N., Malka, G., Wolf, B.C., Avnit-Sagi, T., Lotan-Pompan, M., Weinberger, A., Halpern, Z., Carmi, S., Fu, J., Wijmenga, C., Zhernakova, A., Elinav, E., and Segal, E.: Environment dominates over host genetics in shaping human gut microbiota. Nature 555, 210-215 (2018).
- Sanchez-Blanco, A., Rodriguez-Matellan, A.,

Gonzalez-Paramas, A., Gonzalez-Manzano, S., Kim, S.K., and Mollinedo, F.: Dietary and microbiome factors determine longevity in Caenorhabditis elegans. Aging (Albany NY) 8, 1513-1539 (2016).

- Seidel, J. and Valenzano, D.R.: The role of the gut microbiome during host ageing. F1000Res. 7, F1000 Faculty Rev-1086 (2018).
- Smith, P., Willemsen, D., Popkes, M., Metge, F., Gandiwa, E., Reichard, M., and Valenzano, D.R.: Regulation of life span by the gut microbiota in the short-lived African turquoise killifish. Elife 6, e27014 (2017).
- Sommer, F. and Backhed, F.: The gut microbiota--masters of host development and physiology. Nat. Rev. Microbiol. 11, 227-238 (2013).
- Tchkonia, T., Zhu, Y., van Deursen, J., Campisi, J., and Kirkland, J.L.: Cellular senescence and the senescent secretory phenotype: therapeutic opportunities. J. Clin. Invest. 123, 966-972 (2013).
- Thaiss, C.A., Levy, M., Korem, T., Dohnalova, L., Shapiro, H., Jaitin, D.A., David, E., Winter, D.R., Gury-BenAri, M., Tatirovsky, E., Tuganbaev, T., Federici, S., Zmora, N., Zeevi, D., Dori-Bachash, M., Pevsner-Fischer, M., Kartvelishvily, E., Brandis, A., Harmelin, A., Shibolet, O., Halpern, Z., Honda, K., Amit, I., Segal, E., Elinav, E.: Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations. Cell 167, 1495-1510 (2016).
- Vaidya, A., Mao, Z., Tian, X., Spencer, B., Seluanov, A., and Gorbunova, V.: Knock-in reporter mice demonstrate that DNA repair by non-homologous end joining declines with age. PLoS Genet 10, e1004511 (2014).
- Valenzano, D.R., Aboobaker, A., Seluanov, A., and Gorbunova, V.: Non-canonical aging model systems and why we need them. EMBO J. 36, 959-963 (2017).
- Weindruch, R., Walford, R.L., Fligiel, S., and Guthrie, D.: The retardation of aging in mice by dietary restriction: longevity, cancer, immunity and lifetime energy intake. J. Nutr. 116, 641-654 (1986)..
- Weinstein, J.A., Jiang, N., White, R.A., 3rd,

Fisher, D.S., and Quake, S.R.: Highthroughput sequencing of the zebrafish antibody repertoire. Science 324, 807-810 (2009).

Weir, H.J., Yao, P., Huynh, F.K., Escoubas, C.C., Goncalves, R.L., Burkewitz, K., Laboy, R., Hirschey, M.D., and Mair, W.B.: Dietary Restriction and AMPK Increase Lifespan via Mitochondrial Network and Peroxisome Remodeling. Cell Metab. 26, 884-896 (2017).

- Wikoff, W.R., Anfora, A.T., Liu, J., Schultz, P.G., Lesley, S.A., Peters, E.C., and Siuzdak, G.: Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. Proc. Natl. Acad. Sci. USA 106, 3698-3703 (2009).
- Zhao, Q. and Elson, C.O.: Adaptive immune education by gut microbiota antigens. Immunology 154, 28-37 (2018).

CONTRASTING CONSERVED AND TISSUE-SPECIFIC RESPONSES TO AGEING IN VERTEBRATES

DANIELA ESSER*, LENA BEST*, and CHRISTOPH KALETA

Research Group Medical Systems Biology, Institute for Experimental Medicine, University of Kiel, Kiel, Germany

SUMMARY

Ageing is the most important risk factor for diseases accounting for more than 90% of mortality in western countries. While genomic damage as the root cause of cancer is well established, the primary causes of most degenerative diseases of ageing such as cardiovascular diseases, neurodegenerative diseases and type 2 diabetes are still unknown. In a previous work, we have shown that there is a common core of ageing-associated gene regulation, conserved across vertebrate species and tissues.

In this work, we explored the association between this conserved core signature of ageing and the actual manifestation of ageing in individual tissues in more detail. To this end, we characterized transcriptomic signatures of ageing in individual tissues and investigated the contribution of the microbiome as a potential driver of ageing-associated inflammation. While core versus tissue-specific signatures showed only small differences in phylogenetic conservation or network connectivity, a considerable percentage of genes was found exclusively in tissue-specific signatures. However, on the functional level, tissue-specific signatures showed a very high overlap with the core signature of ageing. Moreover, cancer-regulated genes were enriched among genes of the core signature while they were depleted among the tissue-specific signatures. In contrast, for genes deregulated in degenerative diseases, we observed the tendency of an enrichment amongst genes of the tissue-specific signatures. Through comparisons of ageing signatures to genes regulated in response to microbial colonization, we observed a significant overlap with genes of the core signature of ageing.

Overall these results indicate that tissue ageing on the transcriptomic level is mostly driven by the core signature of ageing and that the microbiome is a potential modulator of this signature.

INTRODUCTION

Ageing is associated with a continuous functional decline and is recognized as a dominant factor driving the pathology of diseases contributing to the majority of mortality in western countries (*Lopez* et al., 2006; *López-Otín* et al.,

2013). There is a remarkable conservation of ageing-associated pathologies such as cancer, cardiovascular disease and cognitive decline across vertebrates (*Dean* et al., 1981; *Genade* et al., 2005; *Pettan-Brewer* and *Treuting*, 2011).

^{*} Both authors contributed equally

However, despite a large number of studies investigating ageing from the molecular (López-Otín et al., 2013) to the population level (Jones et al., 2014), a coherent picture of the processes that are driving ageing and their individual contribution to the pathogenesis of ageing diseases is just emerging (López-Otín et al., 2013). An often-observed feature of ageing is a low-grade inflammatory phenotype ("inflammaging") (Chung et al., 2009) that is characterized by a minor activation of the immune system even in absence of a detectable pathogen (Franceschi and Campisi, 2014). Inflammaging is an important risk factor for morbidity and mortality in the elderly (Franceschi et al., 2000).

A key problem in the study of the ageing phenotype is its remarkable complexity which makes a clear distinction between causal and consequential pathological changes a challenging task (López-Otín et al., 2013; Bousquet et al., 2014; Franceschi and Campisi, 2014; Castellani et al., 2016). In particular, while genomic mutations as a key event in cancer are well established (Valko et al., 2004), the root causes of degenerative ageing diseases such as cardiovascular diseases or neurodegenerative diseases have yet to be identified (Carretero and Oparil, 2000; Galkina and Lev, 2009; Dexter and Jenner, 2013; Reitz and Mayeux, 2014). One contributing factor is that ageing diseases are most often studied in isolation in the particular organ system they affect (Castellani et al., 2016), while extensive comorbidities between ageing diseases (Fried et al., 1999) hint at general underlying mechanisms not restricted to individual organ systems (*Masoro*, 2005; *Bousquet* et al., 2014).

In order to elucidate common pathomechanisms underlying ageing diseases, we recently have analysed a comprehensive next-generation sequencing data set of vertebrate ageing covering four organisms (human, mouse, zebrafish and the killifish Nothobranchius furzeri) across up to four tissues (brain, liver, skin, blood) (Aramillo Irizar et al., 2018). Using a novel method assessing differential activity directly on a functional level, we identified a highly conserved core of ageing-associated transcriptional regulation including an induction of inflamprocesses and a downmatory regulation of cell-cycle as well as developmental processes. While many of these processes have been implicated in ageing pathologies in individual species before (Lee et al., 2000; Teta et al., 2005; Chung et al., 2009), we found that they are part of a conserved core of ageing-associated regulation in the studied vertebrates.

By investigating the association between transcriptional signatures of ageing with differential gene expression observed in ageing related diseases, we found that ageing opposes transcriptional changes of cancer but is aligned with alterations in degenerative disorders. Intriguingly, these results are in line with epidemiological data showing a peak in cancer incidence between 70 to 80 years of age (Harding et al., 2012), while the incidence of degenerative diseases increases up to the oldest age groups (*Berzlanovich* et al., 2005; Harding et al., 2012). Thus, ageing moves the transcriptome away from cancer but brings it closer to that of degenerative diseases. On the functional level, we found that immune- and cell cycle-related processes most strongly contributed to this antagonism, which is supported by their central role in ageing pathologies (Campisi, 2013; Jurk et al., 2014). We observed a similar antagonism on the genetic level whereby for the majority of shared risk loci between cancer and degenerative diseases the allele that predisposed to

cancer protected against degenerative diseases and vice versa (Aramillo Irizar et al., 2018). We found the strongest signals close to central regulators of immune function and cellular senescence such as the P16/Ink4A locus, a master regulator of cellular senescence (Congrains et al., 2013), the Lnk/SH2B3 protein, a key regulator of cytokine signalling (Devallière and Charreau, 2011), and VAMP8 a central regulator of autophagy (Diao and *Liu*, 2016).

Besides ageing-associated changes in the host, ageing and inflammation might also be triggered by changes in microbiome composition in the elderly. Bacterial composition changes continuously throughout lifetime (O'Toole and Jeffery, 2015) and microbial diversity continuously becomes richer in humans with increasing age (Lozupone et al., 2012). Diseases associated with ageing often involve or are preceded by inflammaging, which can be aggravated by changes in the microbial composition (Fransen et al., 2017). A microbiome transfer experiment from old to young germ-free mice led to the upregulation of TNF- α and a dysregulation of pathways involved in the immune Inflammaging response. supporting bacteria were enriched after that treatment, including a higher amount of Akkermansia, TM7 bacteria, and Proteobacteria (Thevaranjan et al., 2018). A similar outcome can be observed when germ-free and old mice are cohoused (Fransen et al., 2017). Common inflammaging biomarkers, such as circulating pro-inflammatory cytokines, cannot be detected in germ-free mice and treating old mice with anti-TNF could alleviate microbiomedriven inflammation (Thevaranjan et al., 2018). In addition, in the ageing microbiome, less bacteria are able to synthesize β -glucuronidases, which are key modulators of epithelial cell

toxicity caused by drugs (*Langille* et al., 2014). Moreover, the amount of monosaccharide-utilizing compared to polysaccharide-using bacteria positively correlates with age in mice (*Langille* et al., 2014).

The immune system of the host can be altered by a modified microbial composition caused by environmental factors including lifestyle and diet. As consequence, a disturbed gut homeostasis can increase the risk for diseases (Langille et al., 2014; Fulde et al., 2018). This is especially the case in elderly humans, in which poor health is associated with imbalances in the microbiome (*Claesson* et al., 2012). Thus, species distribution in the microbial flora is strongly associated with phenotypes of the host such as inflammation, the ability of independent living, sarcopenia as well as geriatric depression. For instance, independently living human individuals are characterized by a higher number of bacteria, which are able to perform biosynthesis of short chain fatty acids (*Claesson* et al., 2012). Taken together, host-microbiome interactions are potential key modulators of health and contribute to the physical conditions of the host.

While we have characterized the conserved transcriptomic signature of ageing in great detail in our previous work, we did not explore the actual manifestation of this signature in individual tissues (Aramillo Irizar et al., 2018). Thus, the aim of this work is to investigate conserved tissue-specific ageing in more detail. We aim to investigate two central hypotheses in ageing research in more detail, 1) whether there is a common mechanism underlying ageing in all tissues, which would point to a common driver of ageing (for instance the microbiome) or 2) whether ageing on the organismal level is basically the sum of functional deterioration in individual tissues. In the first case, we would expect that tissuespecific ageing presents just a tissuespecific manifestation of the changes we observe in the core ageing signature. In the second case, we would expect strong differences in ageing between tissues and the core signature representing the least common denominator. Moreover, we explored the potential role of the microbiome as an important trigger for ageing-associated inflammation and thereby as a potential key driver of the observed transcriptomic signature of ageing.

MATERIAL AND METHODS

Data

To identify sets of conserved ageingassociated genes, we used a similar approach as described in a previous study, in which we searched for commonly regulated processes across species (Aramillo Irizar et al., 2018). More specifically, we used expression values of genes that were orthologue across all the considered four species. We determined orthologues across all four species (zebrafish, killifish, human and mouse) using the R-package orthology. We considered only genes with an orthologue in all four species which had detectable expression across all studied tissues and species (RPKM>0). After removing genes that did not pass the ANOVA-based testing procedure 3748 genes remained [see the Methods section of (Aramillo Irizar et al., 2018) for details]. In order to reduce bias due to different reference gene sets, all analyses described in this work were performed using these 3748 genes as basis. The core ageing signature comprised genes, which were conserved over all tissues and species. Genes with a conserved pattern for all species but only one tissue were assigned to the tissue ageing signatures. The specific ageing signatures contained all genes, which were associated with ageing only for one tissue in one species.

Differentially expressed genes were calculated for 13 cancer types investigated in the consortia 'International Cancer Genome Consortium' (ICGC) (Hudson et al., 2010) and 'The Cancer Genome Atlas' (TCGA). The following tissues were considered: lung (LUADbreast (BRCA-US), US). prostate (PRAD-US), uterus (UCEC-US), kidney (KIRC-US), head and neck (HNSC-US), colorectal (COAD-US), liver (LIHC-US), bladder (BLCA-US), skin (SKCM-US), cervix (CESC-US), pancreas (PAAD-US), and ovary (OV-US). Read counts per gene were downloaded from the ICGC data portal (https://dcc.icgc.org/) (International Cancer Genome Consortium et al., 2010). Sample groups were compared with DeSeq2 v1.8.2. Thereby, gene outliers were replaced with trimmed mean value. All other disease data sets were collected from published studies and processed as described by *Aramillo* Irizar et al. (2018). Genes differentially expressed between germ-free and conventionally raised mice were taken from Pan et al. (2018). Only transcription data of adult mice (12 to 16 weeks of age) were considered in our study.

Enrichment analyses

Gene set enrichment analyses for KEGG pathways and transcription factor binding sites (TFBS) were performed with the online tool innateDB (*Lynn* et al., 2008). To test for significance, a hypergeometric test was applied. Gene set enrichment analyses for Gene Ontology (GO) terms were conducted with the online tool g:profiler (*Reimand* et al., 2016). Exclusively the GO class "Biological processes" was considered. All p-values were corrected for multiple testing using the Benjamini Hochberg approach (*Benjamini* and *Hochberg*, 1995). Protein-protein interaction networks were created with the R package igraph v1.2.1 based on the String database v10 using medium confidence for connection predictions.

Enrichment of ageing signature genes in disease-associated genes

The enrichment score was calculated as [(<number of genes in ageing signature> / <number of disease associated genes>) / (<number of genes in ageing signature> / <number of all investigated genes>)]. Thereby, the regulation direction was not considered.

Calculation of evolutionary conservation per gene

The latest Version (May 8th, 2015) of phylogenetic p-values (phyloP) from

the PHAST software (*Pollard* et al., 2010) for multiple alignments of 99 vertebrate genomes against the human genome (*Siepel* et al., 2005) were obtained from the University of California, Santa Cruz's webserver under the address:

"<u>http://hgdownload.soe.ucsc.edu/golde</u> <u>nPath/hg38/phyloP100way/</u>".

We evaluated phyloP-100-way genetic conservation scores for every single base position of the signatures' genes. The phyloP scores of genes found in any of our ageing signature were averaged over all base-pairs of that gene, leaving us with one mean phyloP score per gene. Chromosome, gene and exon structures were based on the latest version of the human genome (hg38). The mean scores per gene were then compared across signatures.

RESULTS

While we determined commonly differentially regulated processes in our previous work (Aramillo Irizar et al., 2018), we applied this approach directly on the gene expression level to identify genes that show a conserved regulation across four vertebrate species (Homo sapiens, Mus musculus, Danio rerio and Nothobranchius furzeri) and across tissues (blood, brain, liver and skin). Thus, we obtained a core signature of ageing comprising 985 genes and tissue specific signatures comprising 789 genes for blood, 1423 genes for brain, 826 genes for liver and 1281 genes for skin (referred to as tissue signatures). Moreover, we compared those signatures to differentially expressed genes from ageing tissues in mice and humans (referred to as specific signatures) (Aramillo Irizar et al., 2018). Following the two central hypotheses, we investigated the overlap between the core and the tissue signatures of ageing (Figure 1). We found that the signatures of blood and liver were smaller in size than the core signature, while the brain and skin signatures were larger. Tissue signatures showed an average overlap of 38 % with the core genes. Thus, there is a large overlap of the tissue signatures with the overall core signature of ageing. Considering the specific signatures, we found that they showed an average overlap of 61 % with either core or tissue signatures demonstrating that the tissue-specific manifestation of ageing in individual species shows a high concordance with conserved ageing in the corresponding tissue across species (Figure 1).

Conservation and connectivity of the individual ageing signatures

In a first step, we investigated the phylogenetic conservation of genes belonging to the individual signatures. Please



Figure 1: Number of genes involved in each ageing signature. A large proportion of the tissue ageing signature and the specific ageing signature is already enclosed in a more general ageing signature.

note that when considering tissue signatures, we excluded genes also belonging to the core signature. We found no significant differences regarding the phylogenetic conservation between the core signature and the tissue signatures (Figure 2A). Subsequently, we investigated the network context of the genes belonging to the ageing signatures based on their connectivity in the String database (Szklarczyk et al., 2015). Only interactions between genes belonging to one of the ageing signatures were considered. We found that genes belonging to the core signature tended to have fewer connections to other genes than the tissue signatures (Figure 2B-D). This was especially reflected in a higher number of nodes (= genes) with at least 50 different interaction partners, the so-called hub genes, in the tissue ageing signatures (Figure 2C). Also, from the ten proteins with highest connectivity, only one belonged to the core ageing signature (108 interaction partners). However, with 87 % the vast majority of proteins of the core signature remained unconnected.

Functional characterization of core vs. tissue-specific ageing

Next, we functionally characterized the genes belonging to individual tissues. We determined significantly enriched processes as well as enriched transcription factor binding sites in the individual ageing signatures.

Following the results from our previous work, particularly immune related pathways and cell cycle processes were enriched among the different ageing signatures. This is also in agreement with the central role of these



Figure 2: Characteristics of ageing signatures. (A) Genetic conservation of age-linked genes across species. The phyloP scores of genes found in the ageing signatures were averaged over all base-pairs of a gene, based on the human reference (hg38), and compared across signatures. (B) Number of connections within the interaction network based on all ageing-associated genes (core, tissue, and specific signature). Number of connections to all other nodes (independent of the signature) were counted. For better clarity, the y-axis is constrained to a plotting range of 0 % to 10 % and the x-axis to a maximal number of 100 connections (maximal number of connections is 309, three genes had more than 100 connections across all signatures). (C) Number of hub genes (genes with more than 50 connections) within the network described in (B). (D) Average number of connections within the network described in (B).

processes in ageing pathologies (*Chung* et al., 2009; Campisi, 2013). We performed gene set enrichment analyses based on gene ontology (GO) terms using once the complete tissue signatures and once the tissue signatures excluding genes of the core signature. We found the highest number of enriched processes for the core signature even in comparison to the brain and skin signatures which contained more genes (Figure 3A). Importantly, we found that most of the enriched processes in the tissue signatures were also enriched in the core signature. This was true both, when considering tissue signature genes not belonging to the core signature only, and when considering the full gene sets. Thus, a mean of 65.2 % of the functional groups (GO terms) enriched in the tissue signature were also enriched in the core signature when considering only genes specific to the tissue signature and 70.6 % when considering the entire gene set.

Testing for the enrichment of transcription factor binding sites (TFBS), we found an enrichment of transcription factors in the core signature, known to play a central role in the pathogenesis of ageing diseases, such as HIF-1 (involved in ischemic disease,



Figure 3: (A) Functional enrichment analysis using gene ontology (GO) cellular function terms. Number of GO terms enriched for each ageing signature. (B) Transcription factor enrichment analysis. Number of TFBS enriched for each ageing signature.

tumour angiogenesis), E2F-4 DP-2 (involved in tumour suppression, colorectal cancer), and E3F1 DP-2 (involved in tumour suppression). We observed a larger number of enriched transcription factors for the complete versus the reduced tissue signatures (Figure 3B). However, for all tissues except brain the enriched transcription factors showed a strong overlap with the core ageing signature: on average 54.6 % of the transcription factors enriched in the reduced tissue signatures were also enriched in the core signature and on average 58.1 % of transcription factors were enriched in the full gene sets.

Disease-specific characterization of ageing signatures

Since ageing is the most important risk factor for human disease (*Lopez* et al., 2006; *López-Otín* et al., 2013), we determined the enrichment of disease-regulated gene sets with genes belonging to the individual signatures of ageing. Specifically, we investigated to which extent genes, which are differentially expressed in cancer or ageing-

associated degenerative diseases. showed an enrichment among genes belonging to the core or the tissuespecific signatures (Figure 4). While cancer-associated genes showed a significant enrichment among genes forming the core signature of ageing (Fisher's exact test p-value = 0.0461), they showed a significant depletion among the tissue-specific signatures (Fisher's exact test p-value = 0.0187). In contrast, for degenerative diseases the enrichment among genes belonging to the core ageing signature was less pronounced, while we observed a larger number of cases of enrichment among gene sets belonging to the tissuespecific ageing signature. In this context it is also important to emphasize that cancer-associated genes exclusively originated from the affected tissue while deregulated genes from degenerative diseases originated mostly from blood expression data [cf. (Aramillo Irizar et al., 2018) for details]. Thus, the comparison to the blood signature is likely most representative for the degenerative disease signatures.



Figure 4: (A) Enrichment of genes belonging to the individual signatures among genes deregulated in different ageing diseases. (B) Frequency of enrichment of disease-deregulated genes in core and tissue signatures. For the three tissue signatures, average values across the four tissues are shown.

The microbiome is a modulator of the core signature of ageing

Next, we investigated to which extent the different ageing signatures were associated with the host response to microbial colonization. Hence, we compared the signatures to gene expression differences in colon of adult mice possessing a microbiome compared to germ-free mice (*Pan* et al., 2018) (Figure 5). Of the 3748 genes considered for the ageing signatures, 355 showed a differential expression in germ-free versus conventionally raised mice. We found that in particular the core signature of ageing showed an enrichment of

genes differentially expressed during microbial colonization (Fisher's exact test p-value = 0.022) while there was no enrichment of the other signatures. In contrast, the core signature of ageing showed a significant enrichment of microbiome-regulated genes in comparison to brain and liver signatures. Intriguingly, comparing the direction of changes between colonized mice versus germ-free mice with the direction of changes observed in the ageing signature, 68 % of the genes belonging to the core signature showed an opposing direction of regulation during colonization.

DISCUSSION

While we identified a common signature of ageing in our previous work (*Aramillo Irizar* et al., 2018), we here explored the tissue-specific representation of this signature. Specifically, we investigated two partially competing hypotheses in ageing research that 1) ageing is driven by a common underlying process across all tissues or 2) that ageing on the organismal level is



Figure 5: Comparison between ageing signatures and differentially expressed genes in response to microbial colonization. "All" represents the total percentage of all considered genes differentially expressed upon microbial colonization. For each signature the fraction of genes showing the same or opposing direction of regulation are indicated. "Same direction of regulation" corresponds to genes that pointed in the same direction of response in germfree versus colonized mice like in young vs. old mice.

mostly a consequence of functional deteriorations in individual organs.

While we found mostly no differences in phylogenetic conservation between the core and tissue-specific signatures of ageing, the tissue-specific signatures tended to show a stronger degree of interactions. On the functional and the regulatory level tissuespecific signatures showed a very strong overlap to the core signature of ageing: Most processes or transcription factor binding sites enriched for genes belonging to the tissue ageing signatures showed also an enrichment for the genes of the core ageing signature. This supports the hypothesis that ageing in individual tissues probably just represents different tissue-level manifestations of the same underlying processes driving the core ageing signature. On the molecular level, the observed

differences in the tissue-specific signatures compared to the core ageing signature represent probably just tissuespecific downstream consequences of a process' deregulations by the core ageing signature. These downstream consequences could arise, for instance, due to the induction of tissue-specific compensatory processes or due to a reallocation of cellular resources away from tissue-specific processes to processes that need to be induced as part of the core ageing signature.

Testing for enrichment of diseaseregulated genes among the individual signatures revealed that cancerregulated genes tend to show a stronger enrichment among genes belonging to the core signature of ageing compared to the tissue-specific signatures of ageing. In contrast, degenerative diseases of ageing did not show this tendency.

In our previous work we hypothesized that the antagonism between cancer and degenerative diseases might be driven by an accumulation of genomic damage with age (Aramillo Irizar et al., 2018). This accumulation of damage leads to an induction of processes aimed at the suppression of potentially malignant cells which, while to some extent preventing the proliferation of potential cancer cells, drives tissue degeneration and thereby degenerative diseases as a side effect. This hypothesis is well in line with our observation of the enrichment of disease-specific gene sets in the individual ageing signatures: While cancer-deregulated genes as a potential causative factor for the ageing signature are enriched among the core signature of ageing, genes deregulated in degenerative diseases do not show this tendency.

We moreover tested the influence of the microbiome as a potential driver of the core ageing signature. The core ageing signature was significantly enriched for genes differentially expressed between germ-free and conventionally raised mice, whereas no enrichment was found with the tissuespecific signatures of ageing. Importantly, microbial colonization was associated with an expression signature opposing the direction of regulation observed in the core signature of ageing. Microbial colonization seems to lead to a rejuvenation of the colon which is likely explained by a larger colon mass in conventional versus germ-free animals (Wostmann, 1981) thus requiring a stronger proliferation while the core signature of ageing is

associated with an inhibition of cellular proliferation. These observations are in some contrast to previous observations about an extended lifespan of germfree animals (*Thevaranjan* et al., 2018). However, the lifespan-effect is contextdependent. Thus, under caloric restriction germ-free mice show a shortened lifespan (*Tazume* et al., 1991). Moreover, to our knowledge lifespan effects of microbial colonization (i.e. after maturation) have not been tested yet.

Overall, these analyses tend to favour the first hypothesis postulating that ageing is driven by a common process across organs. The tissue-specific manifestation of disease-processes potentially represent downstream consequences of a deregulation of cellular processes in response to the core signature of ageing. Specifically, on a functional and regulatory level the tissuespecific signatures of ageing showed a very high degree of similarity to the core signature of ageing. Moreover, the enrichment of cancer-deregulated genes in the core signature of ageing and a depletion among tissue-specific signatures strongly supports our hypothesis that processes that are actually geared to suppress the proliferation of potential cancer cells are one of the main drivers of ageing-associated regulation conserved across tissues. This is also supported by the enrichment of genes deregulated in degenerative diseases in the tissue-specific signatures, which we suppose arose as a manifestation of the downstream consequences of the core signature in the individual tissues.

ACKNOWLEDGEMENTS

The authors acknowledge support by the Excellence Cluster "Inflammation at Interfaces" (EXC306) and the Collaborative Research Centre "Origin and Function of Metaorganisms" (SFB1182).

LITERATURE

- Aramillo Irizar, P., Schäuble, S., Esser, D., Groth, M., Frahm, C., Priebe, S., Baumgart, M., Hartmann, N., Marthandan, S., Menzel, U., Müller, J., Schmidt, S., Ast, V., Caliebe, A., König, R., Krawczak, M., Ristow, M., Schuster, S., Cellerino, A., Diekmann, S., Englert, C., Hemmerich, P., Sühnel, J., Guthke, R., Witte, O.W., Platzer, M., Ruppin, E., and Kaleta, C.: Transcriptomic alterations during ageing reflect the shift from cancer to degenerative diseases in the elderly. Nat. Commun. 9, 327 (2018).
- Berzlanovich, A.M., Keil, W., Waldhoer, T., Sim, E., Fasching, P., and Fazeny-Dörner, B.: Do centenarians die healthy? An autopsy study. J. Gerontol. A Biol. Sci. Med. Sci. 60, 862-865 (2005).
- Bousquet, J., Jorgensen, C., Dauzat, M., Cesario, A., Camuzat, T., Bourret, R., Best, N., Anto, J.M., Abecassis, F., Aubas, P., Avignon, A., Badin, M., Bedbrook, A., Blain, H., Bourdin, A., Bringer, J., Camu, W., Cayla, G., Costa, D.J., Courtet, P., Cristol, J.-P., Demoly, P., de la Coussaye, J.-E., Fesler, P., Gouzi, F., Gris, J.-C., Guillot, B., Hayot, M., Jeandel, C., Jonquet, O., Journot, L., Lehmann, S., Mathieu, G., Morel, J., Ninot, G., Pelissier, J., Picot, M.-C., Radier-Pontal, F., Robine, J.-M., Rodier, M., Roubille, F., Sultan, A., Wojtusciszyn, A., Auffray, C., Balling, R., Barbara, C., Cambon-Thomsen, A., Chavannes, N.H., Chuchalin, A., Crooks, G., Dedeu, A., Fabbri, L.M., Garcia-Aymerich, J., Hajjam, J., Melo Gomes, E., Palkonen, S., Piette, F., Pison, C., Price, D., Samolinski, B., Schunemann, H.J., Sterk, P.J., Yiallouros, P., Roca, J., Van de Perre, P., and Mercier, J .: Systems medicine approaches for the definition of complex phenotypes in chronic diseases and ageing. From concept to implementation and policies. Curr. Pharm. Des. 20, 5928-5944 (2014).
- Campisi, J.: Aging, cellular senescence, and cancer. Annu. Rev. Physiol. 75, 685-705 (2013).
- Carretero, O.A. and Oparil, S.: Essential Hypertension: part II: treatment. Circulation 101, 446-453 (2000).
- Castellani, G.C., Menichetti, G., Garagnani, P.,

Giulia Bacalini, M., Pirazzini, C., Franceschi, C., Collino, S., Sala, C., Remondini, D., Giampieri, E., Mosca, E., Bersanelli, M., Vitali, S., Valle, I.F. do, Liò, P., and Milanesi, L.: Systems medicine of inflammaging. Brief. Bioinform. 17, 527-540 (2016).

- Chung, H.Y., Cesari, M., Anton, S., Marzetti, E., Giovannini, S., Seo, A.Y., Carter, C., Yu, B.P., and Leeuwenburgh, C.: Molecular inflammation: underpinnings of aging and age-related diseases. Ageing Res. Rev. 8, 18-30 (2009).
- Claesson, M.J., Jeffery, I.B., Conde, S., Power, S.E., O'Connor, E.M., Cusack, S., Harris, H.M.B., Coakley, M., Lakshminarayanan, B., O'Sullivan, O., Fitzgerald, G.F., Deane, J., O'Connor, M., Harnedy, N., O'Connor, K., O'Mahony, D., van Sinderen, D., Wallace, M., Brennan, L., Stanton, C., Marchesi, J.R., Fitzgerald, A.P., Shanahan, F., Hill, C., Ross, R.P., and O'Toole, P.W.: Gut microbiota composition correlates with diet and health in the elderly. Nature 488, 178-184 (2012).
- Congrains, A., Kamide, K., Ohishi, M., and Rakugi, H.: ANRIL: molecular mechanisms and implications in human health. Int. J. Mol. Sci. 14, 1278-1292 (2013).
- Dean, R.L., 3rd, Scozzafava, J., Goas, J.A., Regan, B., Beer, B., and Bartus, R.T.: Agerelated differences in behavior across the life span of the C57BL/6J mouse. Exp. Aging Res. 7, 427-451 (1981).
- Devallière, J. and Charreau, B.: The adaptor Lnk (SH2B3): an emerging regulator in vascular cells and a link between immune and inflammatory signaling. Biochem. Pharmacol. 82, 1391-1402 (2011).
- Dexter, D.T. and Jenner, P.: Parkinson disease: from pathology to molecular disease mechanisms. Free Radic. Biol. Med. 62, 132-144 (2013).
- Diao, H. and Liu, R.: The Eigencurve is Proper. Duke Math. J. 165, 1381-1395 (2016).
- Franceschi, C., Bonafè, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E., and De Benedictis, G.: Inflamm-aging: an evolutionary perspective on immunosenescence. Ann. NY Acad. Sci. 908, 244-254

(2000).

- Franceschi, C. and Campisi, J.: Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. J. Gerontol. A Biol. Sci. Med. Sci. 69, Suppl. 1, S4-9 (2014).
- Fransen, F., van Beek, A.A., Borghuis, T., Aidy, S.E., Hugenholtz, F., van der Gaastde Jongh, C., Savelkoul, H.F.J., De Jonge, M.I., Boekschoten, M.V., Smidt, H., Faas, M.M., and de Vos, P.: Aged Gut Microbiota Contributes to Systemical Inflammaging after Transfer to Germ-Free Mice. Front. Immunol. 8, 1385 (2017).
- Fried, L.P., Bandeen-Roche, K., Kasper, J.D., and Guralnik, J.M.: Association of comorbidity with disability in older women: the Women's Health and Aging Study. J. Clin. Epidemiol. 52, 27-37 (1999).
- Fulde, M., Sommer, F., Chassaing, B., van Vorst, K., Dupont, A., Hensel, M., Basic, M., Klopfleisch, R., Rosenstiel, P., Bleich, A., Bäckhed, F., Gewirtz, A.T., and Hornef, M.W.: Neonatal selection by Toll-like receptor 5 influences long-term gut microbiota composition. Nature 560, 489-493 (2018).
- Galkina, E. and Ley, K.: Immune and inflammatory mechanisms of atherosclerosis (*). Annu. Rev. Immunol. 27, 165-197 (2009).
- Genade, T., Benedetti, M., Terzibasi, E., Roncaglia, P., Valenzano, D.R., Cattaneo, A., and Cellerino, A.: Annual fishes of the genus Nothobranchius as a model system for aging research. Aging Cell 4, 223-233 (2005).
- Harding, C., Pompei, F., and Wilson, R.: Peak and decline in cancer incidence, mortality, and prevalence at old ages. Cancer 118, 1371-1386 (2012).
- Benjamini, Y. and, Hochberg Y.: Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J. R. Statist. Soc. B 57, 289-300 (1995).
- International Cancer Genome Consortium, Hudson, T.J., Anderson, W., Artez, A., Barker, A.D., Bell, C., Bernabé, R.R., Bhan, M.K., Calvo, F., Eerola, I., Gerhard, D.S., Guttmacher, A., Guyer, M., Hemsley, F.M., Jennings, J.L., Kerr, D., Klatt, P., Kolar, P., Kusada, J., Lane, D.P., Laplace, F., Youyong, L., Nettekoven, G., Ozenberger,

B., Peterson, J., Rao, T.S., Remacle, J., Schafer, A.J., Shibata, T., Stratton, M.R., Vockley, J.G., Watanabe, K., Yang, H., Yuen, M.M.F., Knoppers, B.M., Bobrow, M., Cambon-Thomsen, A., Dressler, L.G., Dyke, S.O.M., Joly, Y., Kato, K., Kennedy, K.L., Nicolás, P., Parker, M.J., Rial-Sebbag, E., Romeo-Casabona, C.M., Shaw, K.M., Wallace, S., Wiesner, G.L., Zeps, N., Lichter, P., Biankin, A.V., Chabannon, C., Chin, L., Clément, B., de Alava, E., Degos, F., Ferguson, M.L., Geary, P., Hayes, D.N., Hudson, T.J., Johns, A.L., Kasprzyk, A., Nakagawa, H., Penny, R., Piris, M.A., Sarin, R., Scarpa, A., Shibata, T., van de Vijver, M., Futreal, P.A., Aburatani, H., Bayés, M., Botwell, D.D.L., Campbell, P.J., Estivill, X., Gerhard, D.S., Grimmond, S.M., Gut, I., Hirst, M., López-Otín, C., Majumder, P., Marra, M., McPherson, J.D., Nakagawa, H., Ning, Z., Puente, X.S., Ruan, Y., Shibata, T., Stratton, M.R., Stunnenberg, H.G., Swerdlow, H., Velculescu, V.E., Wilson, R.K., Xue, H.H., Yang, L., Spellman, P.T., Bader, G.D., Boutros, P.C., Campbell, P.J., Flicek, P., Getz, G., Guigó, R., Guo, G., Haussler, D., Heath, S., Hubbard, T.J., Jiang, T., Jones, S.M., Li, Q., López-Bigas, N., Luo, R., Muthuswamy, L., Ouellette, B.F.F., Pearson, J.V., Puente, X.S., Quesada, V., Raphael, B.J., Sander, C., Shibata, T., Speed, T.P., Stein, L.D., Stuart, J.M., Teague, J.W., Totoki, Y., Tsunoda, T., Valencia, A., Wheeler, D.A., Wu, H., Zhao, S., Zhou, G., Stein, L.D., Guigó, R., Hubbard, T.J., Joly, Y., Jones, S.M., Kasprzyk, A., Lathrop, M., López-Bigas, N., Ouellette, B.F.F., Spellman, P.T., Teague, J.W., Thomas, G., Valencia, A., Yoshida, T., Kennedy, K.L., Axton, M., Dyke, S.O.M., Futreal, P.A., Gerhard, D.S., Gunter, C., Guyer, M., Hudson, T.J., McPherson, J.D., Miller, L.J., Ozenberger, B., Shaw, K.M., Kasprzyk, A., Stein, L.D., Zhang, J., Haider, S.A., Wang, J., Yung, C.K., Cros, A., Liang, Y., Gnaneshan, S., Guberman, J., Hsu, J., Bobrow, M., Chalmers, D.R.C., Hasel, K.W., Joly, Y., Kaan, T.S.H., Kennedy, K.L., Knoppers, B.M., Lowrance, W.W., Masui, T., Nicolás, P., Rial-Sebbag, E., Rodriguez, L.L., Vergely, C., Yoshida, T., Grimmond, S.M.,

Biankin, A.V., Bowtell, D.D.L., Cloonan, N., deFazio, A., Eshleman, J.R., Etemadmoghadam, D., Gardiner, B.B., Kench, J.G., Scarpa, A., Sutherland, R.L., Tempero, M.A., Waddell, N.J., Wilson, P.J., McPherson, J.D., Gallinger, S., Tsao, M.-S., Shaw, P.A., Petersen, G.M., Mukhopadhyay, D., Chin, L., DePinho, R.A., Thayer, S., Muthuswamy, L., Shazand, K., Beck, T., Sam, M., Timms, L., Ballin, V., Lu, Y., Ji, J., Zhang, X., Chen, F., Hu, X., Zhou, G., Yang, Q., Tian, G., Zhang, L., Xing, X., Li, X., Zhu, Z., Yu, Y., Yu, J., Yang, H., Lathrop, M., Tost, J., Brennan, P., Holcatova, I., Zaridze, D., Brazma, A., Egevard, L., Prokhortchouk, E., Banks, R.E., Uhlén, M., Cambon-Thomsen, A., Viksna, J., Ponten, F., Skryabin, K., Stratton, M.R., Futreal, P.A., Birney, E., Borg, A., Børresen-Dale, A.-L., Caldas, C., Foekens, J.A., Martin, S., Reis-Filho, J.S., Richardson, A.L., Sotiriou, C., Stunnenberg, H.G., Thoms, G., van de Vijver, M., van't Veer, L., Calvo, F., Birnbaum, D., Blanche, H., Boucher, P., Boyault, S., Chabannon, C., Gut, I., Masson-Jacquemier, J.D., Lathrop, M., Pauporté, I., Pivot, X., Vincent-Salomon, A., Tabone, E., Theillet, C., Thomas, G., Tost, J., Treilleux, I., Calvo, F., Bioulac-Sage, P., Clément, B., Decaens, T., Degos, F., Franco, D., Gut, I., Gut, M., Heath, S., Lathrop, M., Samuel, D., Thomas, G., Zucman-Rossi, J., Lichter, P., Eils, R., Brors, B., Korbel, J.O., Korshunov, A., Landgraf, P., Lehrach, H., Pfister, S., Radlwimmer, B., Reifenberger, G., Taylor, M.D., von Kalle, C., Majumder, P.P., Sarin, R., Rao, T.S., Bhan, M.K., Scarpa, A., Pederzoli, P., Lawlor, R.A., Delledonne, M., Bardelli, A., Biankin, A.V., Grimmond, S.M., Gress, T., Klimstra, D., Zamboni, G., Shibata, T., Nakamura, Y., Nakagawa, H., Kusada, J., Tsunoda, T., Miyano, S., Aburatani, H., Kato, K., Fujimoto, A., Yoshida, T., Campo, E., López-Otín, C., Estivill, X., Guigó, R., de Sanjosé, S., Piris, M.A., Montserrat, E., González-Díaz, M., Puente, X.S., Jares, P., Valencia, A., Himmelbauer, H., Quesada, V., Bea, S., Stratton, M.R., Futreal, P.A., Campbell, P.J., Vincent-Salomon, A., Richardson, A.L., Reis-Filho, J.S., van de Vijver, M., Thomas, G.,

MassonJacquemier, J.D., Aparicio, S., Borg, A., Børresen-Dale, A.-L., Caldas, C., Foekens, J.A., Stunnenberg, H.G., van't Veer, L., Easton, D.F., Spellman, P.T., Martin, S., Barker, A.D., Chin, L., Collins, F.S., Compton, C.C., Ferguson, M.L., Gerhard, D.S., Getz, G., Gunter, C., Guttmacher, A., Guyer, M., Hayes, D.N., Lander, E.S., Ozenberger, B., Penny, R., Peterson, J., Sander, C., Shaw, K.M., Speed, T.P., Spellman, P.T., Vockley, J.G., Wheeler, D.A., Wilson, R.K., Hudson, T.J., Chin, L., Knoppers, B.M., Lander, E.S., Lichter, P., Stein, L.D., Stratton, M.R., Anderson, W., Barker, A.D., Bell, C., Bobrow, M., Burke, W., Collins, F.S., Compton, C.C., DePinho, R.A., Easton, D.F., Futreal, P.A., Gerhard, D.S., Green, A.R., Guyer, M., Hamilton, S.R., Hubbard, T.J., Kallioniemi, O.P., Kennedy, K.L., Ley, T.J., Liu, E.T., Lu, Y., Majumder, P., Marra, M., Ozenberger, B., Peterson, J., Schafer, A.J., Spellman, P.T., Stunnenberg, H.G., Wainwright, B.J., Wilson, R.K., and Yang, H.: International network of cancer genome projects. Nature 464, 993-998 (2010).

- Jones, O.R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C.G., Schaible, R., Casper, B.B., Dahlgren, J.P., Ehrlén, J., García, M.B., Menges, E.S., Quintana-Ascencio, P.F., Caswell, H., Baudisch, A., and Vaupel, J.W.: Diversity of ageing across the tree of life. Nature 505, 169-173 (2014).
- Jurk, D., Wilson, C., Passos, J.F., Oakley, F., Correia-Melo, C., Greaves, L., Saretzki, G., Fox, C., Lawless, C., Anderson, R., Hewitt, G., Pender, S.L., Fullard, N., Nelson, G., Mann, J., van de Sluis, B., Mann, D.A., and von Zglinicki, T.: Chronic inflammation induces telomere dysfunction and accelerates ageing in mice. Nat. Commun. 2, 4172 (2014).
- Langille, M.G., Meehan, C.J., Koenig, J.E., Dhanani, A.S., Rose, R.A., Howlett, S.E., and Beiko, R.G.: Microbial shifts in the aging mouse gut. Microbiome 2, 50 (2014).
- Lee, C.K., Weindruch, R., and Prolla, T.A.: Gene-expression profile of the ageing brain in mice. Nat. Genet. 25, 294-297 (2000).
- Lopez, A.D., Mathers, C.D., Ezzati, M., Jamison, D.T., and Murray, C.J.L.: Global and regional burden of disease and risk

factors, 2001: systematic analysis of population health data. Lancet 367, 1747-1757 (2006).

- López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G.: The hallmarks of aging. Cell 153, 1194-1217 (2013).
- Lozupone, C.A., Stombaugh, J.I., Gordon, J.I., Jansson, J.K., and Knight, R.: Diversity, stability and resilience of the human gut microbiota. Nature 489, 220-230 (2012).
- Lynn, D.J., Winsor, G.L., Chan, C., Richard, N., Laird, M.R., Barsky, A., Gardy, J.L., Roche, F.M., Chan, T.H.W., Shah, N., Lo, R., Naseer, M., Que, J., Yau, M., Acab, M., Tulpan, D., Whiteside, M.D., Chikatamarla, A., Mah, B., Munzner, T., Hokamp, K., Hancock, R.E.W., and Brinkman, F.S.L.: InnateDB: facilitating systems-level analyses of the mammalian innate immune response. Mol. Syst. Biol. 4, 218 (2008).
- Masoro, E.J.: Overview of caloric restriction and ageing. Mech. Ageing Dev. 126, 913-922 (2005).
- O'Toole, P.W. and Jeffery, I.B.: Gut microbiota and aging. Science 350, 1214-1215 (2015).
- Pan, W.-H., Sommer, F., Falk-Paulsen, M., Ulas, T., Best, P., Fazio, A., Kachroo, P., Luzius, A., Jentzsch, M., Rehman, A., Müller, F., Lengauer, T., Walter, J., Künzel, S., Baines, J.F., Schreiber, S., Franke, A., Schultze, J.L., Bäckhed, F., and Rosenstiel, P.: Exposure to the gut microbiota drives distinct methylome and transcriptome changes in intestinal epithelial cells during postnatal development. Genome Med. 10, 27 (2018).
- Pettan-Brewer, C. and Treuting, P.M.: Practical pathology of aging mice. Pathobiol. Aging Age Relat. Dis. 1, doi: 10.3402/pba.v1i0.7202 (2011).
- Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., and Siepel, A.: Detection of nonneutral substitution rates on mammalian phylogenies. Genome Res. 20, 110-121 (2010).
- Reimand, J., Arak, T., Adler, P., Kolberg, L., Reisberg, S., Peterson, H., and Vilo, J.: g:Profiler-a web server for functional interpretation of gene lists (2016 update). Nucleic Acids Res. 44, W83-89 (2016).

- Reitz, C. and Mayeux, R.: Alzheimer's Disease; Epidemiology. In: Encyclopedia of the Neurological Sciences (Eds.: Aminoff, M.J. and Daroff, R.B.). Academic Press, London: 128-133 (2014).
- Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson, H., Spieth, J., Hillier, L.W., Richards, S., Weinstock, G.M., Wilson, R.K., Gibbs, R.A., Kent, W.J., Miller, W., and Haussler, D.: Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res. 15, 1034-1050 (2005).
- Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic, M., Roth, A., Santos, A., Tsafou, K.P., Kuhn, M., Bork, P., Jensen, L.J., and von Mering, C.: STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic Acids Res. 43, D447-452 (2015).
- Tazume, S., Umehara, K., Matsuzawa, H., Aikawa, H., Hashimoto, K., and Sasaki, S.: Effects of germfree status and food restriction on longevity and growth of mice. Jikken Dobutsu 40, 517-522 (1991).
- Teta, M., Long, S.Y., Wartschow, L.M., Rankin, M.M., and Kushner, J.A.: Very slow turnover of beta-cells in aged adult mice. Diabetes 54, 2557-2567 (2005).
- Thevaranjan, N., Puchta, A., Schulz, C., Naidoo, A., Szamosi, J.C., Verschoor, C.P., Loukov, D., Schenck, L.P., Jury, J., Foley, K.P., Schertzer, J.D., Larché, M.J., Davidson, D.J., Verdú, E.F., Surette, M.G., and Bowdish, D.M.E.: Age-Associated Microbial Dysbiosis Promotes Intestinal Permeability, Systemic Inflammation, and Macrophage Dysfunction. Cell Host Microbe 23, 570 (2018).
- Valko, M., Izakovic, M., Mazur, M., Rhodes, C.J. and Telser, J.: Role of oxygen radicals in DNA damage and cancer incidence. Mol. Cell. Biochem. 266, 37-56 (2004).
- Wostmann, B.S.: The germfree animal in nutritional studies. Ann. Rev. Nutr. 1, 257-279 (1981).

AGEING AND THE MICROBIOME – LESSONS FROM NON-SENESCENT MODELS

BENEDIKT M. MORTZFELD and THOMAS C.G. BOSCH

Zoological Institute, Christian-Albrechts-University Kiel, Kiel, Germany

SUMMARY

Research on the biological processes of ageing is at the brink of a revolution with respect to our understanding of its underlying mechanisms as well as our ability to prevent and cure a wide variety of age-related pathologies. Ageing is ordinarily recognized as a process that results from the combined influence of genetic and epigenetic determinants, life-style associated factors and external events. Studies using model organisms such as budding and fission yeasts, the nematode *Caenorhabditis elegans*, zebrafish and mice have generated significant insight into (epi)genetic pathways involved in ageing. However, developing an integrated understanding of these diverse processes and, thereby, achieving insight into the causes and mechanisms of the ageing process, remain a challenge. Here we review recent studies in non-senescent *Hydra* which demonstrate that a continuously high activity of the transcription factor FoxO contributes to continuous stem cell proliferation and at the same time also supports robust colonization of epithelia with a stable microbiome.

THE SIGNATURE OF AGEING

Numerous studies show that ageing is not a series of stochastic defects but a directed biological program that has a conserved genetic background (Hitt et al., 1999; Kenyon, 2010; Jeck et al., 2012; *López-Otín* et al., 2013) and is dependent on both genetic and environmental factors (López-Otín et al., 2013; Dato et al., 2017). Remarkably, only two proteins have been consistently associated with human ageing and longevity – Apolipoprotein E (Apo-E) and forkhead-box protein O3 (FoxO3). The genes were initially discovered in candidate gene studies of long-lived French (Apo-E) (Schächter et al., 1994) and long-lived Japanese/Okinawans in Hawaii (FoxO3) (Willcox et al., 2008), by comparing selected candidate gene frequency between long-lived cases (mainly centenarians) and shorter-lived controls. These findings were subsequently replicated in numerous populations worldwide (Morris et al., 2015). Genome-wide association studies (GWAS) utilizing larger numbers of cases and controls confirmed these genes as impacting human longevity but no other candidates have emerged with widespread replication (Broer et al., 2015). Apo-E plays an important role in clearance of cholesterol and lipoproteins from the circulatory system in vertebrates (Phillips, 2014). The gene consists of three alleles – the 'protective' allele $\epsilon 2$, the most common 'neutral' allele $\epsilon 3$ and the 'risk' allele ɛ4. While ɛ2 is associated with reduced risk for CVD and dementia as well as with increased life

span, $\varepsilon 4$ contributes to opposite effects (Schächter et al., 1994; Bennet et al., 2007; Nebel et al., 2011; Lindahl-Jacobsen et al., 2013). FoxO is a conserved transcription factor that shuttles between cytoplasm and nucleus (Van Der Heide et al., 2004) and binds directly to the regulatory sequences of its target genes (Webb et al., 2016). Four mammalian members of the FoxO family are described, FoxO1, FoxO3, FoxO4 and FoxO6. While FoxO6 is rather specific for neurons, the other FoxOs are expressed in most tissues and are causally linked to cell survival, cellular proliferation and DNA damage repair response (Monsalve and Olmos, 2011). In humans, a 1.5 Kb region of chromosome 6, in or near intron 2, contains dozens of non-coding foxO3 single nucleotide polymorphisms (SNPs), in high linkage disequilibrium (close association) (Donlon et al., 2012). These foxO3 SNPs vary in frequency by ethnicity, but one or more SNPs are directly associated with life span in all populations. This finding has been observed in case-control studies of centenarians and shorter-lived controls as well as cohort studies of older individuals (Broer et al., 2015; Santos-Lozano et al., 2016). Furthermore, since the FoxO protein and its target sites are highly conserved, analyses of human, mouse, C. elegans and Drosophila homologs recently uncovered FoxO as being a central hub in regulating a network of ageing-related genes (Webb et

al., 2016). Functional analyses in model organisms verified that the level of FoxO expression is indeed directly linked to life span without detectable costs for the individuals (Kenyon et al., 1993; Hwangbo et al., 2004; Schaible and Sussman, 2013). Several human tissues experience synchronized changes in gene expression during the ageing process which were shown to be a direct cause of age-related diseases (Yang et al., 2015). Furthermore, also the immune system is affected by the ageing process. "Immunosenescence" describes the deteriorating function of the immune system which results in increased susceptibility of elderly populations to infection, autoimmune diseases and cancer (Pawalec, 1999; Castle, 2000). The breakdown of mechanical barriers like the epithelia of lung, skin or intestinal tract leads to increased incidences of pathogenic invasion (Gomez et al., 2005). Combined with the declined functionality of immune cells and dysregulation of central components of innate immunity, elderly individuals observe a lower effectiveness in coping with the increased inflammation status, consequently resulting in higher morbidity and mortality (Franceschi et al., 2005; Rosenstiel et al., 2008). Although the understanding of FoxO signalling antagonizing ageing appears crucial to explain life span and health span in humans, the causative role of FoxO can only be addressed in model organisms.

MICROBES MATTER DURING THE AGEING PROCESS

All living beings are metaorganisms, associated with myriads of viruses, archaea and bacteria (*Zilber-Rosenberg* and *Rosenberg*, 2008; *Bosch* and *McFall-Ngai*, 2011). High throughput sequencing studies on humans showed that all epithelia are colonized with different, distinct microbial communities (*Eckberg* et al., 2005; *Grice* and *Segre*, 2011). Furthermore, many severe and chronic diseases, *e.g.* irritable bowel syndrome, inflammatory bowel disease, allergy or asthma, are associated with an altered bacterial

composition on the affected site (Khosravi and Mazmanian, 2013; Fujimura and Lynch, 2015). Studies in model organisms show that symbiotic bacteria can provide essential amino acids to the host (Sandström and Moran, 1999; Shigenobu et al., 2000), detoxify harmful substances (Chaucheyras-Durand et al., 2010) or degrade complex carbohydrates and thereby make them accessible to the host (Warnecke et al., 2007; Keeney and Finlay, 2011). Additionally, the composition of the gut microbiota regulates the nutrition uptake and is involved in weight control in mice and humans (Ley et al., 2005, 2006). Therefore, an organism's health status is reliant on an intact gut microflora (Goodman et al., 2009; Blottière et al., 2013; Khosravi and Mazmanian, 2013). Recent research also underlines the importance of bacterial colonization on host behaviour (Vuong et al., 2017) as well as fundamental developmental processes involving cell proliferation and ımmune system maturation (Rakoff-Nahoum et al., 2004; Mazmanian et al., 2005; Bates et al., 2006; Cheesman et al., 2011; Hill et al., 2016).

Interestingly, the human microbiome is not stable throughout life but experiences significant changes with age. First inoculation occurs during birth as new-borns are colonized with birth canal specific taxa (*Dominguez-Bello* et al., 2016). Disruption of this transmission via caesarean section or prenatal use of antibiotics may increase the risk of coeliac disease, asthma or obesity (Couzin-Frankel et al., 2010; Decker et al., 2010; Ajslev et al., 2011; *Metsälä* et al., 2015) and therefore have a huge impact on the individual's health. The intestinal microbiota resembles an adult stage already at the age of three (Yatsunenko et al., 2012), while the community of the skin microbes still undergoes drastic changes in puberty before it reaches a robust state (Oh et al., 2012). Over adulthood the bacterial communities are very stable but start to change again in elderly cohorts. The ageing process affects the structure of the human gut microbiota which results in a decrease in species diversity and a higher risk of pathogenic infections (Keller and Surawicz, 2014). In model organisms the structure of the bacterial community is discussed to have a direct influence on the ageing process as it can be protective against pathogens, shapes the nutrient landscape and affects the inflammation status (Heintz and Mair, 2014). In the short-lived African turquoise killifish the microbial community of young donors is able to extend life span in older individuals by preventing the decrease in microbial diversity associated with host ageing (Smith et al., 2017). Concordantly, latest research in C. elegans describes the positive impact of certain bacterial genotypes on host life span and health as non-essential bacterial compounds, e.g. colanic acid, are able to regulate mitochondrial dynamics and unfolded protein response (UPR^{mt}) (*Han* et al., 2017).

NON-SENESCENT MODEL ORGANISMS TO STUDY THE INTERACTION BETWEEN FOXO AND MICROBES

While most ageing research has been done in short-living models, some animals show no ageing phenotype and are considered as non-senescent. The freshwater polyp *Hydra* (Figure 1A) has an estimated life span that exceeds 1400 years (*Jones* et al., 2014). It belongs to the class of *Hydrozoa* within



Figure 1: Conserved transcription factor FoxO controls non-senescence and innate immunity in Hydra. (A) Non-senescent Hydra. Continuous proliferation of the stem cell compartment provides the potential of an unlimited life span and clonal growth by asexual budding. (B) Model depicting the transcription factor FoxO as the key regulator of tissue maintenance and as link to the innate immune system. In response to environmental or bacterial signals FoxO can be shuttled between the transcriptionally inactive state in the cytoplasm and the active form in the nucleus. FoxO targets include genes of tissue maintenance and AMPs that directly loop back to the associated microbes. Therefore, FoxO is not only an ageing antagonist but could also be involved in maintenance of the metaorganism (from: *Mortzfeld* and *Bosch*, 2017).

the phylum of *Cnidaria*, the sister group of bilaterians, and therefore holds a basal position in the animal tree of life (Martindale et al., 2002). Due to three everlasting stem cell lineages that give rise to about 20 cell types (Bosch, 2009), the animals are able to continuously reproduce asexually via budding (Figure 1A). A single homolog of the longevity gene foxO was found to be expressed in all three stem cell lineages in Hydra (Hemmrich et al., 2012). While earlier Hydra studies associated the transcription factor with apoptosis (Lasi et al., 2010) and stress resistance (Bridge et al., 2010), functional analyses uncovered FoxO's role in both stem cell regulation and innate immunity (Boehm et al., 2012). In situ hybridizations of the cnidarians Clytia and Hydra revealed foxO expression in the area of cell proliferation (Chevalier et al., 2006; Boehm et al, 2012). Epithelial FoxO deficiency for the latter resulted in a significant increase of differentiated cell mass in relation to the stem cell compartment which resembles the ageing phenotype in senescent animals. Inversely, an overexpression in the interstitial stem cell lineage caused not only an increase in cell proliferation, but also expression of stem cell marker genes in terminally differentiated cells (Boehm et al., 2012). FoxO's capacity in other model organisms [Drosophila (Hwangbo et al., 2004) and C. elegans (Kenyon et al., 1993)] to delay the onset of age-related processes and to extend life span underlines its conserved potential in stem cell control and tissue maintenance, especially in the absence of classical stem cell

factors like Oct-3/4 or SOX2 (*Chapman* et al., 2010). Remarkably, the changes in tissue homeostasis were accompanied by altered expression of antimicrobial peptides (AMPs), effectors of the innate immune system.

Microbial colonization has been studied intensively in chidarians as they have a huge impact on marine ecology but also offer a great opportunity to study bacteria-host interactions in organisms of lower complexity. Intracellular photosynthetic dinoflagellates (e.g. Symbiodinium) allow corals to populate nutrient-poor environments (Muscatine and Porter, 1977), however, recent research has shown that basal metazoans in addition to eukaryotic photobionts also harbour extracellular, species-specific bacterial communities (Rohwer et al., 2002; Littman et al., 2009; Franzenburg et al., 2013a). In the case of *Hydra* even closely related species are colonized by specific microbiomes that mirror the phylogenetic relatedness between their host species (phylosymbiosis; *Brooks* et al., 2016). The species-specific microbiomes in cnidarians are extremely stable and still resemble the ones of animals from the wild even after years of culturing under laboratory conditions involving artificial water and feeding (Fraune and Bosch, 2007; Franzenburg et al., 2013a; Mortzfeld et al., 2016). The acquisition of bacterial symbionts occurs during early development and establishes in a robust, reproducible pattern (Franzenburg et al., 2013b; Mortzfeld et al., 2016). Specific bacterial taxa may not only be acquired horizontally (Apprill et al., 2009; Sharp et al., 2010) but in some cases also via vertical transmission (parent to offspring) (Fraune et al., 2010; Sharp et al., 2012). Certain taxa of free-living bacteria can even be attracted by the host using chemotactic gradients (*Tout* et al., 2015). Underlining the importance of the microbiome, a study with germ-free animals showed that Hydra is protected from fungal infections by its bacterial colonizers. Interestingly, recolonization experiments proofed that a combination of the two most abundant bacteria of the naturally occurring microbiota [Curvibacter sp. 75% and Duganella sp. 10% (Franzenburg et al., 2013a)] was most protective, while monoassociations resulted in insufficient effects (Fraune et al., 2015).

AGEING ANTAGONIST FOXO CONTROLS BOTH STEM CELLS AND MICROBES

Besides its well-known conserved function as major tissue regulator and ageing antagonist, FoxO modulates the innate immune system in various model organisms including mouse 2013), Drosophila (Seiler et al., (Becker et al., 2010), C. elegans (Libina et al., 2003) and Hydra (Boehm et al., 2012). In mice FoxO signalling has been shown to reduce susceptibility to bacterial infections by reducing oxidative stress and induction of inflammatory cytokines (*Joseph* et al., 2016).

Furthermore, FoxO transcription factors directly regulate TLR3-mediated innate immune responses as well as the expression of AMPs and thereby contribute to pathogen clearance in the respiratory tract (*Seiler* et al., 2013). Also in *Drosophila* AMPs are well-known effector molecules of the innate immune system and important regulators of the bacterial colonizers. Here, oral bacterial infection induces FoxO activity in the intestine, while impaired FoxO signalling decreases resistance to

intestinal infections. The inability to raise the expression level of AMPs leads to an elevated bacterial load and a decline in survival (Fink et al., 2015). In *Hydra*, the microbiome is selectively assembled by a species-specific combination of AMPs which are predominantly expressed in epithelial cells (Fraune et al., 2010; Franzenburg et al., 2013a). FoxO-deficient Hydra polyps show in addition to defects in stem cell maintenance a severe change of the immune status and drastically altered expression of AMPs (Boehm et al., 2012). Remarkably, loss of tissue homeostasis as well as AMP-deficiency compromise the ability to select for microbial communities resembling the polyps' native microbiota (Fraune et al., 2009; Franzenburg et al., 2013a). Interestingly, a recent study indicates that FoxO signalling may also be involved in the establishment of symbiosis in the early developmental phase of the sea anemone Aiptasia (Wolfowicz et al., 2016).

Taken together, the conserved transcription factor FoxO appears to combine two functions crucially involved in ageing and health in metazoans (Figure 1B): FoxO is responsible for stem cell regulation, including tissue maintenance and renewal, and controls the innate immune system. In response to environmental (including bacterial) signals, FoxO switches between a transcriptionally inactive state in the cytoplasm and an active form in the nucleus thereby serving as an intracellular control board for environmental signals. The capabilities of the FoxO transcription factor to extend life span and control effectors of the immune system demonstrate a strong and unique mechanism of cross-regulation of tissue homeostasis and innate immunity.

By exploring epithelial FoxO lossof-function mutants, we recently made two important discoveries (*Mortzfeld* et al., 2018). First, deficiency in FoxO signalling leads to dysregulation of multiple AMP families. Most genes encoding epithelially expressed AMP families including Hydramacin, Kazal and Arminin respond with downregulation to FoxO-deficiency. Only one gene (contig 45266) was found to be upregulated in FoxO deficient animals, suggesting a mainly activating function of FoxO signalling on AMP expression and innate immunity. Second, FoxO loss-of-function polyps were more susceptible to colonization of foreign bacteria and impaired in selection for bacteria resembling the native microbi-Therefore, FoxO-induced deome. crease in AMP expression is correlating with differences in microbial colonization and highlights the inhibitory action of AMPs against non-commensal bacteria. In a state of intact FoxO signalling secretion of numerous AMP families provides a highly selective milieu and shapes the microbial composition in a species-specific manner (Figure 1B). FoxO-deficiency reduces the expression of AMPs, which results in a decreased selection pressure on colonizing taxa and in establishment of higher abundances of foreign bacteria in the community. Consequently, especially during the process of colonization, reduced expression of FoxO compromises the resilience of the microbiome.

CONCLUSIONS

Stem cell maintenance and immunity are two of the prominent areas in

ageing research. Observations in different organisms indicate that, in contrast
to the essentially static genome, the microbiome is rather dynamic throughout life history. Our observations in nonsenescent *Hydra* add support to the view that (i) there is a need to consider the holobiotic nature of an organism when thinking about longevity, that (ii) the microbial environment matters in the context of senescence and contributes to complex processes such as ageing; and that (iii) the hub regulator FoxO presents a direct link between age-related processes and microbial colonization. In the newly discovered world of metaorganisms, stem cell proliferation and immunity are part of a global program, which seeks to fuse stem cell biology with ecological concepts and the rules governing the interactions between an organism and its microbial environment (*Mortzfeld* and *Bosch*, 2017; *Mortzfeld* et al., 2018).

ACKNOWLEDGEMENT

This work was supported by the Deutsche Forschungsgemeinschaft (DFG) (CRC1182 "Origin and Function of Metaorganisms", DFG grant BO 848/17-1, and grants from the DFG Cluster of Excellence program "Inflammation at Interfaces"). TCGB gratefully appreciates support from the Canadian Institute for Advanced Research (CIFAR).

LITERATURE

- Ajslev, T.A., Andersen, C.S., Gamborg, M., Sørensen, T.I., and Jess, T.: Childhood overweight after establishment of the gut microbiota: the role of delivery mode, prepregnancy weight and early administration of antibiotics. Int. J. Obes. (Lond). 35, 522-529 (2011).
- Apprill, A., Marlow, H.Q., Martindale, M.Q., and Rappé, M.S.: The onset of microbial associations in the coral Pocillopora meandrina. ISME J. 3, 685-699 (2009).
- Bates, J.M., Mittge, E., Kuhlman, J., Baden, K.N., Cheesman, S.E., and Guillemin, K.: Distinct signals from the microbiota promote different aspects of zebrafish gut differentiation. Dev. Biol. 297, 374-386 (2006).
- Becker, T., Loch, G., Beyer, M., Zinke, I., Aschenbrenner, A.C., Carrera, P., Inhester, T., Schultze, J.L., and Hoch, M.: FOXOdependent regulation of innate immune homeostasis. Nature 463, 369-373 (2010).
- Bennet, A.M., Di Angelantonio, E., Ye, Z., Wensley, F., Dahlin, A., Ahlbom, A., Keavney, B., Collins, R., Wiman, B., de

Faire, U., and Danesh, J.: Association of apolipoprotein E genotypes with lipid levels and coronary risk. JAMA 298, 1300-1311 (2007).

- Blottière, H.M., de Vos, W.M., Ehrlich, S.D., and Doré, J.: Human intestinal metagenomics: state of the art and future. Curr. Opin. Microbiol. 16, 234-239 (2013).
- Boehm, A.M., Khalturin, K., Anton-Erxleben, F., Hemmrich, G., Klostermeier, U.C., Lopez-Quintero, J.A., Oberg, H.H., Puchert, M., Rosenstiel, P., Wittlieb, J., and Bosch, T.C.: FoxO is a critical regulator of stem cell maintenance in immortal Hydra. Proc. Natl. Acad. Sci. USA 109, 19697-19702 (2012).
- Bosch, T.C.G.: Hydra and the evolution of stem cells. BioEssays 31, 478-486 (2009).
- Bosch, T.C.G. and McFall-Ngai, M.J.: Metaorganisms as the new frontier. Zoology 114, 185-190 (2011).
- Bourne, D.G., Morrow, K.M., and Webster, N.S.: Insights into the Coral Microbiome: Underpinning the Health and Resilience of Reef Ecosystems. Annu. Rev. Microbiol.

70, 317-340 (2016).

- Bridge, D., Theofiles, A.G., Holler, R.L., Marcinkevicius, E., Steele, R.E., and Martínez, D.E.: FoxO and stress responses in the cnidarian Hydra vulgaris. PLoS One 5, e11686 (2010).
- Broer, L., Buchman, A.S., Deelen, J., Evans, D.S., Faul, J.D., Lunetta, K.L., Sebastiani, P., Smith, J.A., Smith, A.V., Tanaka, T., Yu, L., Arnold, A.M., Aspelund, T., Benjamin, E.J., De Jager, P.L., Eirkisdottir, G., Evans, D.A., Garcia, M.E., Hofman, A., Kaplan, R.C., Kardia, S.L., Kiel, D.P., Oostra, B.A., Orwoll, E.S., Parimi, N., Psaty, B.M., Rivadeneira, F., Rotter, J.I., Seshadri, S., Singleton, A., Tiemeier, H., Uitterlinden, A.G., Zhao, W., Bandinelli, S., Bennett, D.A., Ferrucci, L., Gudnason, V., Harris, T.B., Karasik, D., Launer, L.J., Perls, T.T., Slagboom, P.E., Tranah, G.J., Weir, D.R., Newman, A.B., van Duijn, C.M., and Murabito, J.M.: GWAS of Longevity in CHARGE Consortium Confirms APOE and FOXO3 Candidacy. J. Gerontol. A Biol. Sci. Med. Sci. 70, 110-118 (2015).
- Brooks, A.W., Kohl, K.D., Brucker, R.M., van Opstal, E.J., and Bordenstein, S.R.: Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host Evolutionary History. PLoS Biol 14, e2000225 (2016).
- Castle, S.C.: Clinical relevance of age-related immune dysfunction. Clin. Infect. Dis. 31, 578-585 (2000).
- Chapman, J.A., Kirkness, E.F., Simakov, O., Hampson, S.E., Mitros, T., Weinmaier, T., Rattei, T., Balasubramanian, P.G., Borman, J., Busam, D., Disbennett, K., Pfannkoch, C., Sumin, N., Sutton, G.G., Viswanathan, L.D., Walenz, B., Goodstein, D.M., Hellsten, U., Kawashima, T., Prochnik, S.E., Putnam, N.H., Shu, S., Blumberg, B., Dana, C.E., Gee, L., Kibler, D.F., Law, L., Lindgens, D., Martinez, D.E., Peng, J., Wigge, P.A., Bertulat, B., Guder, C., Nakamura, Y., Ozbek, S., Watanabe, H., Khalturin, K., Hemmrich, G., Franke, A., Augustin, R., Fraune, S., Hayakawa, E., Hayakawa, S.,

Hirose, M., Hwang, J.S., Ikeo, K., Nishimiya-Fujisawa, С., Ogura, A., Takahashi, T., Steinmetz, P.R., Zhang, X., Aufschnaiter, R., Eder, M.K., Gorny, A.K., Salvenmoser, W., Heimberg, A.M., Wheeler, B.M., Peterson, K.J., Böttger, A., Tischler, P., Wolf, A., Gojobori, T., Remington, K.A., Strausberg, R.L., Venter, J.C., Technau, U., Hobmayer, B., Bosch, T.C., Holstein, T.W., Fujisawa, T., Bode, H.R., David, C.N., Rokhsar, D.S., and Steele, R.E.: The dynamic genome of Hydra. Nature 464, 592-596 (2010).

- Chaucheyras-Durand, F., Masséglia, S., Fonty, G., and Forano, E.: Influence of the composition of the cellulolytic flora on the development of hydrogenotrophic microorganisms, hydrogen utilization, and methane production in the rumens of gnotobiotically reared lambs. Appl. Environ. Microbiol. 76, 7931-7937 (2010).
- Cheesman, S.E., Neal, J.T., Mittge, E., Seredick, B.M., and Guillemin, K.: Epithelial cell proliferation in the developing zebrafish intestine is regulated by the Wnt pathway and microbial signaling via Myd88. Proc. Natl. Acad. Sci. USA 108, 4570-4577 (2011).
- Chevalier, S., Martin, A., Leclère, L., Amiel, A., and Houliston, E.: Polarised expression of FoxB and FoxQ2 genes during development of the hydrozoan Clytia hemisphaerica. Dev. Genes Evol. 216, 709-720 (2006).
- Couzin-Frankel, J.: Bacteria and asthma: untangling the links. Science 330, 1168-1169 (2010).
- Dato, S., Rose, G., Crocco, P., Monti, D., Garagnani, P., Franceschi, C., and Passarino, G.: The genetics of human longevity: an intricacy of genes, environment, culture and microbiome. Mech. Ageing Dev. 165, 147-155 (2017).
- Decker, E., Engelmann, G., Findeisen, A., Gerner, P., Laass, M., Ney, D., Posovszky, C., Hoy, L., and Hornef, M.W.: Cesarean delivery is associated with celiac disease but not inflammatory bowel disease in children. Pediatrics 125, e1433-40 (2010).

- Dominguez-Bello, M.G., De Jesus-Laboy, K.M., Shen, N., Cox, L.M., Amir, A., Gonzalez, A., Bokulich, N.A., Song, S.J., Hoashi, M., Rivera-Vinas, J.I., Mendez, K., Knight, R., and Clemente, J.C.: Partial restoration of the microbiota of cesareanborn infants via vaginal microbial transfer. Nat. Med. 22, 250-253 (2016).
- Donlon, T.A., Curb, J.D., He, Q., Grove, J.S., Masaki, K.H., Rodriguez, B., Elliott, A., Willcox, D.C., and Willcox, B.J.: FOXO3 gene variants and human aging: coding variants may not be key players. J. Gerontol. A Biol. Sci. Med. Sci. 67, 1132-1139 (2012).
- Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R., Nelson, K.E., and Relman, D.A.: Diversity of the Human Intestinal Microbial Flora. Science 308, 1635-1638 (2005).
- Fink, C., Hoffmann, J., Knop, M., Li, Y., Isermann, K., and Roeder, T.: Intestinal FoxO signaling is required to survive oral infection in Drosophila. Mucosal Immunol. 9, 927-936 (2015).
- Franceschi, C., Olivieri, F., Marchegiani, F., Cardelli, M., Cavallone, L., Capri, M., Salvioli, S., Valensin, S., De Benedictis, G., Di Iorio, A., Caruso, C., Paolisso, G., and Monti, D.: Genes involved in immune response/inflammation, IGF1/insulin pathway and response to oxidative stress play a major role in the genetics of human longevity: The lesson of centenarians. Mech. Ageing Dev. 126, 351-361 (2005).
- Franzenburg, S., Walter, J., Künzel, S., Wang, J., Baines, J.F., Bosch, T.C.G., and Fraune, S.: Distinct antimicrobial peptide expression determines host species-specific bacterial associations. Proc. Natl. Acad. Sci. USA 110, E3730-8 (2013a).
- Franzenburg, S., Fraune, S., Altrock, P.M., Künzel, S., Baines, J.F., Traulsen, A., and Bosch, T.C.G.: Bacterial colonization of Hydra hatchlings follows a robust temporal pattern. ISME J. 7, 781-790 (2013b).
- Fraune, S. and Bosch, T.C.G.: Long-term maintenance of species-specific bacterial microbiota in the basal metazoan Hydra.

Proc. Natl. Acad. Sci. USA 104, 13146-13151 (2007).

- Fraune, S., Abe, Y., and Bosch, T.C.G.: Disturbing epithelial homeostasis in the metazoan Hydra leads to drastic changes in associated microbiota. Environ. Microbiol. 11, 2361-2369 (2009).
- Fraune, S., Augustin, R., Anton-Erxleben, F., Wittlieb, J., Gelhaus, C., Klimovich, V.B., Samoilovich, M.P., and Bosch, T.C.G.: In an early branching metazoan, bacterial colonization of the embryo is controlled by maternal antimicrobial peptides. Proc. Natl. Acad. Sci. USA 107, 18067-18072 (2010).
- Fraune, S., Anton-Erxleben, F., Augustin, R., Franzenburg, S., Knop, M., Schröder, K., Willoweit-Ohl, D., and Bosch, T.C.G.: Bacteria-bacteria interactions within the microbiota of the ancestral metazoan Hydra contribute to fungal resistance. ISME J. 9, 1543-1556 (2015).
- Fraune, S., Forêt, S., and Reitzel, A.M.: Using Nematostella vectensis to Study the Interactions between Genome, Epigenome, and Bacteria in a Changing Environment. Front. Mar. Sci. 3, 148 (2016).
- Fujimura, K.E. and Lynch, S.V.: Microbiota in allergy and asthma and the emerging relationship with the gut microbiome. Cell Host Microbe 17, 592-602 (2015).
- Gomez, C.R., Boehmer, E.D., and Kovacs, E.J.: The aging innate immune system. Curr. Opin. Immunol. 17, 457-462 (2005).
- Goodman, A.L., McNulty, N.P., Zhao, Y., Leip, D., Mitra, R.D., Lozupone, C.A., Knight, R., and Gordon, J.I.: Identifying genetic determinants needed to establish a human gut symbiont in its habitat. Cell Host Microbe 6, 279-289 (2009).
- Grice, E.A. and Segre, J.A.: The skin microbiome. Nat. Rev. Microbiol. 9, 244-253 (2011).
- Han, B., Sivaramakrishnan, P., Lin, C.J., Neve, I.A.A., He, J., Tay, L.W.R., Sowa, J.N., Sizovs, A., Du, G., Wang, J., Herman, C., and Wang, M.C.: Microbial Genetic Composition Tunes Host Longevity. Cell 169, 1249-1262 (2017).
- Heintz, C. and Mair, W.: You are what you

host: microbiome modulation of the aging process. Cell 156, 408-411 (2014).

- Hemmrich, G., Khalturin, K., Boehm, A.M., Puchert, M., Anton-Erxleben, F., Wittlieb, J., Klostermeier, U.C., Rosenstiel, P., Oberg, H.H., Domazet-Lošo, T., Domazet-Loso, T., Sugimoto, T., Niwa, H., and Bosch, T.C.: Molecular signatures of the three stem cell lineages in hydra and the emergence of stem cell function at the base of multicellularity. Mol. Biol. Evol. 29, 3267-3280 (2012).
- Hill, J.H., Franzosa, E.A., Huttenhower, C., and Guillemin, K.: A conserved bacterial protein induces pancreatic beta cell expansion during zebrafish development. Elife 5, e20145 (2016).
- Hitt, R., Young-xu, Y., Silver, M., and Perls, T.: Centenarians: the older you get, the healthier you have been. Lancet 354, 2215 (1999).
- Hwangbo, D.S., Gershman, B., Tu, M.P., Palmer, M., Tatar, M.: Drosophila dFOXO controls lifespan and regulates insulin signalling in brain and fat body. Nature 429, 562-566 (2004).
- Jeck, W.R., Siebold, A.P., and Sharpless, N.E.: Review: a meta-analysis of GWAS and ageassociated diseases. Aging Cell 11, 727-731 (2012).
- Jones, O.R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C.G., Schaible, R., Casper, B.B., Dahlgren, J.P., Ehrlén, J., García, M.B., Menges, E.S., Quintana-Ascencio, P.F., Caswell, H., Baudisch, A., and Vaupel, J.W.: Diversity of ageing across the tree of life. Nature 505, 169-173 (2014).
- Joseph, J., Ametepe, E.S., Haribabu, N., Agbayani, G., Krishnan, L., Blais, A., and Sad, S.: Inhibition of ROS and upregulation of inflammatory cytokines by FoxO3a promotes survival against Salmonella typhimurium. Nat. Commun. 7, 12748 (2016).
- Keeney, K.M. and Finlay, B.B.: Enteric pathogen exploitation of the microbiota-generated nutrient environment of the gut. Curr. Opin. Microbiol. 14, 92-98 (2011).
- Keller, J.M. and Surawicz, C.M.: Clostridium difficile infection in the elderly. Clin.

Geriatr. Med. 30, 79-93 (2014).

- Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtlang, R.: A C. elegans mutant that lives twice as long as wild type. Nature 366, 461-464 (1993).
- Kenyon, C.J.: The genetics of ageing. Nature 464, 504-512 (2010).
- Khosravi, A. and Mazmanian, S.K.: Disruption of the gut microbiome as a risk factor for microbial infections. Curr. Opin. Microbiol. 16, 221-227 (2013).
- Lasi, M., David, C.N., and Böttger, A.: Apoptosis in pre-Bilaterians: Hydra as a model. Apoptosis 15, 269-278 (2010).
- Ley, R.E., Backhed, F., Turnbaugh, P., Lozupone, C.A., Knight, R.D., and Gordon, J.I.: Obesity alters gut microbial ecology. Proc. Natl. Acad. Sci. USA 102, 11070-11075 (2005).
- Ley, R., Turnbaugh, P., Klein, S., and Gordon, J.: Microbial ecology: human gut microbes associated with obesity. Nature 444, 1022-1023 (2006).
- Libina, N., Berman, J.R., and Kenyon, C.: Tissue-specific activities of C. elegans DAF-16 in the regulation of lifespan. Cell 115, 489-502 (2003).
- Lindahl-Jacobsen, R., Tan, Q., Mengel-From, J., Christensen, K., Nebel, A., and Christiansen, L.: Effects of the APOE ɛ2 allele on mortality and cognitive function in the oldest old. J. Gerontol. A Biol. Sci. Med. Sci. 68, 389-394 (2013).
- Littman, R.A., Willis, B.L., Pfeffer, C., and Bourne, D.G.: Diversities of coral-associated bacteria differ with location, but not species, for three acroporid corals on the Great Barrier Reef. FEMS Microbiol. Ecol. 68, 152-163 (2009).
- López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G.: The hallmarks of aging. Cell 153, 1194-1217 (2013).
- Martindale, M., Finnerty, J.R., and Henry, J.Q.: The Radiata and the evolutionary origin of the bilaterian body plan. Mol. Phylogenet. Evol. 24, 358-365 (2002).
- Mazmanian, S.K., Cui, H.L., Tzianabos, A.O., and Kasper, D.L.: An immunomodulatory

molecule of symbiotic bacteria directs maturation of the host immune system. Cell 122, 107-118 (2005).

- Metsälä, J., Lundqvist, A., Virta, L.J., Kaila, M., Gissler, M., and Virtanen, S.M.: Prenatal and post-natal exposure to antibiotics and risk of asthma in childhood. Clin. Exp. Allergy 45, 137-145 (2015).
- Monsalve, M. and Olmos, Y.: The complex biology of FOXO. Curr. Drug Targets 12, 1322-1350 (2011).
- Morris, B.J., Willcox, D.C., Donlon, T.A., and Willcox, B.J.: FOXO3: A Major Gene for Human Longevity--A Mini-Review. Gerontology 61, 515-525 (2015).
- Mortzfeld, B.M., Urbanski, S., Reitzel, A.M., Künzel, S., Technau, U., and Fraune, S.: Response of bacterial colonization in Nematostella vectensis to development, environment and biogeography. Environ. Microbiol. 18, 1764-1781 (2016).
- Mortzfeld, B. and Bosch, T.C.G.: Eco-Ageing: stem cells and microbes are controlled by ageing antagonist FoxO. Curr. Opin. Microbiol. 38, 181-187 (2017).
- Mortzfeld, B.M., Taubenheim, J., Fraune, S., Klimovich, A.V., and Bosch, T.C.G.: Stem cell transcription factor FoxO controls microbiome resilience in Hydra. Front Microbiol., 9, 629 (2018).
- Muscatine, L. and Porter, J.W.: Reef corals: mutualistic symbioses adapted to nutrientpoor environments. Bioscience 27, 454-460 (1977).
- Nebel, A., Kleindorp, R., Caliebe, A., Nothnagel, M., Blanché, H., Junge, O., Wittig, M., Ellinghaus, D., Flachsbart, F., Wichmann, H.E., et al.: A genome-wide association study confirms APOE as the major gene influencing survival in long-lived individuals. Mech. Ageing Dev. 132, 324-330 (2011).
- Oh, J., Conlan, S., Polley, E.C., Segre, J.A., and Kong, H.H.: Shifts in human skin and nares microbiota of healthy children and adults. Genome Med. 4, 77 (2012).
- Pawelec, G.: Immunosenescence: impact in the young as well as the old? Mech. Ageing Dev. 108, 1-7 (1999).

- Phillips, M.C.: Apolipoprotein e isoforms and lipoprotein metabolism. IUBMB Life 66, 616-623 (2014).
- Rakoff-Nahoum, S., Paglino, J., Eslami-Varzaneh, F., Edberg, S., and Medzhitov, R.: Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. Cell 118, 229-241 (2004).
- Rohwer, F., Seguritan, V., Azam, F., and Knowlton, N.: Diversity and distribution of coral-associated bacteria. Mar. Ecol. Prog. Ser. 243, 1-10 (2002).
- Rosenstiel, P., Derer, S., Till, A., Häsler, R., Eberstein, H., Bewig, B., Nikolaus, S., Nebel, A., and Schreiber, S.: Systematic expression profiling of innate immune genes defines a complex pattern of immunosenescence in peripheral and intestinal leukocytes. Genes Immun. 9, 103-114 (2008).
- Sandström, J. and Moran, N.: How nutritionally imbalanced is phloem sap for aphids? Entomol. Exp. Appl. 91, 203-210 (1999).
- Santos-Lozano, A., Santamarina, A., Pareja-Galeano, H., Sanchis-Gomar, F., Fiuza-Luces, C., Cristi-Montero, C., Bernal-Pino, A., Lucia, A., and Garatachea, N.: The genetics of exceptional longevity: Insights from centenarians. Maturitas 90, 49-57 (2016).
- Schächter, F., Faure-Delanef, L., Guénot, F., Rouger, H., Froguel, P., Lesueur-Ginot, L., and Cohen, D.: Genetic associations with human longevity at the APOE and ACE loci. Nat. Genet. 6, 29-32 (1994).
- Schaible, R. and Sussman, M.: FOXO in aging: Did evolutionary diversification of FOXO function distract it from prolonging life? Bioessays 35, 1101-1110 (2013).
- Schroeder, L.A. and Callaghan, W.M.: Thermal tolerance and acclimation of two species of Hydra. Limnol. Ocean. 26, 690-696 (1981).
- Seiler, F., Hellberg, J., Lepper, P.M., Kamyschnikow, A., Herr, C., Bischoff, M., Langer, F., Schafers, H.J., Lammert, F., Menger, M.D., Bals, R., and Beisswenger, C.: FOXO transcription factors regulate innate immune mechanisms in respiratory epithelial cells. J. Immunol. 190, 1603-1613 (2013).

- Sharp, K.H., Ritchie, K.B., Schupp, P.J., Ritson-Williams, R., and Paul, V.J.: Bacterial acquisition in juveniles of several broadcast spawning coral species. PLoS One 5, e10898 (2010).
- Sharp, K.H., Distel, D., and Paul, V.J.: Diversity and dynamics of bacterial communities in early life stages of the Caribbean coral Porites astreoides. ISME J. 6, 790-801 (2012).
- Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y., and Ishikawa, H.: Genome sequence of the endocellular bacterial symbiont of aphids Buchnera sp. APS. Nature 407, 81-86 (2000).
- Smith, P., Willemsen, D., Popkes, M., Metge, F., Gandiwa, E., Reichard, M., and Valenzano, D.R.: Regulation of life span by the gut microbiota in the short-lived African turquoise killifish. Elife 6, e27014 (2017).
- Tout, J., Jeffries, T.C., Petrou, K., Tyson, G.W., Webster, N.S., Garren, M., Stocker, R., Ralph, P.J., and Seymour, J.R.: Chemotaxis by natural populations of coral reef bacteria. ISME J. 9, 1764-1777 (2015).
- Van Der Heide, L.P., Hoekman, M.F.M., and Smidt, M.P.: The ins and outs of FoxO shuttling: mechanisms of FoxO translocation and transcriptional regulation. Biochem. J. 380, 297-309 (2004).
- Vuong, H.E., Yano, J.M., Fung, T.C., and Hsiao, E.Y.: The Microbiome and Host Behavior. Annu. Rev. Neurosci. 40, 21-49 (2017).
- Warnecke, F., Luginbühl, P., Ivanova, N., Ghassemian, M., Richardson, T.H., Stege, J.T., Cayouette, M., McHardy, A.C., Djordjevic, G., Aboushadi, N., Sorek, R., Tringe, S.G., Podar, M., Martin, H.G., Kunin, V., Dalevi, D., Madejska, J., Kirton, E., Platt, D., Szeto, E., Salamov, A., Barry, K., Mikhailova, N., Kyrpides, N.C., Matson, E.G., Ottesen, E.A., Zhang, X., Hernández, M., Murillo, C., Acosta, L.G., Rigoutsos, I., Tamayo, G., Green, B.D., Chang, C., Rubin, E.M., Mathur, E.J., Robertson, D.E., Hugenholtz, P., and Leadbetter, J.R.: Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher ter-

mite. Nature 450, 560-565 (2007).

- Webb, A.E., Kundaje, A., and Brunet, A.: Characterization of the direct targets of FOXO transcription factors throughout evolution. Aging Cell 15, 673-685 (2016).
- Willcox, B.J., Donlon, T.A., He, Q., Chen, R., Grove, J.S., Yano, K., Masaki, K.H., Willcox, D.C., Rodriguez, B., and Curb, J.D.: FOXO3A genotype is strongly associated with human longevity. Proc. Natl. Acad. Sci. USA 105, 13987-13992 (2008).
- Williams, R.B.: Starlet sea anemone: Nematostella vectensis. In: The IUCN invertebrate red data book (Eds.: Collins, N.M., Hughes, S.A., Pyle, R.M., and Wells, S.). IUCN, Gland, Switzerland: 43-46 (1983).
- Wolfowicz, I., Baumgarten, S., Voss, P.A., Hambleton, E.A., Voolstra, C.R., Hatta, M., and Guse, A.: Aiptasia sp. larvae as a model to reveal mechanisms of symbiont selection in cnidarians. Sci. Rep. 6, 32366 (2016).
- Yang, J., Huang, T., Petralia, F., Long, Q., Zhang, B., Argmann, C., Zhao, Y., Mobbs, C.V., Schadt, E.E., Zhu, J., and Tu, Z.: Synchronized age-related gene expression changes across multiple tissues in human and the link to complex diseases. Sci. Rep. 5, 15145 (2015).
- Yatsunenko, T., Rey, F.E., Manary, M.J., Trehan, I., Dominguez-Bello, M.G., Contreras, M., Magris, M., Hidalgo, G., Baldassano, R.N., Anokhin, A.P., Heath, A.C., Warner, B., Reeder, J., Kuczynski, J., Caporaso, J.G., Lozupone, C.A., Lauber, C., Clemente, J.C., Knights, D., Knight, R., and Gordon, J.I.: Human gut microbiome viewed across age and geography. Nature 486, 222-227 (2012).
- Ziegler, M., Seneca, F.O., Yum, L.K., Palumbi, S.R., and Voolstra, C.R.: Bacterial community dynamics are linked to patterns of coral heat tolerance. Nat. Commun. 8, 1-8 (2017).
- Zilber-Rosenberg, I. and Rosenberg, E.: Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. FEMS Microbiol. Rev. 32, 723-735 (2008).

AGEING AND GUT TO BRAIN SIGNALLING

WOLFGANG KUNZE

The McMaster Brain-Body Institute and Department of Psychiatry, St Joseph's Healthcare, Hamilton, ON, Canada.

INTRODUCTION

There are several major gut-brain communication pathways. These include signalling via the vagus and spinal nerves, immune factors via cytokines, endocrine signalling via gut hormones, and microbial products that reach the brain via bloodstream (*Holzer* and *Farzi*, 2014). In this brief essay, the focus is on the effects of ageing on gut to brain signalling via the vagus.

AGEING GENERAL

There is a general functional decline in all mammalian physiological systems in advanced old age. For the intestine, this includes a decrease in motility, small intestine permeability and mucosal defence (Calvani, 2018). In addition, there are associated changes in lifestyle and dietary habits, reduced somatic mobility and increased probability of hospitalisation and medication use (Han et al., 2017). There may also be deficits in memory and increases in anxiety in humans (Han et al., 2017) or mice (Scott et al., 2017). Other agerelated changes include alterations in the HPA axis with decreased negative feedback mechanisms and increased basal glucocorticoid release (Calvani et

al., 2018). Circadian rhythm disruption and altered sleep/wake cycles have been also been reported to increase in prevalence in old age (Hood and Amir, 2017; Calvani et al., 2018). The microbiome and gut to brain signalling has increasingly been focused on in the ageing research field because of the changing nature of the microbiome during the lifespan and recognition that how the intestine signals to the enteric and central nervous systems alters during ageing. Thus, to understand how the microbiome influences ageing and devise potential remedial interventions it is necessary to understand the gut to brain neural signalling pathway.

MICROBIOME

Eli Metchnikoff proposed that people in Eastern Europe live longer because they consume lactic acid bacteria (*Metchnikoff*, 1908). The link between microbiota and longevity is also supported by the observation that germfree mice live longer than conventional controls (*Dinan* and *Cryan*, 2017). This association does not mean that the intestinal microbes themselves age. However, here is a decrease in intestinal microbial diversity in the elderly with, for example, an increase in *Bifidobacteria*, but a decrease in their relative proportion (*Claesson* et al., 2011; *Dinan* and *Cryan*, 2017; *Calvani*

et al., 2018). A confounding factor is that age-related alterations in diet can lead to changes in the resident microbiome (*David* et al., 2014). It is at present not established entirely whether changes in microbiota are the cause or result of host ageing.

There is some experimental evidence that for the nematode Caenorhabditis elegans, intestinal microbes might manipulate lifespan. The prolongevity effects of metformin in C. elegans appear to depend on the suppression of folate metabolism by the worm's intestinal E. coli bacteria (Cabreiro et al., 2013). In the absence of intestinal E. coli, the C. elegans' lifespan was extended, but metformin reduced its lifespan. In addition, the effect of supplementary E. coli was dependent on the strain used; metformin resistant E. coli produce short lifespans in response to metformin, but metformin sensitive E. coli strains impaired folate metabolism and slowed ageing (*Cabreiro* et al., 2013). In a separate C. elegans model, genetic manipulation of *E. coli* to increase their secretion of the polysaccharide colanic acid increased the worm's lifespan by action on the host's mitochondrial homeostasis (increases fragmentation) and unfolded protein response. Purified colanic acid alone was sufficient to promote longevity (Han et al., 2017). Dietary colanic acid also increased lifespan of

Drosophila melanogaster (Han et al., 2017). In other experiments, it was shown that *C. elegans* fed on NO-deficient bacteria have a reduced lifespan whereas worms exposed to NO donors had extended lifespans (*Gusarov* et al., 2013). These remarkable results imply that, for *C. elegans* (and possibly for *Drosophila*); intestinal microbes have a causal effect on the host lifespan.

The resident and transient gut microbiota can also influence gut to brain signalling via neuroactive molecules that they generate and release. Intestinal bacteria can release a variety of neurotransmitters including GABA, noradrenaline, dopamine, acetylcholine and 5-HT. These neurotransmitters can cross the mucosal layer to act on neuronal fibres innervating the epithelium (Dinan and Cryan, 2017; Calvani et al., 2018). Short-chain fatty acids produced by intestinal microbes may affect the enteric nervous system or brain directly or with respect to butyrate via epigemodulation through histone netic deacetylases (Dinan and Cryan, 2017; *Calvani* et al., 2018). Reduced levels of short-chain fatty acids associated with old age have been associated with increased susceptibility to intestinal inflammatory disorders, and it has been speculated that short-chain fatty acid availability may affect human longevity (*Nagpal* et al., 2018).

INFLAMMATION

Ageing is associated with raised background inflammation (inflammageing). Inflammageing is associated with increased risks for morbidity and mortality and age-related diseases (*Franceschi* and *Campisi*, 2014). The increased background inflammation appears to be accompanied by reduced inflammatory response to acute challenges (*Franceschi* and *Campisi*, 2014) perhaps because the acute inflammatory response starts off from a higher background level to a given ceiling. A counterpoint is that chronic inflammation in the very elderly may have an adaptive role possibly underlying tissue remodelling; for example, healthy centenarians can have an increased background inflammation with associated hypercoagulability, but this may be balanced by other anti-inflammatory mechanisms (*Franceschi* and *Campisi*, 2014).

It has been reported that intestinal commensal bacteria that maintain immune tolerance tend to be reduced in aged people, but opportunistic bacteria that stimulate intestinal inflammation are often increased in number (Nagpal et al., 2018). The old age inflammatory phenotype can be manipulated experimentally. For example, when the resident microbiome was transplanted from old to young germ-free mice, there was increased inflammation in the intestines, increased leakage of bacterial components into the circulation and increased T-cell activation systemically (Fransen et al., 2017). The importance of inflammageing in gut function is illustrated by the IP injection of the tumour necrosis factor alpha antagonist etanercept in aged mice, which reversed the old age related decrease in colon serotonin transporter and slowing of intestinal motility (Patel et al., 2017).

Inflammageing is manifest in the nervous system through the activation of microglia (the tissue macrophages of the brain) and may be associated with neurodegenerative pathologies such as Parkinson's disease (Jvothi et al., 2015). The roles of microglia are complex with metabolic and neuroprotective functions, but in old age the microglial sensome can shift from being mainly protective to favouring neuroinflammation (Hickman et al., 2013). Microglia activation and/or development are influenced by the gut microbiome (Erny et al., 2015), since germfree mice have defective and immature microglia, and antibiotic treated mice also had immature microglia. Reconstitution of microbiota from donor specific pathogen free mice restored microglia function (Erny et al., 2015). The influence that the microbiome appears to have on brain microglia might explain, at least to some extent, the shift in old age to a pro-inflammatory microglial phenotype.

GUT MOTILITY

Chronic constipation, with or without incontinence, tends to increase in prevalence with old age (*Higgins* and *Johanson*, 2004; *De Giorgio* et al., 2015; *Ranson* and *Saffrey*, 2015). In north America, and after age 65 y, 16 or 26% (male vs female) have reported chronic constipation and this rose to 26 or 34% at the 84th year (*De Giorgio* et al., 2015). Beyond psychosocial and economic factors, it is believed that ageing of the gut itself can underlie these numbers.

Changes in gastrointestinal function with ageing occur in both humans and animal models and the upper and lower gut are most at risk. The oesophagus, stomach, colon, and rectum are targeted with difficulties in swallowing and defaecation; there is also an increase in intra-colonic pressure (reviewed in Hall, 2002; Wade, 2002; Soenen et al., 2016). Aged rats have fewer colonic migrating motor complexes as measured by electrodes in fasting conscious animals (Metugriachuk et al., 2006). Similarly, aged (2 y old) mice have reduced total faecal output with decreased water content in the pellets compared to their younger (3 mo) counterparts (*Patel* et al., 2012). In particular, there was a decrease in velocity of epoxy coated pellet movement in the colon and an increase in impaction (Patel et al., 2012).

Since the enteric nervous system (ENS) controls gut motility and secretion, it is reasonable to ask whether the increased prevalence of constipation in the elderly is accompanied by alterations or reductions in number of enteric neurons. Indeed, reductions in the number of myenteric neurons have been reported for the elderly. Along the gut there is an apparent 34% decrease in the number of myenteric neurons in the elderly, with the largest reduction (>38%) reported for the duodenum (*de* Souza et al., 1993). In the human colon myenteric plexus, there was a decrease in the number of choline acetyltransferase positive neurons while neuronal nitric oxide positive neurons did not decline in number (Bernard et al., 2009). Hanani et al. (2004) reported an increase with age in the proportion of colonic myenteric ganglia with cavities and a decrease in the proportion of normal ganglia.

Similar reductions in myenteric neurons have been reported for animals. Moreover, numerous animal studies suggest that the ENS is more susceptible to age-related degeneration than other nervous systems (Saffrey, 2013). Aged Fisher 344 rats showed significant reductions in the number of myenteric neurons (and associated glia) in both small and large intestines, except for the rectum (*Phillips* et al., 2004). The same was true for mice when young (3 mo) were compared to old (24 mo) ones (*El-Salhy* et al., 1999). An analogous reduction in neurons (5 vs 25 mo old) has been reported for the myenteric and submucous plexuses in the small intestine of the guinea pig (Phillips and Powley, 2007; Zanesco and *Souza*, 2011).

The enteric nervous system is a complete nervous system in the sense that it contains primary afferent, interand motor neurons. Also, peristalsis and mixing occur ex vivo after all nervous connections with the extrinsic nervous systems have been severed, demonstrating that the enteric nervous system can function independently. There are several functional classes of specialised neurons within the ENS subserving different functions (Kunze and Furness, 1999), but only one class, intrinsic primary afferent neurons (IPANs), is both chemo- and mechanosensitive and serves as an intramural gatekeeper relaying more than two thirds of signals originating from luminal contents to the afferent vagus nerve (Perez-Burgos et al., 2014). The remainder (< one third) of afferent chemoceptive vagal signals derive from direct innervation of the intestinal epithelium by the vagus. These considerations make it important to ask, with respect to gut to brain signalling, whether there are enteric neurons particularly sensitive or resistant to the effects of ageing.

Cholinergic (choline acetyltransferase (ChAT) immunopositive) enteric neurons appear to be especially vulnerable to ageing-related decrease when compared to other histochemical phenotypes (Camilleri et al., 2008). The number of ChAT positive human myenteric neurons decreased with age while nNOS neurons do not appear to change (Bernard et al., 2009). In rodents too, ChAT positive were preferentially lost (after 12 mo of age) with nicotinamideadenine dinucleuotide phosphate diaphorase positive neurons (nNOS containing) being spared (Phillips et al., 2003; Phillips and Powley, 2007). Intriguingly, this loss of enteric cholinergic neurons is paralleled by a preferential decrease in cholinergic innervation in the aged brain (Casu 2002). However, et al., enteric cholinergic neurons belong to several functional classes including IPANs, excitatory motor neurons and interneurons, and the question arises which of these classes are the most vulnerable.

It has been suggested that IPANs are most susceptible to neurodegeneration with age (Wade, 2002; Wade and Cowen, 2004), exhibiting degeneration to a greater extent than other phenotypes, e.g. serotonergic interneurons or nitrergic inhibitory motor neurones. The evidence for this is incomplete, relying mainly on the decrease in old age of vitamin D-dependent 28 kDa calcium binding protein (calbindin) positive neurons or the decrease in substance P positive neurites in the guineapig or rodent myenteric plexus (*Wade*, 2002; Wade and Cowen, 2004). Because IPANs are positive for substance P, choline acetyltransferase (*Phillips* et al., 2003) and calbindin there is at least circumstantial evidence that IPANs may be principally sensitive to the effects of ageing.

Ageing also affects the neurites (processes) of myenteric neurons. Agerelated development of dystrophic nerve fibers alters the shape of human (*Hanani* et al., 2004) and guinea pig enteric ganglia (Abalo et al., 2007), presumably due to the presence of swollen fibres. In aged mice (>18 mo old) calbindin containing and nNOS immunoreactive neurons developed multiple swollen processes when compared to young mice (3 mo) (Gamage et al., 2013). Other signs of degeneration and pathology include the accumulation in guinea pig or rat of lipofuscin (pigment granules composed of lipid-containing residues of lysosomal digestion) (*Phillips* et al., 2004; Abalo et al., 2007) and the presence of α -synuclein immunoreactive aggregates and hyperphosphorylated microtubule-associated Tau protein (Phillips et al., 2009). It should be noted that glial cells and interstitial cells of Cajal also decrease with age, although neuronal reduction seems to occur first (Phillips and Powley, 2007; Camilleri et al., 2008; Saffrey, 2013, 2014).

Despite neural loss and degeneration, the ageing gut and the enteric nervous system have a significant functional reserve since intestinal motility, although reduced, appears largely intact until the animal is very old (*Saffrey*, 2013).

EXTRINSIC AFFERENTS

The intestine is innervated by nerve fibres whose cell bodies lie outside the gut wall. These fibres run within mesenteric neurovascular bundles that branch off arterial arcades supplying intestinal segments. The nerve bundles contain vagal, and spinal and sympathetic nerve fibres.

Single unit (extracellular action potentials from single nerve fibres) mesenteric nerve fibre recordings have been made using *ex vivo* segments of human ileum or sigmoid colon (*Yu* et al., 2016). For these recordings, background firing rates correlated negatively with age (from 24 to 77 y old), and number of single units showing burst firing patterns also decline with age. There was also a decrease in the number of substance P containing nerve fibres (presumably either intrinsic or extrinsic) innervating the luminal mucosa (Yu et al., 2016). Paradoxically, there was also an increase in the density of mucosal mast cells and ileal enterochromaffin cell numbers with age, which the authors attributed to a compensatory mechanism for the sensory neurodegeneration (Yu et al., 2016). These results parallel others that report decreased intestinal visceral pain perception in old age (Lasch et al., 1997; Lagier et al., 1999).

The diminished mesenteric nerve fibre discharge seen in elderly humans has been replicated in ageing mice. Single and multiunit spike recordings have been made from the mesenteric nerve of young and aged (3 mo vs 12 and 24 mo old) C57bl/6 mice (Keating et al., 2016). Baseline mesenteric multiunit spiking activity was significantly decreased for both jejunal and colonic mesenteric afferent fibres in old (24 mo) compared to young (3 mo) mice. Mesenteric nerve fibre responses to intraluminal fluid distension were also reduced in old mice. The TRPV1 receptor is also a mechanoreceptor in the intestine mediating the response of mesenteric afferents to fluid distension (Rong et al., 2004). It is therefore of interest that intestinal TRPV1 receptor expression was diminished in the older mice (*Keating* et al., 2016).

There appears to be little functional data on the effects of ageing on the afferent vagus compared to the mixed mesenteric nerve. There are, however, reports in the literature that the vagus nerve of old animals displays degenerative anatomical changes. In aged rats, vagal afferents appear to have swollen varicosities in fibres innervating the myenteric plexus, smooth muscle and mucosa (Phillips and Powley, 2007). It is not established that there is an actual decrease in the number of vagal fibre endings supplying the myenteric plexus but there are dystopic changes including dilations and swellings (*Phillips* et al., 2010) in the NIH Fisher 344 rat model of ageing. The extent of the terminal arbors is also reduced compared to young rats. Similar degenerative changes are seen in vagal afferents to mucosal villi and the musculature (*Phillips* et al., 2010). It is thought that this reduction or the pathological changes in vagal innervation results from a decrease in trophic factors released from the targets of vagal innervation (Phillips et al., 2010) in particular NT-4 (*Phillips* et al., 2010).

Sympathetic nerve fibres originate from prevertebral sympathetic ganglia to innervate the ENS, intestinal arteries and smooth muscle (Phillips and Powley, 2007). Tyrosine hydroxylase staining can be used to identify the sympathetic innervation of the intestine, and positive fibres that supply the myenteric and submucous plexuses of aged (24 mo old) rats display axonopathies such as swollen axons and small sparse terminals (*Phillips* and *Powley*, 2007). Functionally, old age has been associated with decreased ability of the sympathetic nervous system to adapt to environmental or interoception of stimuli (Hotta and Uchida, 2010).

INTRAMURAL GATEKEEPER

90 to 95% of sensory neuron processes innervating the intestinal epithelium arise from the ENS, with the rest originating from neurons whose somata are located outside the intestine (*Keast* et al., 1984; *Ekblad* et al., 1987). In agreement with this anatomical data on epithelial innervation density, is the recent discovery that more than two thirds of vagal afferent signals evoked by a luminal probiotic is relayed to the vagus via the enteric neurons (*Perez-Burgos* et al., 2014). Neuroactive luminal molecules first excite juxtaepithelial neurites belonging to IPANs whose cell bodies are located within



Figure 1: Gut to brain afferent signalling pathway. Sensory neurons within the enteric nervous system respond to luminal stimuli and relay information to the brain via a nicotinic sensory synapse that activates the afferent vagus.

the enteric nervous system. The excited IPANs release acetylcholine and perhaps other neurotransmitters to activate vagal intraganglionic laminar endings (IGLEs) which closely surround and abut the IPANs (Berthoud et al., 1997; Perez-Burgos et al., 2014). The IPAN to IGLE nicotinic sensory synapse is perfectly positioned to act as a gatekeeper to regulate microbial to brain signalling (Figure 1). Accordingly, the amount of information transmitted to the brain via the vagus would be markedly influenced by whether IPANs are refractory or readily responsive to luminal stimuli, and by the density of IPAN sensory innervation of the epithelium.

Given the important role of enteric IPANs in gut to brain signalling, the vulnerability of the enteric nervous system to old age in terms of numbers and degeneration would have a significant impact on the amount and quality of information reaching the brain from the gut. Thus, even if the number of vagal afferent fibres are not appreciably reduced it can be predicted that there would be decreased constitutive and stimulus evoked vagal afferent responses to luminal microbial stimuli in old age. Exacerbating these potential impediments in gut to brain signalling there is the added problem of decreased microbial diversity (Nagpal et al., 2018) demonstrated in old age. In summary, there are several reasons why gut microbes to brain signalling via the vagus is compromised in old age, with a degenerating or dysfunctional enteric nervous system being a major contributor to the reduced signalling.

LITERATURE

- Abalo, R., Vera, G., Rivera, A.J., and Martin, M.I.: Age-related changes in the gastrointestinal tract: a functional and immunohistochemical study in guinea-pig ileum. Life Sci. 80, 2436-2445 (2007).
- Bernard, C.E., Gibbons, S.J., Gomez-Pinilla, P.J., Lurken, M.S., Schmalz, P.F., Roeder, J.L., Linden, D., Cima, R.R., Dozois, E.J., Larson, D.W., Camilleri, M., Zinsmeister, A.R., Pozo, M.J., Hicks, G.A., and Farrugia, G.: Effect of age on the enteric nervous system of the human colon. Neurogastroenterol. Motil. 21, 746-e746 (2009).
- Berthoud, H.R., Patterson, L.M., Neumann, F., and Neuhuber, W.L.: Distribution and structure of vagal afferent intraganglionic laminar endings (IGLEs) in the rat gastrointestinal tract. Anat. Embryol. (Berl.) 195, 183-191 (1997).
- Cabreiro, F., Au, C., Leung, K.Y., Vergara-Irigaray, N., Cocheme, H.M., Noori, T., Weinkove, D., Schuster, E., Greene, N.D., and Gems, D.: Metformin retards aging in C. elegans by altering microbial folate and methionine metabolism. Cell 153, 228-239 (2013).
- Calvani, R., Picca, A., Lo Monaco, M.R., Landi, F., Bernabei, R., and Marzetti E.: Of Microbes and Minds: A Narrative Review on the Second Brain Aging. Front. Med. (Lausanne) 5, 53 (2018).
- Camilleri, M., Cowen, T., and Koch, T.R.: Enteric neurodegeneration in ageing. Neurogastroenterol. Motil. 20, 418-429 (2008).
- Casu, M.A., Wong, T.P., De Koninck, Y., Ribeiro-da-Silva, A., and Cuello, A.C.: Aging causes a preferential loss of cholinergic innervation of characterized neocortical pyramidal neurons. Cereb. Cortex 12, 329-337 (2002).
- Claesson, M.J., Cusack, S., O'Sullivan, O., Greene-Diniz, R., de Weerd, H., Flannery, E., Marchesi, J.R., Falush, D., Dinan, T., Fitzgerald, G., Stanton, C., van Sinderen, D., O'Connor, M., Harnedy, N., O'Connor, K., Henry, C., O'Mahony, D., Fitzgerald, A.P., Shanahan, F., Twomey, C., Hill, C.,

Ross, R.P., and O'Toole, P.W.: Composition, variability, and temporal stability of the intestinal microbiota of the elderly. Proc. Natl. Acad. Sci. USA 108, Suppl. 1, 4586-4591 (2011).

- David, L.A., Maurice, C.F., Carmody, R.N., Gootenberg, D.B., Button, J.E., Wolfe, B.E., Ling, A.V., Devlin, A.S., Varma, Y., Fischbach, M.A., Biddinger, S.B., Dutton, R.J., and Turnbaugh, P.J.: Diet rapidly and reproducibly alters the human gut microbiome. Nature 505, 559-563 (2014).
- De Giorgio, R., Ruggeri, E., Stanghellini, V., Eusebi, L.H., Bazzoli, F., and Chiarioni, G.: Chronic constipation in the elderly: a primer for the gastroenterologist. BMC Gastroenterol. 15, 130 (2015).
- de Souza, R.R., Moratelli, H.B., Borges, N., and Liberti, E.A.: Age-induced nerve cell loss in the myenteric plexus of the small intestine in man. Gerontology 39, 183-188 (1993).
- Dinan, T.G. and Cryan, J.F.: Gut instincts: microbiota as a key regulator of brain development, ageing and neurodegeneration. J. Physiol. 595, 489-503 (2017).
- Ekblad, E., Winther, C., Ekman, R., Hakanson, R., and Sundler, F.: Projections of peptidecontaining neurons in rat small intestine. Neuroscience 20, 169-188 (1987).
- El-Salhy, M., Sandstrom, O., and Holmlund, F.: Age-induced changes in the enteric nervous system in the mouse. Mech. Ageing Dev. 107, 93-103 (1999).
- Erny, D., Hrabě de Angelis, A., Jaitin, D., Wieghofer, P., Staszewski, O., David, E., Keren-Shaul, H., Mahlakoiv, T., Jakobshagen, K., Buch, T., Schwierzeck, V., Utermöhlen, O., Chun, E., Garrett, W., McCoy, K., Diefenbach, A., Staeheli, P., Stecher, B., Amit, I., and Prinz, M.: Host microbiota constantly control maturation and function of microglia in the CNS. Nat. Neurosci. 18, 965-977 (2015).
- Franceschi, C. and Campisi, J.: Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. J.

Gerontol. A Biol. Sc.i Med. Sci. 69, Suppl. 1, S4-9 (2014).

- Fransen, F., van Beek, A.A., Borghuis, T., Aidy, S.E., Hugenholtz, F., van der Gaastde Jongh, C., Savelkoul, H.F.J., De Jonge, M.I., Boekschoten, M.V., Smidt, H., Faas, M.M., and de Vos, P.: Aged Gut Microbiota Contributes to Systemical Inflammaging after Transfer to Germ-Free Mice. Front. Immunol. 8, 1385 (2017).
- Gamage, P.P., Ranson, R.N., Patel, B.A., Yeoman, M.S., and Saffrey, M.J.: Myenteric neuron numbers are maintained in aging mouse distal colon. Neurogastroenterol. Motil, 25, e495-e505 (2013).
- Gusarov, I., Gautier, L., Smolentseva, O., Shamovsky, I., Eremina, S., Mironov, A., and Nudler, E.: Bacterial Nitric Oxide Extends the Lifespan of C. elegans. Cell 152, 818-830 (2013).
- Hall, K.E.: Aging and neural control of the GI tract. II. Neural control of the aging gut: can an old dog learn new tricks? Am. J. Physiol. Gastrointest. Liver Physiol 283, G827-832 (2002).
- Han, B., Sivaramakrishnan, P., Lin, C.-C.J., Neve, I.A.A., He, J., Tay, L.W.R., Sowa, J.N., Sizovs, A, Du, G., Wang, J., Herman, C., and Wang, M.C.: Microbial Genetic Composition Tunes Host Longevity. Cell 169, 1249-1262 (2017).
- Hanani, M., Fellig, Y., Udassin, R., and Freund, H.R.: Age-related changes in the morphology of the myenteric plexus of the human colon. Auton. Neurosci. 113, 71-78 (2004).
- Hickman, S.E., Kingery, N.D., Ohsumi, T.K., Borowsky, M.L., Wang, L.C., Means, T.K., and El Khoury, J.: The microglial sensome revealed by direct RNA sequencing. Nat. Neurosci. 16, 1896-1905 (2013).
- Higgins, P.D. and Johanson, J.F.: Epidemiology of constipation in North America: a systematic review. Am. J. Gastroenterol. 99, 750-759 (2004).
- Holzer, P. and Farzi, A.: Neuropeptides and the microbiota-gut-brain axis. Adv. Exp. Med. Biol. 817, 195-219 (2014).
- Hood, S. and Amir, S.: The aging clock:

circadian rhythms and later life. J. Clin. Invest. 127, 437-446 (2017).

- Hotta, H. and Uchida, S.: Aging of the autonomic nervous system and possible improvements in autonomic activity using somatic afferent stimulation. Geriatr. Gerontol. Int. 10, Suppl 1, S127-136 (2010).
- Jyothi, H.J., Vidyadhara, D.J., Mahadevan, A., Philip, M., Parmar, S.K., Manohari, S.G., Shankar, S.K., Raju, T.R., and Alladi, P.A.: Aging causes morphological alterations in astrocytes and microglia in human substantia nigra pars compacta. Neurobiol. Aging 36, 3321-3333 (2015).
- Keast, J.R., Furness, J.B., and Costa, M.: Origins of peptide and norepinephrine nerves in the mucosa of the guinea pig small intestine. Gastroenterology 86, 637-644 (1984).
- Keating, C., Nocchi, L., Yu, Y., Donovan, J., and Grundy, D.: Ageing and gastrointestinal sensory function: Altered colonic mechanosensory and chemosensory function in the aged mouse. J Physiol. 594, 4549-4564 (2016).
- Kunze, W.A. and Furness, J.B.: The enteric nervous system and regulation of intestinal motility. Annu. Rev. Physiol. 61, 117-142 (1999).
- Lagier, E., Delvaux, M., Vellas, B., Fioramonti, J., Bueno, L., Albarede, J.L., and Frexinos, J.: Influence of age on rectal tone and sensitivity to distension in healthy subjects. Neurogastroenterol. Motil. 11, 101-107 (1999).
- Lasch, H., Castell, D.O., and Castell, JA.: Evidence for diminished visceral pain with aging: studies using graded intraesophageal balloon distension. Am. J. Physiol. 272, G1-3 (1997).
- Metchnikoff, E: The Prolongation of Life: Optimistic Studies. English translation, edited by Peter Chalmers Mitchell. G.P. Putnam's Sons, New York & London, (1908).
- Metugriachuk, Y., Marotta, F., Pavasuthipaisit, K., Kuroi, O., Tsuchiya, J., Lorenzetti, A., Fesce, E., and Minelli, E.: The aging gut motility decay: may symbiotics be acting as "implantable" biologic pace-makers? Rejuvenation Res. 9, 342-345 (2006).

- Nagpal, R., Mainali, R., Ahmadi, S., Wang, S., Singh, R., Kavanagh, K., Kitzman, D.W., Kushugulova, A., Marotta, F., and Yadav, H.: Gut microbiome and aging: Physiological and mechanistic insights. Nutr. Healthy Aging, 4, 267-285 (2018).
- Patel, B., Patel, N., Fidalgo, S., Ranson, R., Saffrey, J., and Yeoman, M.: Age-related changes in colonic motility, faecal output and the properties of faecal pellets in the mouse. Neurogastroenterol. Motil. 24, 129 (2012).
- Patel, B.A., Fidalgo, S., Wang, C., Parmar, L., Mandona, K., Panossian, A., Flint, M.S., Ranson, R.N., Saffrey, M.J., and Yeoman, M.S.: The TNF-α antagonist etanercept reverses age-related decreases in colonic SERT expression and faecal output in mice. Sci. Rep. 7, 42754 (2017).
- Perez-Burgos, A., Mao, Y., Bienenstock, J., and Kunze, W.: The gut-brain axis rewired: adding a functional vagal nicotinic "sensory synapse". FASEB J. 28, 3064-3074 (2014).
- Phillips R, Walter G, Ringer B, Higgs K and Powley T. (2009). Alpha-synuclein immunopositive aggregates in the myenteric plexus of the aging Fischer 344 rat. Exp Neurol 220, 109-119.
- Phillips, R.J., Kieffer, E.J., and Powley, T.L.: Aging of the myenteric plexus: neuronal loss is specific to cholinergic neurons. Auton. Neurosci. 106, 69-83 (2003).
- Phillips, R.J., Kieffer, E.J., and Powley, T.L.: Loss of glia and neurons in the myenteric plexus of the aged Fischer 344 rat. Anat. Embryol. (Berl.) 209, 19-30 (2004).
- Phillips, R.J. and Powley, T.L.: Innervation of the gastrointestinal tract: patterns of aging. Auton. Neurosci. 136, 1-19 (2007).
- Phillips, R.J., Walter, G.C., and Powley, T.L.: Age-related changes in vagal afferents innervating the gastrointestinal tract. Auton. Neurosci. 153, 90-98 (2010).

Ranson, R.N. and Saffrey, M.J.: Neurogenic

mechanisms in bladder and bowel ageing. Biogerontology 16, 265-284 (2015).

- Rong, W., Hillsley, K., Davis, J.B., Hicks, G., Winchester, W.J., and Grundy, D.: Jejunal afferent nerve sensitivity in wild-type and TRPV1 knockout mice. J. Physiol. 560, 867-881 (2004).
- Saffrey, M.J.: Cellular changes in the enteric nervous system during ageing. Dev. Biol. 382, 344-355 (2013).
- Saffrey, M.J.: Aging of the mammalian gastrointestinal tract: a complex organ system. Age (Dordr.) 36, 9603 (2014).
- Scott, K.A., Ida, M., Peterson, V.L., Prenderville, J.A., Moloney, G.M., Izumo, T., Murphy, K., Murphy, A., Ross, R.P., Stanton, C., Dinan, T.G., and Cryan, J.F.: Revisiting Metchnikoff: Age-related alterations in microbiota-gut-brain axis in the mouse. Brain Behav. Immun. 65, 20-32 (2017).
- Soenen, S., Rayner, C., Jones, K., and Horowitz, M.: The ageing gastrointestinal tract. Curr. Opin. Clin. Nutr. Metab. Care 19, 12-18 (2016).
- Wade, P.R.: Aging and neural control of the GI tract. I. Age-related changes in the enteric nervous system. Am. J. Physiol. Gastrointest. Liver Physiol. 283, G489-495 (2002).
- Wade, P.R. and Cowen, T.: Neurodegeneration: a key factor in the ageing gut. Neurogastroenterol. Motil. 16, Suppl. 1, 19-23 (2004).
- Yu, Y., Daly, D., Adam, I., Kitsanta, P., Hill, C., Wild, J., Shorthouse, A., Grundy, D., and Jiang, W.: Interplay between mast cells, enterochromaffin cells, and sensory signaling in the aging human bowel. Neurogastroenterol. Motil. 28, 1465-1479 (2016).
- Zanesco, M. and Souza, R.: Morphoquantitative study of the submucous plexus (of Meissner) of the jejunum-ileum of young and old guinea pigs. Arq. Neuropsiquiatr. 69, 85-90 (2011).

STROKE AND MICROBIOTA – CURRENT RESEARCH AND CONCEPTS

JOSEF ANRATHER

Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY, USA

SUMMARY

Stroke is a highly prevalent cerebrovascular disease with limited treatment options. Although therapeutic interventions have evolved over the last 30 years, stroke remains the second most common cause of death and the leading cause for long-term disability worldwide. While primarily a disease caused by acute ischaemia to the brain, stroke pathology is characterized by a strong immune component. The intestinal microbiota and gut immune system have recently been linked to the inflammatory processes observed after stroke. It has been shown that intestinal microbiota composition strongly affects the polarization state of immune cells found in Payer's patches and submucosa of the small intestine. This may result in an imbalance of regulatory and effector type T-cells in the gut. Because gut immune cells traffic to the brain after stroke they influence the local immune response triggered by ischaemic injury by damping or increasing inflammation. In the context of stroke, the communication between brain and gut is bidirectional and stroke can impact the composition of intestinal microbiota. In addition, intestinal microbiota can become a source of opportunistic infectious agents and contribute to post-stroke infections thereby increasing morbidity and mortality. In this review I will summarize current concepts on the role of intestinal microbiota and the intestinal immune system in stroke, review the effects stroke and ageing has on the gut microbiome, and describe studies that identified commensal intestinal microbiota as a source of infectious bacteria participating in post-stroke infections of the respiratory tract.

INTRODUCTION

Ischaemic stroke is a highly prevalent and devastating disease with limited therapeutic options (*Henninger* et al., 2010). Most therapeutic efforts have focused on counteracting the intrinsic brain mechanisms leading to ischaemic injury (*Suwanwela* and *Koroshetz*, 2007). Unfortunately, these efforts have met with limited success in large clinical trials, reflecting, in part, an incomplete understanding of the pathological processes triggered by cerebral ischaemia (*Moskowitz* et al., 2010). Inflammation is a key component in the pathophysiology of ischaemic stroke (*Iadecola* and *Anrather*, 2011). Numerous experimental approaches have concentrated on developing the therapeutic potential of immunomodulation (*Macrez* et al., 2011). However, our understanding of the interaction between resident brain cells and peripheral immune cells infiltrating the postischaemic brain, and their role in tissue damage and repair is still incomplete (*Macrez* et al., 2011). The peripheral

immune system plays an essential role in the pathophysiology of stroke, involving both innate and adaptive immune cells (Iadecola and Anrather, 2011). In turn, neuro-humoral brain signals generated during stroke induce immunosuppression, which contributes to post-stroke morbidity by increasing the risk for infection in stroke patients (Meisel et al., 2005). The immune system is forged by the continuous interaction with commensal microbiota that populate the epithelial surfaces and this interaction is essential for immune cell development, maintenance and function (Mazmanian et al., 2005; Hill and Artis, 2010). Specifically, intestinal commensal bacteria, the most abundant

symbiotic compartment in the body, have emerged as a potent regulator of the immune system (Nishio and Honda, 2012). In particular, the microbiota exerts powerful effects on lymphocyte populations, such as regulatory T-cells (Treg) and $\gamma\delta$ T-cells, which are involved in the mechanisms of cerebral ischaemic injury (Iadecola and Anrather, 2011). Conversely, stroke also affects the peripheral immune system and alters the composition of the intestinal microbiome. This sets the stage for complex multidirectional interactions within the microbiota-gutbrain axis that is likely to affect stroke outcome.

EFFECTS OF MICROBIOTA ON IMMUNE CELLS INVOLVED IN STROKE PATHOPHYSIOLOGY

Accumulating evidence suggests that commensal microbes influence the outcome of diseases in which the immune system is involved. For example, germfree (GF) animals are protected from experimental autoimmune encephalomyelitis (Ochoa-Repáraz et al., 2009; Berer et al., 2011; Lee et al., 2011), ankylosing spondylitis (Sinkorová et al., 2008), or rheumatoid arthritis (Wu et al., 2010; Chappert et al., 2013). Increasing evidence implicates systemic immunity in the pathophysiology of ischaemic brain injury (Iadecola and Anrather, 2011). Thus, cells associated with innate and adaptive immunity play a defining role in the outcome of cerebral ischaemia (*Iadecola* and *Anrather*, 2011). As reviewed below, intestinal microbes are critical for the development, expansion and activity of immune cells implicated in stroke pathophysiology.

Lymphocytes

Th1 and IL-17 secreting T-cells are detrimental in the early phase of cere-

bral ischaemia and lymphocyte-deficient mice are protected in models of focal ischaemia (Hurn et al., 2007; Kleinschnitz et al., 2010). The mechanism of T-cell mediated brain injury after ischaemia does not involve classical antigen-mediated T-cell activation and the cytotoxic activity might be related innate T-cell functions to (Kleinschnitz et al., 2010). Accordingly, IL-17 secreting $\gamma\delta$ T-cells, that do not undergo classical antigen-dependent T-cell activation, have been shown to contribute to ischaemic injury (Shichita et al., 2009; Gelderblom et al., 2012;). Peripheral development of T-cells is greatly affected by microbiota (*Kuhn* and *Stappenbeck*, 2013). Beside Th17 cells (*Ivanov* et al., 2006), intestinal $\gamma\delta$ T-cells residing in the epithelial layer and submucosa (lamina propria) are also a major source for IL 17 (Roark et al., 2008; Korn and Petermann, 2012; Hou et al., 2013). IL-17producing $\gamma \delta T$ -cells are responsible for the colitis seen in Treg deficient mice and the disease is abrogated in $\gamma\delta T$ -

cell-deficient or antibiotic-treated mice, suggesting activation of $\gamma\delta T$ -cells by commensal bacteria (Park et al., 2010). Moreover, intestinal IL-17-producing $\gamma\delta$ T-cells are reduced in antibiotictreated mice (*Duan* et al., 2010; *Park* et al., 2010). While effector T-lymphocytes may contribute to focal ischaemic injury (Yilmaz et al., 2006; Hurn et al., 2007), Treg can have a protective effect by down-regulating post-ischaemic inflammation. Treg appear in the ischaemic tissue after the acute phase and confer neuroprotection by IL-10 secretion, an effect that might be antigen independent (Liesz et al., 2009, 2013; Planas and Chamorro, 2009; Stubbe et al., 2013). Intestinal Treg express high levels of IL-10, which is indispensable for maintaining intestinal homeostasis by suppressing Th17 differentiation (Chaudhry et al., 2009; Huber et al., 2011), yoT-cell proliferation (Park et al., 2010), and myeloid cell activation (Takeda et al., 1999). The gut microbiota has a role also in the development of intestinal and extraintestinal Treg. For example, colonic FoxP3⁺ Treg cells are reduced in GF mice (Round and Mazmanian, 2010; Atarashi et al., 2013). Furthermore, in a mouse model of EAE, the generation of Treg from splenic and lymph node lymphocytes was increased in GF animals (Lee et al., 2011). Therefore, mucosa-microbial interactions influence Treg lymphocytes in the gut as in lymphoid organs.

Myeloid cells

In experimental stroke, as in human stroke, monocytes/macrophages infil-

trate the brain early after ischaemia and persist for several weeks (Schilling et al., 2003; Gelderblom et al., 2009). Inhibition of post-ischaemic monocyte recruitment by interfering with CCR2 and CX3CR1 chemokine receptor utilization is beneficial in experimental stroke (Hughes et al., 2002; Soriano et al., 2002; Denes et al., 2008; Schilling et al., 2009). Monocyte/macrophage development is influenced by signals Systemic intestinal bacteria. from monocyte counts are normal in GF mice, but they are functionally impaired and do not mount an IFNy response after bacterial stimulation (Ganal et al., 2012). Dendritic cells (DC) have been implicated in stroke pathology (Felger et al., 2010; Gelderblom et al., 2017). Intestinal DC and macrophages are essential to maintain a tolerogenic environment in the gut and are strong inducers of Treg (Coombes et al., 2007; Sun et al., 2007). Neutrophils play a part in post-ischaemic inflammation by limiting tissue perfusion due to intravascular clogging (*del* Zoppo et al., 1991; Dawson et al., 1996), destabilizing the BBB by releasing MMP9 (Rosell et al., 2008; Ludewig et al., 2013), and by generating ROS and NO (Garcia-Bonilla et al., 2014). Neutrophils isolated from the bone marrow of GF or antibiotictreated mice exhibit impaired ex vivo bacterial killing (Clarke et al., 2010). All together, there is ample evidence that the intestinal environment shapes the activation state of immune cells involved in the immune response to ischaemic brain injury.

ALTERED INTESTINAL IMMUNE STATUS AND ITS EFFECTS ON ISCHAEMIC STROKE

The role of altered intestinal microbial composition on stroke outcome has been investigated in mice undergoing transient focal ischaemia by intraluminal obstruction of the middle cerebral artery (MCA) (*Benakis* et al.,

2016). In this study mice were treated with amoxicillin (amoxicillin/clavulanic acid; AC) to alter the microbiome resulting in elimination of Clostridia and *Bacteroidetes* with concomitant expansion of *Proteobacteria*. The authors observed remarkable neuroprotection both on the anatomical and functional level. Because protection was not observed after one week of AC treatment, a time point when microbial alterations were already present, let the authors to hypothesize that changes in the immune system leading to an altered inflammatory response after stroke might be at the basis of the observed neuroprotection. Analysis of the intestinal immune system revealed an increase in Treg and a decrease in IL17⁺ $\gamma\delta$ T-cells, whereas Th17 cells were not affected. In addition, the authors could show that DC isolated from AC treated animals showed higher expression of the "tolerogenic" marker protein CD103 and were more potent in inducing Treg when co-cultured with naïve CD4⁺ T-cells. After MCA occlusion IL17⁺ $\gamma\delta$ T-cells were increased in the meninges of control but not ACtreated mice indicating that the deleterious IL17 response was blocked in mice with altered microbiota. Accordingly, the protective effect of AC treatment was absent in IL17 deficient mice. Similar alterations in T-cell homeostasis resulting in reduced Treg and increased Th1 and Th17 cells were observed after ischaemic stroke in mice and post-stroke immune changes could be prevented by faecal matter transplants from healthy donors (Singh et al., 2016). It was also shown that Tcells trafficked from the intestine to the brain after stroke indicating that the polarization state of intestinal T-cells might determine the course of post-ischaemic inflammation in the brain (Benakis et al., 2016; Singh et al., 2016). A recent study by Singh and coworkers (Singh et al., 2018) compared GF animals with GF animals recolonized with conventional SPF flora in a permanent focal ischaemia model in mice. Interestingly, GF animals showed increased infarct volume and decreased inflammatory response to cerebral ischaemia. Using Rag1^{-/-} mice that lack T and B cells, the authors found that ischaemic lesions in GF Rag1^{-/-} mice were not different from lesions of recolonized Rag^{-/-} mice, albeit the lesions were smaller than in wild-type mice. These studies identified disturbances of the intestinal immune homeostasis as an important contributor to ischaemic brain injury.

EFFECTS OF STROKE ON THE INTESTINAL MICROBIOME

The relationship between microbiota and stroke is complex, given that ischaemic brain injury has been found to alter gut microbiota composition. After stroke, up to 50% of patients experience gastrointestinal symptoms, including dysphagia, gastrointestinal bleeding, and constipation. Stroke is likely to alter the intestinal microbial environment by altering intestinal epithelial permeability, motility, mucus biosynthesis and the immune system. Large ischaemic infarcts in mice after transient MCA occlusion altered the gut microbiome partially by inducing intestinal paralysis leading to reduced microbial diversity and overgrowth of *Bacteroidetes* species (*Singh* et al., 2016). Another study found that after stroke, relative abundance of *Peptococcaceae* increased in the mouse caecum, whereas the proportion of *Prevotellacea* decreased (*Houlden* et al., 2016). This alteration was paralleled by augmented release of noradrenaline and the reduction of mucoprotein-producing goblet cells. Investigating the mucosa-associated microbiome it was found that microbial communities within the mucosa were significantly different between sham-operated and post-stroke mice at 24 hours following surgery (*Stanley* et al., 2018). Microbiota composition was substantially different in all sections of the gastrointestinal mucosa. The main changes in mucosal microbiota composition were due to increased abundance of *Akkermansia muciniphila* and clostridial species while operational taxonomic units (OTUs) potentially belonging to the *Barnesiella* genus were reduced.

ROLE OF INTESTINAL MICROBIOTA IN POST-STROKE INFECTIONS

Secondary infections, including pneumonia and urinary tract infections, are major complications in stroke patients. Pneumonia occurring in up to 20% of all stroke patients is the most common serious medical complication in stroke care resulting in a 2.5-fold increased mortality rate (Meisel et al., 2005). Post-stroke immunodepression has been recognized as a key factor in facilitating infections. Immunodepression in stroke patients manifests itself in reduced peripheral blood lymphocyte counts and impaired T- and natural killer (NK)-cell activity (Meisel et al., 2005). Experimental stroke in rodents induced a rapid and extensive apoptotic loss of lymphocytes in lymphoid organs and peripheral blood (Prass et al., 2003). The effects were linked to increased noradrenalin and glucocorticoid secretion through overactivation of the hypothalamic-pituitary-adrenal axis. Moreover, stroke also reduced intestinal lymphocyte counts likely resulting in suppression of the intestinal immune system and possibly facilitating bacterial invasion (Schulte-Herbrüggen et al., 2009). Further it has been shown that the majority of bacteria detected in stroke patients who developed infections were common commensal bacteria that normally reside in the intestinal tract including Escherichia coli, Enterococcus spp., and Morganella morganii (Stanley et al., 2016). Using the transient MCA occlusion model in mice, the same study found that loss of intestinal epithelial barrier function, which was observed as early as 3 hours after stroke, facilitated the translocation of facultative pathogenic bacteria into the blood stream from where they may have colonized the respiratory tract. Using bioinformatic algorithms it was predicted that the small intestine and liver were the most likely origins of microbial communities present in the lung of poststroke mice.

STROKE, MICROBIOME AND AGEING

Age has a strong effect on stroke outcome and the composition of the intestinal microbiome. Aged (18-20 month) mice show increased *Firmicutes* and reduced *Bacteroidetes* (*Spychala* et al., 2018). The ratio of *Firmicutes* to *Bac*- *teroidetes* (F:B) increased 9-fold compared to young (8-12 week) mice. Interestingly, stroke increased the F:B ratio in aged as well young mice. When young mice were transplanted with faecal microbiota harvested from aged mice, stroke outcome was worse. On the other hand, aged mice transplanted with a "young microbiome" performed better than control mice. It was also found that short chain fatty acids, potential mediators of neuroprotective effects afforded by commensal microbiota, were decreased in faecal matter of aged mice and levels could be recovered after faecal transplantation. Aged mice are also more likely to develop sepsis after cerebral ischaemia than young animals (*Crapser* et al., 2016). While high bacterial burden was detected in the mesenteric lymph nodes and spleens of both young and aged mice after stroke, substantial bacterial colonization of liver and lung was found only in aged stroke mice and not in those from young. Increased bacterial dissemination in aged mice was associated with increased gut permeability, lymphopenia, and increased plasma levels of inflammatory markers as compared to young mice.

CONCLUSIONS

Increasing evidence points to a role of intestinal microbiota in stroke pathophysiology and outcome. Recent studies have identified a gut-brain immune axis as an essential component of the inflammatory response to stroke. The microbiota-gut-brain axis is complex and shows a strong bi-directional interaction. Stroke disturbs intestinal motility and epithelial barrier function and alters the composition of intestinal microbiota. Therefore, it is likely that stroke itself induced changes of the microbial environment contribute to stroke pathology in the subacute and chronic stages of the disease. The current research also evokes many questions. Is the presence of certain microbiota a risk factor for stroke? Could one predict stroke outcome based on gut microbial composition? Besides the immune axis, are there humoral or neural pathways engaged in microbiotabrain communications that will affect stroke? Future studies are likely to address some of these questions.

LITERATURE

- Atarashi, K., Tanoue, T., Oshima, K., Suda, W., Nagano, Y., Nishikawa, H., Fukuda, S., Saito, T., Narushima, S., Hase, K., Kim, S., Fritz, J.V., Wilmes, P., Ueha, S., Matsushima, K., Ohno, H., Olle, B., Sakaguchi, S., Taniguchi, T., Morita, H., Hattori, M., and Honda, K.: Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. Nature 500, 232-236 (2013).
- Benakis, C., Brea, D., Caballero, S., Faraco, G., Moore, J., Murphy, M., Sita, G., Racchumi, G., Ling, L., Pamer, E.G., Iadecola, C., and Anrather, J.: Commensal microbiota affects ischemic stroke outcome by regulating intes-

tinal $\gamma\delta$ T cells. Nat. Med. 22, 516-523 (2016).

- Berer, K., Mues, M., Koutrolos, M., Rasbi, Z.A., Boziki, M., Johner, C., Wekerle, H., and Krishnamoorthy, G.: Commensal microbiota and myelin autoantigen cooperate to trigger autoimmune demyelination. Nature 479, 538-541 (2011).
- Chappert, P., Bouladoux, N., Naik, S., and Schwartz, R.H.: Specific gut commensal flora locally alters T cell tuning to endogenous ligands. Immunity 38, 1198-1210 (2013).
- Chaudhry, A., Rudra, D., Treuting, P., Samstein, R.M., Liang, Y., Kas, A., and Ruden-

sky, A.Y.: CD4+ regulatory T cells control TH17 responses in a Stat3-dependent manner. Science 326, 986-991 (2009).

- Clarke, T.B., Davis, K.M., Lysenko, E.S., Zhou, A.Y., Yu, Y., and Weiser, J.N.: Recognition of peptidoglycan from the microbiota by Nod1 enhances systemic innate immunity. Nat. Med. 16, 228-231 (2010).
- Coombes, J.L., Siddiqui, K.R.R., Arancibia-Cárcamo, C.V., Hall, J., Sun, C.-M., Belkaid, Y., and Powrie, F.: A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. J. Exp. Med. 204, 1757-1764 (2007).
- Crapser, J., Ritzel, R., Verma, R., Venna, V.R., Liu, F., Chauhan, A., Koellhoffer, E., Patel, A., Ricker, A., Maas, K., Graf, J., and McCullough, L.D.: Ischemic stroke induces gut permeability and enhances bacterial translocation leading to sepsis in aged mice. Aging (Albany NY) 8, 1049-1063 (2016).
- Dawson, D.A., Ruetzler, C.A., Carlos, T.M., Kochanek, P.M., and Hallenbeck, J.M.: Polymorphonuclear leukocytes and microcirculatory perfusion in acute stroke in the SHR. Keio J. Med. 45, 248-252; discussion 252-253 (1996).
- del Zoppo, G.J., Schmid-Schönbein, G.W., Mori, E., Copeland, B.R., and Chang, C.M.: Polymorphonuclear leukocytes occlude capillaries following middle cerebral artery occlusion and reperfusion in baboons. Stroke 22, 1276-1283 (1991).
- Denes, A., Ferenczi, S., Halász, J., Környei, Z., and Kovács, K.J.: Role of CX3CR1 (fractalkine receptor) in brain damage and inflammation induced by focal cerebral ischemia in mouse. J. Cereb. Blood Flow Metab. 28, 1707-1721 (2008).
- Duan, J., Chung, H., Troy, E., and Kasper, D.L.: Microbial colonization drives expansion of IL-1 receptor 1-expressing and IL-17-producing gamma/delta T cells. Cell Host Microbe 7, 140-150 (2010).
- Felger, J.C., Abe, T., Kaunzner, U.W., Gottfried-Blackmore, A., Gal-Toth, J., McEwen, B.S., Iadecola, C., and Bulloch, K.: Brain

dendritic cells in ischemic stroke: time course, activation state, and origin. Brain Behav. Immun. 24, 724-737 (2010).

- Ganal, S.C., Sanos, S.L., Kallfass, C., Oberle, K., Johner, C., Kirschning, C., Lienenklaus, S., Weiss, S., Staeheli, P., Aichele, P., and Diefenbach, A.: Priming of natural killer cells by nonmucosal mononuclear phagocytes requires instructive signals from commensal microbiota. Immunity 37, 171-186 (2012).
- Garcia-Bonilla, L., Moore, J.M., Racchumi, G., Zhou, P., Butler, J.M., Iadecola, C., and Anrather, J.: Inducible nitric oxide synthase in neutrophils and endothelium contributes to ischemic brain injury in mice. J. Immunol. 193, 2531-2537 (2014).
- Gelderblom, M., Leypoldt, F., Steinbach, K., Behrens, D., Choe, C.-U., Siler, D.A., Arumugam, T.V., Orthey, E., Gerloff, C., Tolosa, E., and Magnus, T.: Temporal and spatial dynamics of cerebral immune cell accumulation in stroke. Stroke. 40, 1849-1857 (2009).
- Gelderblom, M., Weymar, A., Bernreuther, C., Velden, J., Arunachalam, P., Steinbach, K., Orthey, E., Arumugam, T.V., Leypoldt, F., Simova, O., Thom, V., Friese, M.A., Prinz, I., Hölscher, C., Glatzel, M., Korn, T., Gerloff, C., Tolosa, E., and Magnus, T.: Neutralization of the IL-17 axis diminishes neutrophil invasion and protects from ischemic stroke. Blood 120, 3793-3802 (2012).
- Gelderblom, M., Gallizioli, M., Ludewig, P., Thom, V., Arunachalam, P., Rissiek, B., Bernreuther, C., Glatzel, M., Korn, T., Arumugam, T.V., Sedlacik, J., Gerloff, C., Tolosa, E., Planas, A.M., and Magnus, T.: IL-23 (Interleukin-23)-Producing Conventional Dendritic Cells Control the Detrimental IL-17 (Interleukin-17) Response in Stroke. Stroke 49, 155-164 (2017).
- Henninger, N., Kumar, R., and Fisher, M.: Acute ischemic stroke therapy. Expert Rev. Cardiovasc. Ther. 8, 1389-1398 (2010).
- Hill, D.A. and Artis, D.: Intestinal bacteria and the regulation of immune cell homeostasis. Annu. Rev. Immunol. 28, 623-667 (2010).
- Hou, L., Jie, Z., Desai, M., Liang, Y., Soong,

L., Wang, T., and Sun, J.: Early IL-17 production by intrahepatic T cells is important for adaptive immune responses in viral hepatitis. J. Immunol. 190, 621-629 (2013).

- Houlden, A., Goldrick, M., Brough, D., Vizi, E.S., Lénárt, N., Martinecz, B., Roberts, I.S., and Denes, A.: Brain injury induces specific changes in the caecal microbiota of mice via altered autonomic activity and mucoprotein production. Brain Behav. Immun. 57, 10-20 (2016).
- Huber, S., Gagliani, N., Esplugues, E., O'Connor, W., Huber, F.J., Chaudhry, A., Kamanaka, M., Kobayashi, Y., Booth, C.J., Rudensky, A.Y., Roncarolo, M.G., Battaglia, M., and Flavell, R.A.: Th17 cells express interleukin-10 receptor and are controlled by Foxp3⁻ and Foxp3⁺ regulatory CD4+ T cells in an interleukin-10-dependent manner. Immunity 34, 554-565 (2011).
- Hughes, P.M., Allegrini, P.R., Rudin, M., Perry, V.H., Mir, A.K., and Wiessner, C.: Monocyte chemoattractant protein-1 deficiency is protective in a murine stroke model. J. Cereb. Blood Flow Metab. 22, 308-317 (2002).
- Hurn, P.D., Subramanian, S., Parker, S.M., Afentoulis, M.E., Kaler, L.J., Vandenbark, A.A., and Offner, H.: T- and B-cell-deficient mice with experimental stroke have reduced lesion size and inflammation. J. Cereb. Blood Flow Metab. 27, 1798-1805 (2007).
- Iadecola, C. and Anrather, J.: The immunology of stroke: from mechanisms to translation. Nat. Med. 17, 796-808 (2011).
- Ivanov, I.I., McKenzie, B.S., Zhou, L., Tadokoro, C.E., Lepelley, A., Lafaille, J.J., Cua, D.J., and Littman, D.R.: The orphan nuclear receptor RORgammat directs the differentiation program of proinflammatory IL-17+ T helper cells. Cell 126, 1121-1133 (2006).
- Kleinschnitz, C., Schwab, N., Kraft, P., Hagedorn, I., Dreykluft, A., Schwarz, T., Austinat, M., Nieswandt, B., Wiendl, H., and Stoll, G.: Early detrimental T-cell effects in experimental cerebral ischemia are neither related to adaptive immunity nor thrombus formation. Blood 115, 3835-3842 (2010).

- Korn, T. and Petermann, F.: Development and function of interleukin 17-producing $\gamma\delta$ T cells. Ann. N Y Acad. Sci. 1247, 34-45 (2012).
- Kuhn, K.A., andStappenbeck, T.S.: Peripheral education of the immune system by the colonic microbiota. Semin. Immunol. 25, 364-369 (2013).
- Lee, Y.K., Menezes, J.S., Umesaki, Y., and Mazmanian, S.K.: Proinflammatory T-cell responses to gut microbiota promote experimental autoimmune encephalomyelitis. Proc. Natl. Acad. Sci. USA 108, Suppl 1, 4615-4622 (2011).
- Liesz, A., Suri-Payer, E., Veltkamp, C., Doerr, H., Sommer, C., Rivest, S., Giese, T., and Veltkamp, R.: Regulatory T cells are key cerebroprotective immunomodulators in acute experimental stroke. Nat. Med. 15, 192-199 (2009).
- Liesz, A., Zhou, W., Na, S.-Y., Hämmerling, G.J., Garbi, N., Karcher, S., Mracsko, E., Backs, J., Rivest, S., and Veltkamp, R.: Boosting regulatory T cells limits neuroinflammation in permanent cortical stroke. J. Neurosci. 33, 17350-17362 (2013).
- Ludewig, P., Sedlacik, J., Gelderblom, M., Bernreuther, C., Korkusuz, Y., Wagener, C., Gerloff, C., Fiehler, J., Magnus, T., and Horst, A.K.: Carcinoembryonic antigen-related cell adhesion molecule 1 inhibits MMP-9-mediated blood-brain-barrier breakdown in a mouse model for ischemic stroke. Circ. Res. 113, 1013-1022 (2013).
- Macrez, R., Ali, C., Toutirais, O., Le Mauff, B., Defer, G., Dirnagl, U., and Vivien, D.: Stroke and the immune system: from pathophysiology to new therapeutic strategies. Lancet Neurol. 10, 471-480 (2011).
- Mazmanian, S.K., Liu, C.H., Tzianabos, A.O., and Kasper, D.L.: An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system. Cell 122, 107-118 (2005).
- Meisel, C., Schwab, J.M., Prass, K., Meisel, A., and Dirnagl, U.: Central nervous system injury-induced immune deficiency syndrome. Nat. Rev. Neurosci. 6, 775-786 (2005).
- Moskowitz, M.A., Lo, E.H., and Iadecola, C.:

The science of stroke: mechanisms in search of treatments. Neuron. 67, 181-198 (2010).

- Nishio, J. and Honda, K.: Immunoregulation by the gut microbiota. Cell. Mol. Life Sci. 69, 3635-3650 (2012).
- Ochoa-Repáraz, J., Mielcarz, D.W., Ditrio, L.E., Burroughs, A.R., Foureau, D.M., Haque-Begum, S., and Kasper, L.H.: Role of gut commensal microflora in the development of experimental autoimmune encephalomyelitis. J. Immunol. 183, 6041-6050 (2009).
- Park, S.-G., Mathur, R., Long, M., Hosh, N., Hao, L., Hayden, M.S., and Ghosh, S.: T regulatory cells maintain intestinal homeostasis by suppressing $\gamma\delta$ T cells. Immunity 33, 791-803 (2010).
- Planas, A.M. and Chamorro, A.: Regulatory T cells protect the brain after stroke. Nat. Med. 15, 138-139 (2009).
- Prass, K., Meisel, C., Höflich, C., Braun, J., Halle, E., Wolf, T., Ruscher, K., Victorov, I.V., Priller, J., Dirnagl, U., Volk, H.-D., and Meisel, A.: Stroke-induced immunodeficiency promotes spontaneous bacterial infections and is mediated by sympathetic activation reversal by poststroke T helper cell type 1-like immunostimulation. J. Exp. Med. 198, 725-736 (2003).
- Roark, C.L., Simonian, P.L., Fontenot, A.P., Born, W.K., and O'Brien, R.L.: Gammadelta T cells: an important source of IL-17. Curr. Opin. Immunol. 20, 353-357 (2008).
- Rosell, A., Cuadrado, E., Ortega-Aznar, A., Hernández-Guillamon, M., Lo, E.H., and Montaner, J.: MMP-9-positive neutrophil infiltration is associated to blood-brain barrier breakdown and basal lamina type IV collagen degradation during hemorrhagic transformation after human ischemic stroke. Stroke 39, 1121-1126 (2008).
- Round, J.L. and Mazmanian, S.K.: Inducible Foxp3+ regulatory T-cell development by a commensal bacterium of the intestinal microbiota. Proc. Natl. Acad. Sci. USA 107, 12204–12209 (2010).
- Schilling, M., Besselmann, M., Leonhard, C., Mueller, M., Ringelstein, E.B., and Kiefer, R.: Microglial activation precedes and pre-

dominates over macrophage infiltration in transient focal cerebral ischemia: a study in green fluorescent protein transgenic bone marrow chimeric mice. Exp. Neurol. 183, 25-33 (2003).

- Schilling, M., Strecker, J.-K., Ringelstein, E.B., Schäbitz, W.-R., and Kiefer, R.: The role of CC chemokine receptor 2 on microglia activation and blood-borne cell recruitment after transient focal cerebral ischemia in mice. Brain Res. 1289, 79-84 (2009).
- Schulte-Herbrüggen, O., Quarcoo, D., Meisel, A., and Meisel, C.: Differential affection of intestinal immune cell populations after cerebral ischemia in mice. Neuroimmunomodulation 16, 213-218 (2009).
- Shichita, T., Sugiyama, Y., Ooboshi, H., Sugimori, H., Nakagawa, R., Takada, I., Iwaki, T., Okada, Y., Iida, M., Cua, D.J., Iwakura, Y., and Yoshimura, A.: Pivotal role of cerebral interleukin-17-producing gammadeltaT cells in the delayed phase of ischemic brain injury. Nat. Med. 15, 946-950 (2009).
- Singh, V., Roth, S., Llovera, G., Sadler, R., Garzetti, D., Stecher, B., Dichgans, M., and Liesz, A.: Microbiota Dysbiosis Controls the Neuroinflammatory Response after Stroke. J. Neurosci. 36, 7428-7440 (2016).
- Singh, V., Sadler, R., Heindl, S., Llovera, G., Roth, S., Benakis, C., and Liesz, A.: The gut microbiome primes a cerebroprotective immune response after stroke. J. Cereb. Blood Flow Metab. 38, 1293-1298 (2018).
- Sinkorová, Z., Capková, J., Niederlová, J., Stepánková, R., and Sinkora, J.: Commensal intestinal bacterial strains trigger ankylosing enthesopathy of the ankle in inbred B10.BR (H-2(k)) male mice. Hum. Immunol. 69, 845-850 (2008).
- Soriano, S.G., Amaravadi, L.S., Wang, Y.F., Zhou, H., Yu, G.X., Tonra, J.R., Fairchild-Huntress, V., Fang, Q., Dunmore, J.H., Huszar, D., and Pan, Y.: Mice deficient in fractalkine are less susceptible to cerebral ischemia-reperfusion injury. J. Neuroimmunol. 125, 59-65 (2002).
- Spychala, M.S., Venna, V.R., Jandzinski, M., Doran, S.J., Durgan, D.J., Ganesh, B.P.,

Ajami, N.J., Putluri, N., Graf, J., Bryan, R.M., and McCullough, L.D.: Age-related changes in the gut microbiota influence systemic inflammation and stroke outcome. Ann. Neurol., doi: 10.1002/ana.25250 (2018).

- Stanley, D., Mason, L.J., Mackin, K.E., Srikhanta, Y.N., Lyras, D., Prakash, M.D., Nurgali, K., Venegas, A., Hill, M.D., Moore, R.J., and Wong, C.H.Y.: Translocation and dissemination of commensal bacteria in post-stroke infection. Nat. Med. 22, 1277-1284 (2016).
- Stanley, D., Moore, R.J., and Wong, C.H.Y.: An insight into intestinal mucosal microbiota disruption after stroke. Sci Rep. 8, 568 (2018).
- Stubbe, T., Ebner, F., Richter, D., Engel, O., Randolf Engel, O., Klehmet, J., Royl, G., Meisel, A., Nitsch, R., Meisel, C., and Brandt, C.: Regulatory T cells accumulate and proliferate in the ischemic hemisphere for up to 30 days after MCAO. J. Cereb. Blood Flow Metab. 33, 37-47 (2013).
- Sun, C.-M., Hall, J.A., Blank, R.B., Bouladoux,

N., Oukka, M., Mora, J.R., and Belkaid, Y.: Small intestine lamina propria dendritic cells promote de novo generation of Foxp3 T reg cells via retinoic acid. J. Exp. Med. 204, 1775-1785 (2007).

- Suwanwela, N. and Koroshetz, W.J.: Acute ischemic stroke: overview of recent therapeutic developments. Annu. Rev. Med. 58, 89-106 (2007).
- Takeda, K., Clausen, B.E., Kaisho, T., Tsujimura, T., Terada, N., Förster, I., and Akira, S.: Enhanced Th1 activity and development of chronic enterocolitis in mice devoid of Stat3 in macrophages and neutrophils. Immunity 10, 39-49 (1999).
- Wu, H.-J., Ivanov, I.I., Darce, J., Hattori, K., Shima, T., Umesaki, Y., Littman, D.R., Benoist, C., and Mathis, D.: Gut-residing segmented filamentous bacteria drive autoimmune arthritis via T helper 17 cells. Immunity 32, 815-827 (2010).
- Yilmaz, G., Arumugam, T.V., Stokes, K.Y., and Granger, D.N.: Role of T lymphocytes and interferon-gamma in ischemic stroke. Circulation 113, 2105-2112 (2006).

THE MICROBIOME, IMMUNE DEVELOPMENT, CHRONIC INFLAMMATION, AND CANCER

TAYLOR N. TIBBS¹*, LACEY R. LOPEZ¹*, and JANELLE C. ARTHUR^{1,2,3}

¹Department of Microbiology and Immunology, ²Center for Gastrointestinal Biology and Disease, and ³Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

INTRODUCTION

Our microbiota is an integral part of our mammalian selves. Indeed, all multi-cellular eukaryotic hosts across the tree of life have an essential and characteristic microbiota that influences host development and resistance to disease. Within complex organisms such as vertebrates, we harbour numerous microbial communities whose composition and function are relevant to their habitat at different body sites, such as the intestines ("gut"), skin, and oral cavity. Our gut microbiota is perhaps the best studied, most abundant, and arguably, the most influential microbiota that impacts host phenotypes. In the recent decades, the development of several scientific tools has exponentially increased our understanding of the microbiota and interactions with its human host. These include model organisms, most notably "gnotobiotic" laboratory mice that are born and raised germ-free (GF) and then colonized with individual strains or groups of microbes. Through the use of GF and gnotobiotic mice, we have been able to demonstrate causality of specific microbes and microbial groups with distinct processes of immune development and non-infectious diseases like chronic inflammation and cancer, among others. To validate the

physiologic relevance of observations made in model organisms with human disease, we can now survey the human microbiota at an unprecedented depth using culture-independent molecular methods (i.e. targeted 16S "microbiome" sequencing, metagenomics, metatranscriptomics, and metabolomics) coupled with sophisticated bioinformatics pipelines. An important finding from population studies of the microbiome has revealed that the compositional fluctuations in an individual's microbiome over time are less substantial than inter-individual differences at a particular stage in development. However, the developmental changes that occur during early life and over an individual's lifespan certainly shape the composition and function of the microbiota. Indeed, the composition and functional capabilities of the microbiota shape host development. The focus of the 2018 Old Herborn University Seminar Series 32 is "Ageing and the Microbiome". Accordingly, in this chapter we discuss the current state of knowledge regarding the influence of our mammalian microbiota on the immune system, chronic inflammation, and a prominent disease of the ageing – cancer.

^{*}Both first authors contributed equally to this chapter

THE IMMUNE SYSTEM AND MICROBIOME IN AGEING

The immune system is our major host defence system that is educated early in life to distinguish harmful stimuli, including bacteria. It is broadly subdivided into two branches, the innate and adaptive immune systems. The innate immune response is the first line of defence towards invading pathogens. For example, this branch of the immune system is involved when host cell pattern recognition receptors are triggered by conserved pathogen specific molecules which promote an immediate and broad-spectrum protective response to pathogens. Conversely, the adaptive immune response relies on antigen presenting cells (APCs) to present bacterial products to effector T and B lymphocytes and create a pathogen-specific immunological memory and response. Through precise coordination, these systems provide host defence against foreign invaders. However, to efficiently mediate its response, the immune system must accurately distinguish between normal host-associated organisms and those that are potentially deleterious (Chaplin, 2010). Over the course of our lifetime, our immune system encounters a diverse range of stimuli in a variety of contexts, which challenges the ability of the immune system to differentiate between self and nonself (Ponnappan and Ponnappan, 2011). Alterations in this response can result in the development of a variety of diseases that include inflammatory bowel disease (IBD) and cancer (Narendra et al., 2013; Gálvez, 2014). As these immune responses are shaped over a lifetime, it indicates that age impacts the recognition of stimuli, which could transfer towards inappropriately reacting to residential host microbes. Inappropriate reactions to normal commensal bacteria may underlie a variety of pathogenic processes.

Microbial influences on immune development

Establishment of the microbiota begins from birth and continues until postweaning (Bergström et al., 2014). During establishment, the microbiota is highly diverse and prone to fluctuations based on environmental and dietary changes (Penders et al., 2006). After 2-3 years of age, this complex community stabilizes with the majority of bacterial community members remaining unchanged throughout the lifespan of an individual (Rajilić-Stojanović et al., 2013). After stabilization, the core human microbiota mainly comprises the following phyla: *Bacteroidetes* and *Fir*micutes, with a smaller abundance of Actinobacteria, Proteobacteria, and Verrucomicrobia (Arumugam et al., 2011). The ageing process strongly impacts the composition of the microbiota as individuals with increased frailty, an assessment of biological age based on current health status and life expectancy, lose bacterial diversity and form a Bacteroidetes dominant population (*Claesson* et al., 2011). While not directly correlated to chronological ageing, this loss of diversity most often begins between 75-80 years of age and is a form of dysbiosis that potentiates disease development (Biagi et al., 2010). These results are a generality from a diverse population, and only consider numerical age (lifespan) rather than the individual's ageing-related health status (health span). Despite this correlation, it is important to remember that the microbiota play a largely protective role in disease development by initiating and educating the host immune system (Pickard et al., 2017).

Early observations in GF mice demonstrated that the host microbiota is essential for the maturation of the immune system (*Clavel* et al., 2017; *Pickard* et al., 2017). In the absence of

a microbiota, GF mice have several immunological defects, including reduced lymphoid cell numbers and function (Fiebiger et al., 2016). For example, GF mice have less T helper type 1 (Th1) cells compared to their conventionalized counterparts (Wu and Wu, 2012). Th1 cells promote cell-mediated immune responses and phagocyte-dependent inflammation to target intracellular pathogens (Damsker et al., 2010). Th1 responses in GF mice can be restored through host colonization with a variety of microbes, including Listeria monocytogenes which promotes Th1 development through macrophage production of the T-cell-stimulating factor, interleukin 12 (IL-12) (Hsieh et al., 1993). In the gut, Th1 responses are driven by intracellular bacteria like L. monocytogenes (Atarashi et al., 2015). Additionally, GF mice have a reduced number of T helper type 17 (Th17) cells. Th17 cells are like Th1 cells in that they are pro-inflammatory; however, they drive the production of IL-17, a cytokine which mediates defence against extracellular pathogens and auto-immune disease (Damsker et al., 2010; Wu and Wu, 2012). Colonization of GF mice with intestinally adherent pathogens, such as Clostridia-related segmented filamentous bacteria (SFB), induces the development of Th17 cells in the small intestine by driving the release of serum amyloid A from intestinal epithelial cells (IECs). The release of serum amyloid A results in the production of innate lymphoid cell group 3 activating cytokines which upregulate the Th17 response (*Ivanov* et al., 2009). Fine-tuning of Th1 and Th17 responses is essential for immune tolerance towards the host microbiota, as seen in the case of IBD where aberrant populations of Th1 and Th17 cells lead to en-(Gálvez, hanced pathology 2014). Therefore, underdevelopment of these responses may underlie the pathogenesis of other diseases associated with chronic inflammation, such as cancer (*Bailey* et al., 2014; *Vinay* et al., 2015).

The absence of a microbiota impacts most, if not all, aspects of the immune system (*Round* and *Mazmanian*, 2009). However, it is not fully understood where the window of opportunity lies for correcting many of these immune deficiencies. One study examining colonic invariant natural killer T (iNKT) cell populations revealed that this opportunity for modulation likely occurs in early life (An et al., 2014). At birth, GF mice have an enriched population of colonic lamina propria iNKT cells compared to specific-pathogen-free (SPF) mice (*Olszak* et al., 2012). iNKT cells are pro-inflammatory and mediate cell tolerance to commensal microbes (*Chandra* and *Kronenberg*, 2015). Colonization of adult (>5 weeks of age) GF mice with a complex microbiota does not influence the number and activity of iNKT cells (An et al., 2014). However, if the colonization occurs when GF mice are neonates, the number of iNKT cells is reduced and their later activation is well-controlled (An et al., 2014). This early education of the colonic iNKT cell population is important for limiting morbidity associated with IBD (An et al., 2014). This supports the idea that exposure to specific microbes and microbial products is needed within a certain developmental time for the host to appropriately educate their target immune population and prevent disease.

The gastrointestinal tract houses one of the largest and most diverse collection of microbes in the human body (*Thursby* and *Juge*, 2017). In the intestine, a single layer of epithelial cells separates the underlying mucosal immune system from the microbiota. Due to this proximity, much work has focused upon the protective role of the microbiota on inflammatory immune

responses that can lead to intestinal inflammation or colitis. For example, GF mice colonized with a beneficial nontoxin-producing strain of Bacteroides *fragilis* exhibit a marked expansion of regulatory T-cells (Tregs) in the intestinal mucosa (Round and Mazmanian, 2010). This expansion of Tregs limits inappropriate inflammation and is important for preventing autoimmune disease through immunotolerance towards self-antigens (Round and Mazmanian, 2010; Rothstein and Camirand, 2015). This protection is mediated by Bac*teroides spp.* through the production of polysaccharide-A (PSA), a commensal microbial surface molecule presented by APCs to modulate Treg specific immune responses. Indeed, colonization with PSA + B. fragilis protects mice from chemically TNBS-induced and Helicobacter hepaticus-induced colitis (Round and Mazmanian, 2010). Furthermore, colonization of conventional mice with Clostridium butyricum protects against experimentally-induced colitis by promoting the expansion of immunosuppressive interleukin-10 (IL-10) secreting macrophages and mature Tregs (Hayashi et al., 2013). B. fragilis and C. butyricum belong to the most highly represented bacterial phyla within the murine and human microbiota (Bacteroidetes and Firmicutes, respectively) and mechanistically represent some of the many ways the commensal microbiota can modulate host immune responses to prevent the onset of disease (Belkaid and Harrison, 2017; Clavel et al., 2017; Pickard et al., 2017).

The immune system and ageing

The presence of a microbiota in early life is essential for immune system maturation. However, education of the immune response is a lifelong process. Alterations to the innate and adaptive immune systems which occur with increased frailty are linked to a complex biological process known as immunosenescence (Fülöp et al., 2016). Specific changes associated with immunosenescence can best be understood through functional differences within the unique cell types of the innate and adaptive immune systems. For cells of the innate immune system, there are reported functional differences for every major cell type (Castelo-Branco and Soveral, 2014). However, the most distinct differences are within neutrophil and macrophage populations. Neutrophils isolated from the blood of individuals, aged 62-83, displayed reduced phagocytic capabilities and decreased production of reactive oxygen species (ROS) when infected with Staphylococcus aureus (Wenisch et al., 2000). These neutrophils, versus cells isolated from younger adult patients, also had impaired bactericidal activity (Wenisch et al., 2000). Neutrophils are the first line of defence towards invading patho-Therefore, immunosenescence gens. related changes to this cell type suggest an age-related decline in pathogen tolerance (Chaplin, 2010). Similarly, primary macrophages isolated from aged mice (18-24 months old) have impaired phagocytosis and reduced ROS production in response to infection when compared to macrophages isolated from young mice (2-3 months old) (*Davila* et al., 1990; Swift et al., 2001). Additionally, macrophages from aged mice display altered antigen presentation and reduced production of pro-inflammatory cytokines (Herrero et al., 2001; Renshaw et al., 2002). Alterations in macrophage antigen presentation and cytokine release may lead to altered immune signalling between the innate and adaptive immune systems, resulting in a weakened immune response (Chaplin, 2010). Overall, age-related changes to the innate immune system strongly reduce the host's initial response to pathogens.

While changes in the innate immune system have been noted with ageing, alterations to the adaptive immune system may be more pronounced (*Linton* and Dorshkind, 2004; Castelo-Branco and Soveral, 2014). The adaptive immune system is used for long-term protection from environmental insult and invading pathogens. Therefore, longterm education of this subsystem could have additive effects on immunosenescence. B-cells are one of the major cell types of the adaptive immune system. The population of B-cells is divided into plasma cells or memory B-cells. Plasma cells produce pathogen specific antibodies, while memory B-cells provide long-term recognition of antigens (Eibel et al., 2014). Peripheral blood isolated from elderly individuals (aged 86-94 years) showed a reduction in Bcell population diversity, likely due to a decrease in memory B-cells (Gibson et al., 2009). This decline in B-cell diversity was linked to increased frailty and may be used as a predictor for general health status (*Gibson* et al., 2009). A reduction in memory B-cells may cause an inappropriate immune response towards the microbiota, as B-cells are important for establishing the distinction between pathogenic and commensal bacteria (*Eibel* et al., 2014).

T-cells are the second major cell type of the adaptive immune system. Under normal immunological conditions, T-cells differentiate into either T Helper (Th) cells or natural killer T (NKT) cells. This differentiation is based on antigen presentation by APCs and is necessary to form a pathogenspecific immune response (*Chandra* and *Kronenberg*, 2015). Populations of T-cells isolated from the peripheral blood of elderly humans (aged 72-89 years) showed a reduction in the proportion of NKT-cells versus cells isolated from young patients (aged 25-

30 years) (DelaRosa et al., 2002). This finding was matched by a study which showed a reduction in the proportion of NKT-cells isolated from the peripheral blood of adults aged 61 years and over (Jing et al., 2007). Additionally, NKTcells isolated from the liver of aged mice (aged ≥ 20 months) demonstrated a decline in cytotoxic function, and reduced cytokine release versus T-cells isolated from young mice (aged 2 months) (Mocchegiani et al., 2004). Under non-immunosenescence conditions, NKT-cells respond to antigenic activation by robust proliferation and directed cytotoxicity (Chandra and Kronenberg, 2015). Therefore, the reduced proportion of NKT-cells and cytotoxic ability in elderly organisms may exacerbate disease development by weakening the host's response to pathogens.

The host microbiota initiates immune system maturation in early life. However, to keep up with a lifelong antigenic load, the immune response must be fine-tuned and properly educated across the lifespan. Changes to the immune system that occur with age are noted within immunosenescence whereby functional differences develop within innate and adaptive immune cell populations. Despite these observations, it remains unclear how these immunological changes impact cellular crosstalk and overall immunocompetence. On top of this, how the microbiota impacts the immune system during immunosenescence remains to be elucidated. It is likely that changes to the immune system result in an inappropriate response towards commensal microbes, as indicated by diseases like IBD (Sun et al., 2015). Therefore, these inappropriate reactions to the native microbiota may contribute to the development of chronic inflammation and the onset of age-related diseases, such as cancer (*Tilg* et al., 2018).

Cancer is considered a disease of old age. As life expectancy increases, the estimated rate of cancer is predicted to increase by 45% from 2011 to 2030 in the United States (Smith et al., 2009). It is also estimated that by 2030, individuals 65 years and older will contribute to 70% of all cancers in the United States (White et al., 2014). The risk of developing cancer increases dramatically with age as the duration of time in which an individual is exposed to carcinogens increases (Harding et al., 2012), the proliferative capacity of ageing cells decreases (Pompei et al., 2001), and immunological competence decreases (Bonafè et al., 2002). Cancer typically results from a series of genetic mutations or epigenetic modifications that develop sequentially overtime (Loeb et al., 2003). The colon, which harbours the largest and most diverse microbiota of all organs, has the highest incidence rate of all reported cancers in the 85+ population (*Thakkar* et al., 2014). Over the past decade, we have become increasingly aware of the roles that the microbiota play in the development of cancer and modulation of cancer therapies. We have also elucidated several mechanisms underlying the microbial influences on cancer. However, these roles are diverse and seem to influence many aspects of immune and cancer development (Fulbright et al., 2017). Microbes can contribute to the onset and progression of cancer through direct means, such as by producing genotoxins, and indirect means through the modulation of immune responses to tumours and immunotherapy (Rhee et al., 2009; Arthur et al., 2012; Kostic et al., 2013; Boleij et al., 2015; Gopala-krishnan et al., 2018). Additionally, members of native several the microbiota can alter chemotherapeutic

drugs, resulting in unpleasant side effects for the host or even rendering them clinically inert (*Al-Dasooqi* et al., 2011; *Geller* et al., 2017). It is therefore important to divulge the significance of microbial interactions on agerelated diseases, such as cancer, in order to fully understand disease progression and design suitable therapies.

The ability to manipulate the microbiota using GF and gnotobiotic mice has demonstrated the importance of the microbiota on immune system development. The microbiota assist in training the immune system to recognize harmful and non-harmful stimuli, both from the environment and within the self. Here we will discuss known mechanisms by which the microbiota can influence anti-cancer chemotherapeutic activities and anti-tumour immunomodulatory responses. For example, *Clostridia spp.* are capable of suppressing the body's anti-tumour immune responses. Host-derived primary bile acids are converted into secondary bile acids by the gut microbiota, primarily members of the genus Clostridia, and circulated systemically throughout the body via hepatic circulation (Ridlon et al., 2006). Previous work illustrated that secondary bile acids can increase the risk of obesity-associated hepatocellular carcinoma in susceptible mice (Yoshimoto et al., 2013). Recent data suggests that antibiotic elimination of the gut microbiota in mice decreases both primary and metastatic tumours within the liver by facilitating the build-up of primary bile acids, which trigger liver-specific NKT-cell recruitment to target cancer cells (Ma et al., 2018). The profound effects bacteria illicit on cytotoxic immune cells provide key insights on how the native microbiota influence host anti-tumour responses.

Another example, Fusobacterium nucleatum, a Gram-negative oral commensal overrepresented in colorectal carcinoma, can promote tumourigenesis through the modulation of the innate immune system (*Castellarin* et al., 2012; Kostic et al., 2013). A known target is the natural killer (NK) cell, which kills compromised host cells, such as infected or cancerous cells. F. *nucleatum* inhibits the cytotoxicity of NK-cells via the Fusobacterium protein Fap2 which binds the NK-cell inhibitor receptor TIGIT (T-cell immunoglobulin and ITIM or immunoreceptor tyrosine-based inhibition motif domain) (Gur et al., 2015). In addition to targeting the immune system, it should be mentioned that F. nucleatum exerts pro-carcinogenic activities directly on epithelial cells through β-catenin signalling, altering cell fate (Rubinstein et al., 2013). F. nucleatum can also alter the efficacy of chemotherapeutic drugs by inhibiting host cell apoptotic pathways (Yu et al., 2017). F. nucleatum is a prime example of one species of the microbiota that exhibits a variety of different effects on the host to mediate tumourigenesis and hinder cancer therapy. In the next few sections, we will discuss a variety of known bacterial mechanisms that act upon cancer development and treatment.

Cancer immunotherapy and the microbiota

Several independent groups have recently demonstrated that some members of the microbiota play critical roles in determining patient responsiveness to cancer immunotherapy. The exact mechanisms by which individual species of bacteria exhibit these effects are not fully understood. However, current data suggest that bacterial modulation of the immune system may be one critical mode of altering host response to cancer therapy. Recent data regard-

ing anti-PD1 therapy supports this notion. Anti-PD1 treatment is a type of immune checkpoint inhibitor that enhances anti-tumour immune responses by maintaining T-cell activation via blocking the immune inhibitory receptors programmed death ligand-1 and 2 (PDL-1 and PDL-2) (Shields et al., 2017). Anti-PD1 therapy is often prescribed to patients with lung cancer and advanced melanoma. However, the efficacy of anti-PD1 immunotherapy ranges from only 19 to 43% for both cancer types (Jiang et al., 2015; Larkin et al., 2015). Several members of the microbiota are enriched in PD-1 responders, including Bifidobacterium longum, Collinsella aerofaciens, and Enterococcus faecium (Matson et al., 2018). Faecalibacterium, an abundant Gram-positive genus of commensals in the human gut (Shabbir et al., 2016), was also enriched in PD1-responders (Gopalakrishnan et al., 2018). Tumourbearing mice that were given faecal microbiota transplants (FMTs) from PD1-responders exhibited decreased tumour burden and tumour size when receiving anti-PD1 therapy. Faecalibacterium promoted cytotoxic (CD8+) T-cell recruitment to tumours, which may be an important mechanism underlying the ability of this bacterial group to enhance anti-PD1 responses and reduce tumour burden (Gopalakrishnan et al., 2018). Similarly, another group demonstrated that FMTs from PD-1 responders enhanced PD-1 treatment in recipient mice, which was further augmented with oral supplementation of the commensal Akkermansia muciniphila (Routy et al., 2018). Antibiotic treatment reduced the efficacy of PD-1 immunotherapy in mice, consistent with clinical reports of reduced PD-1 efficacy in patients simultaneously taking antibiotics (Routy et al., 2018). These studies demonstrate that multiple species of bacteria have the capability of altering immunotherapeutic responses in patients. Moving forward, it may be critical to consider the contributions of these microbial communities when developing anti-tumour immunotherapies.

Chemotherapy and the microbiota

Multiple members of the microbiota can differentially influence cancer chemotherapy, with some enhancing and some inhibiting the clinical effects of chemotherapeutic drugs. An important early observation was that genotoxic platinum chemotherapies, including oxaliplatin and cisplatin, were ineffective in tumour-bearing GF mice (*lida* et al., 2013). Thus, the presence of a complex microbiota is essential for these chemotherapies (*Iida* et al., 2013). Platinum chemotherapies utilize ROS to induce cytotoxicity. The DNA damage induced by cisplatin is augmented via the production of mitochondrial ROS within cancer cells themselves (Marullo et al., 2013). Data also indicates that the ROS production from tumour-associated inflammatory cells are critical for the efficacy of these platinum chemotherapeutic drugs (*Iida* et al., 2013). It may be that in the absence of a native microbiota, inflammatory cells are not effectively primed to produce ROS during development, leading to shortcomings in ROS production later in life. This data highlights the influential capacity of the microbiota on the host and how in their absence the immune system may have substantial deficits in anti-tumour responses. Conversely, the microbiota can have negative effects on chemotherapeutic efficacy. Deep sequencing for microbes within pancreatic tumour biopsies revealed that 57.5% of pancreatic tumour tissues tested (65 of 113) samples) were positive for bacterial reads, with Gammaproteobacteria being the most abundant (51.7% of reads)

(Geller et al., 2017). Interestingly, 98.4% of Gammaproteobacteria contain genes that encode a specific isoform of the enzyme cytidine deaminase (CDD_L), which has the ability to breakdown gemcitabine and confer chemotherapeutic resistance in tumour tissues (Geller et al., 2017). Accordingly, bacterial migration from the gastrointestinal tract into the pancreatic ducts and tumour tissue may be a significant source of drug failure in clinical pancreatic cancer cases. Gut bacteria are responsible for also re-activating chemotherapeutic drugs in the distal intestine. Irinotecan, a chemotherapeutic drug used to treat colorectal cancer (CRC), is inactivated by the liver, but reactivated into the activate drug by *Clostridia spp.* through β-glucuronidases in the gut (*Stringer* et al., 2009). This re-activation contributes to the typical unpleasant gastrointestinal side effects of irinotecan therapy, including mucositis and diarrhoea (Stringer et al., 2009; Al-Dasooqi et al., 2011). This evidence illustrates the profound impact the native microbiota can have on the response to cancer therapies. Therefore, future treatment plans should account for the influence of these patientspecific microbial factors to ensure successful outcomes.

Direct effects of the microbiota on tumourigenesis

Specific members of the microbiota have the capacity to directly contribute to tumourigenesis (*Schwabe* and *Jobin*, 2013; *Fulbright* et al., 2017). Commensal *Enterobacteriaceae*, including several strains of *Escherichia coli*, are capable of inducing DNA damage in mammalian cells by producing a genotoxin termed colibactin (*Cuevas-Ramos* et al., 2010; *Arthur* et al., 2012). The bacterial polyketide synthase (*pks*) pathogenicity island encoding colibactin is up-regulated in CRC models and the presence of these gene products promotes tumourigenesis by inducing DNA double-stranded breaks (Nougayrede et al., 2006; Arthur et al., 2014; Tomkovich et al., 2017). Colibactin can also induce premature cellular senescence in cells that initially survive the DNA damage (Secher et al., 2013; Cougnoux et al., 2014) Furthermore, the *pks* pathogenicity island is overrepresented in the microbiota of CRC and IBD patients (Arthur et al., 2012; Buc et al., 2013; Prorok-Hamon et al., 2014). This suggests that bacterial genotoxins likely contribute substantially in the development of human cancer and chronic inflammatory diseases.

Bacteria also have the capacity to induce a pro-tumourigenic environment through chronic inflammation. Enterotoxigenic *B. fragilis*, a member of the most abundantly represented genus in the gut, produces its own flavour of toxin called B. fragilis-derived toxin (BFT) (*Wu* et al., 2002). BFT is a zincdependent metalloprotease that can induce colitis and promote tumourigenesis through the generation of ROS and subsequent initiation of DNA damage in epithelial cells (Goodwin et al., 2011). Enterotoxigenic B. fragilis robustly activates Th17 immune responses, which involves the inflammatory cytokine interleukin-17 (IL-17), and may lower host anti-tumour immune responses, encouraging unhindered tumour growth (*Wu* et al., 2009; Geis et al., 2015). Enterotoxigenic B. *fragilis* is overrepresented in patients with CRC when compared to healthy individuals (*Boleij* et al., 2015) and exacerbates tumourigenesis in susceptible mice (*Wu* et al. 2009). Interestingly, the tumourigenic effects of *pks+ E. coli* and enterotoxigenic B. fragilis act synergistically in vivo to quicken tumour onset and increase mortality in susceptible mice beyond the capability

that either species has individually (*Dejea* et al., 2018). Given that the native microbial community is quite complex, the cumulative effects of microbial products on the host may significantly contribute to the onset of cancer.

One of the earliest identified microbial suspects of inflammation-mediated cancer development is Helicobacter pylori. While widely considered a pathogen, H. pylori is estimated to be present in the gastrointestinal tract of over half of the human population worldwide and a major risk factor for gastric adenocarcinoma (Arthur and Jobin, 2011; Hooi et al., 2017). It is estimated that *H. pylori* infection increases the attributable risk of gastric by 73% (Herrera cancer and Parsonnet, 2009). Chronic H. pylori infection results in inflammation and tissue damage by the bacterial virulence factor CagA (cytotoxin-associated gene A), which initiates the development of the hallmark precursory lesions of gastric cancer, including intestinal metaplasia and dysplasia (*Díaz* et al., 2018). It remains unclear why H. pylori infection only progresses to malignancy in a subset of infected individuals; however, it is postulated that host immune responses and the genetics of both host and bacteria contribute to neoplastic development (Polk and *Peek*, 2010). The evidence presented here illustrates the diverse microbial mechanisms that contribute to tumourigenesis, whether that be by directly targeting the DNA for damage through a toxin or by providing an augmented environment for unrestricted cellular proliferation.

In summary, the mechanisms by which the native microbiota influences cancer development and therapy are numerous and diverse. As more data surfaces, it will be imperative to synthesize and apply knowledge on microbial contributions towards cancer development and treatment. By doing so, we can more effectively assess cancer risk and ultimately design more potent therapies.

LITERATURE

- Al-Dasooqi, N., Bowen, J.M., Gibson, R.J., Logan, R.M., Stringer, A.M.,and Keefe, D.M.: Irinotecan-induced alterations in intestinal cell kinetics and extracellular matrix component expression in the dark agouti rat. Int. J. Exp. Pathol. 92, 357-365 (2011).
- An, D., Oh, S.F., Olszak, T., Neves, J.F., Avci, F., Erturk-Hasdemir, D., Lu, X., Zeissig, S., Blumberg, R.S., and Kasper, D.L.: Sphingolipids from a symbiotic microbe regulate homeostasis of host intestinal natural killer T cells. Cell 156, 123-133 (2014).
- Arthur, J.C. and Jobin, C.: The struggle within: microbial influences on colorectal cancer. Inflamm. Bowel Dis. 17, 396-409 (2011).
- Arthur, J.C., Perez-chanona, E., Mühlbauer, M., Tomkovich, S., Uronis, J.M., Fan, T., Campbell, B.J., Abujamel, T., Dogan, B., Rogers, A.B., Rhodes, J.M., Stintzi, A., Simpson, K.W., Hansen, J.J., Keku, T.O., Fodor, A.A., and Jobin, C.: Intestinal inflammation targets cancer-inducing activity of the microbiota. Science 338, 120-123 (2012).
- Arthur, J.C., Gharaibeh, R.Z., Mühlbauer, M., Perez-Chanona, E., Uronis, J.M., McCafferty, J., Fodor, A.A., and Jobin, C.: Microbial genomic analysis reveals the essential role of inflammation in bacteriainduced colorectal cancer. Nat. Commun. 5, 2724 (2014).
- Arumugam, M., Raes, J., Pelletier, E., Le Paslier, D., Yamada, T., Mende, D.R., Fernandes, G.R., Tap, J., Bruls, T., Batto, J.M., Bertalan, M., Borruel, N., Casellas, F., Fernandez, L., Gautier, L., Hansen, T., Hattori, M., Hayashi, T., Kleerebezem, M., Kurokawa, K., Leclerc, M., Levenez, F., Manichanh, C., Nielsen, H.B., Nielsen, T., Pons, N., Poulain, J., Qin, J., Sicheritz-Ponten, T., Tims, S., Torrents, D., Ugarte, E.,

Zoetendal, E.G., Wang, J., Guarner, F., Pedersen, O., de Vos, W.M., Brunak, S., Doré, J.; MetaHIT Consortium, Antolín, M., Artiguenave, F., Blottiere, H.M., Almeida, M., Brechot, C., Cara, C., Chervaux, C., Cultrone, A., Delorme, C., Denariaz, G., Dervyn, R., Foerstner, K.U., Friss, C., van de Guchte, M., Guedon, E., Haimet, F., Huber, W., van Hylckama-Vlieg, J., Jamet, A., Juste, C., Kaci, G., Knol, J., Lakhdari, O., Layec, S., Le Roux, K., Maguin, E., Mérieux, A., Melo Minardi, R., M'rini, C., Muller, J., Oozeer, R., Parkhill, J., Renault, P., Rescigno, M., Sanchez, N., Sunagawa, S., Torrejon, A., Turner, K., Vandemeulebrouck, G., Varela, E., Winogradsky, Y., Zeller, G., Weissenbach, J., Ehrlich, S.D., and Bork, P.: Enterotypes of the human gut microbiome. Nature 473, 174-180 (2011).

- Atarashi, K., Tanoue, T., Ando, M., Kamada, N., Nagano, Y., Narushima, S., Suda, W., Imaoka, A., Setoyama, H., Nagamori, T., Ishikawa, E., Shima, T., Hara, T., Kado, S., Jinnohara, T., Ohno, H., Kondo, T., Toyooka, K., Watanabe, E., Yokoyama, S., Tokoro, S., Mori, H., Noguchi, Y., Morita, H., Ivanov, I.I., Sugiyama, T., Nuñez, G., Camp, J.G., Hattori, M., Umesaki, Y., and Honda, K.: Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial Cells. Cell 163, 367-380 (2015).
- Bailey, S.R., Nelson, M.H., Himes, R.A., Li, Z., Mehrotra, S., and Paulos, C.M.: Th17 cells in cancer: the ultimate identity crisis. Front. Immunol. 5, 1-13 (2014).
- Belkaid, Y. and Harrison, O.J.: Homeostatic immunity and the microbiota. Immunity 46, 562-576 (2017).
- Bergström, A., Skov, T.H., Bahl, M.I., Roager, H.M., Christensen, L.B., Ejlerskov, K.T., Mølgaard, C., Michaelsen, K.F., and Licht, T.R.: Establishment of intestinal microbiota during early life: a longitudinal, explorative
study of a large cohort of Danish infants. Appl. Environ. Microbiol. 80, 2889-2900 (2014).

- Biagi, E., Nylund, L., Candela, M., Ostan, R., Bucci, L., Pini, E., Nikkïla, J., Monti, D., Satokari, R., Franceschi, C., Brigidi, P., and De Vos, W.: Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians. PLoS One 5, e10667 (2010).
- Boleij, A., Hechenbleikner, E.M., Goodwin, A.C., Badani, R., Stein, E.M., Lazarev, M.G., Ellis, B., Carroll, K.C., Albesiano, E., Wick, E.C., Platz, E.A., Pardoll, D.M., and Sears, C.L.: The Bacteroides fragilis toxin gene is prevalent in the colon mucosa of colorectal cancer patients. Clin. Infect. Dis. 60, 208-215 (2015).
- Bonafè, M., Barbi, C., Storci, G., Salvioli, S., Capri, M., Olivieri, F., Valensin, S., Monti, D., Gonos, E.S., De Benedictis, G., and Franceschi, C.: What studies on human longevity tell us about the risk for cancer in the oldest old: data and hypotheses on the genetics and immunology of centenarians. Exp. Gerontol. 37, 1263-1271 (2002).
- Buc, E., Dubois, D., Sauvanet, P., Raisch, J., Delmas, J., Darfeuille-Michaud, A., Pezet, D., and Bonnet, R.: High prevalence of mucosa-associated E. coli producing cyclomodulin and genotoxin in colon cancer. PLoS One 8, e56964 (2013).
- Castellarin, M., Warren, R.L., Freeman, D.J., Dreolini, L., Krzywinski, M., Strauss, J., Barnes, R., Watson, P., Allen-Vercoe, E., Moore, R.A., and Holt, R.A.: Fusobacterium nucleatum infection is prevalent in human colorectal carcinoma. Genome Res. 22, 299-306 (2012).
- Castelo-Branco, C. and Soveral, I.: The immune system and aging: a review. Gynecol. Endocrinol. 30, 16-22 (2014).
- Chandra, S. and Kronenberg, M.: Activation and Function of iNKT and MAIT Cells. Adv. Immunol. 127, 145-201 (2015).
- Chaplin, D.D.: Overview of the Immune Response David. J. Allergy Clin. Immunol. 125, Suppl. 2, S3-23 (2010).
- Claesson, M.J., Cusack, S., O'Sullivan, O.,

Greene-Diniz, R., de Weerd, H., Flannery, E., Marchesi, J.R., Falush, D., Dinan, T., Fitzgerald, G., Stanton, C., van Sinderen, D., O'Connor, M., Harnedy, N., O'Connor, K., Henry, C., O'Mahony, D., Fitzgerald, A.P., Shanahan, F., Twomey, C., Hill, C., Ross, R.P., and O'Toole, P.W.: Composition, variability, and temporal stability of the intestinal microbiota of the elderly. Proc. Natl. Acad. Sci. USA 108, 4586-4591 (2011).

- Clavel, T., Gomes-Neto, J.C., Lagkouvardos, I., and Ramer-Tait, A.E.: Deciphering interactions between the gut microbiota and the immune system via microbial cultivation and minimal microbiomes. Immunol. Rev. 279, 8-22 (2017).
- Cougnoux, A., Dalmasso, G., Martinez, R., Buc, E., Delmas, J., Gibold, L., Sauvanet, P., Darcha, C., Déchelotte, P., Bonnet, M., Pezet, D., Wodrich, H., Darfeuille-Michaud, A., and Bonnet, R.: Bacterial genotoxin colibactin promotes colon tumour growth by inducing a senescenceassociated secretory phenotype. Gut 63, 1932-1942 (2014).
- Cuevas-Ramos, G., Petit, C.R., Marcq, I., Boury, M., Oswald, E., and Nougayrède, J.-P.: Escherichia coli induces DNA damage in vivo and triggers genomic instability in mammalian cells. Proc. Natl. Acad. Sci. USA 107, 11537-11542 (2010).
- Damsker, J.M., Hansen, A.M., and Caspi, R.R.: Th1 and Th17 cells: adversaries and collaborators. Ann. NY Acad. Sci. 1183, 211-221 (2010).
- Davila, D.R., Edwards III, C.K., Arkins, S., Simon, J., and Kelley, K.W.: Interferongamma-induced priming for secretion of superoxide anion and tumor necrosis factoralpha declines in macrophages from aged rats. FASEB J. 4, 2906-2911 (1990).
- Dejea, C.M., Fathi, P., Craig, J.M., Boleij, A., Geis, A.L., Wu, X., DeStefano Shields, C.E., Hechenbleikner, E.M., Huso, D.L., Anders, R.A., Giardiello, F.M., Wick, E.C., Wang, H., Wu, S., Pardoll, D.M., Housseau, F., and Sears, C.L.: Patients with familial adenomatous polyposis harbor

colonic biofilms containing tumorigenic bacteria. Science 359, 592-597 (2018).

- DelaRosa, O., Tarazona, R., Casado, J.G., Alonso, C., Ostos, B., Pena, J., and Solana, R.: Vα24+ NKT cells are decreased in elderly humans. Exp. Gerontol. 37, 213-217 (2002).
- Díaz, P., Valenzuela Valderrama, M., Bravo, J., and Quest, A.F.G.: *Helicobacter pylori* and Gastric Cancer: Adaptive Cellular Mechanisms Involved in Disease Progression. Front. Microbiol. 9, 1-10 (2018).
- Eibel, H., Kraus, H., Sic, H., Kienzler, A.-K., and Rizzi, M.: B cell Biology: An Overview. Curr. Allergy Asthma Rep. 14, 1-10 (2014).
- Fiebiger, U., Bereswill, S., and Heimesaat, M.M.: Dissecting the interplay between intestinal microbiota and host immunity in health and disease: Lessons learned from germfree and gnotobiotic animal models. Eur. J. Microbiol. Immunol. (Bp.) 6, 253-271 (2016).
- Fulbright, L.E., Ellermann, M., and Arthur, J.C.: The microbiome and the hallmarks of cancer. PLoS Pathog. 13, e1006480 (2017).
- Fülöp, T., Dupuis, G., Witkowski, J.M., and Larbi, A.: The Role of Immunosenescence in the Development of Age-Related Diseases. Rev. Invest. Clin. 68, 84-91 (2016).
- Gálvez, J.: Role of Th17 Cells in the Pathogenesis of Human IBD. ISRN Inflamm. 2014, 928461 (2014).
- Geis, A.L., Fan, H., Wu, X., Wu, S., Huso, D.L., Wolfe, J.L., Sears, C.L., Pardoll, D.M., and Housseau, F.: Regulatory T Cell Response to Enterotoxigenic Bacteroides fragilis Colonization Triggers IL17-Dependent Colon Carcinogenesis. Cancer Discov. 5, 1098-1109 (2015).
- Geller, L.T., Barzily-Rokni, M., Danino, T., Jonas, O.H., Shental, N., Nejman, D., Gavert, N., Zwang, Y., Cooper, Z.A., Shee, K., Thaiss, C.A., Reuben, A., Livny, J., Avraham, R., Frederick, D.T., Ligorio, M., Chatman, K., Johnston, S.E., Mosher, C.M., Brandis, A., Fuks, G., Gurbatri, C., Gopalakrishnan, V., Kim, M., Hurd, M.W.,

Katz, M., Fleming, J., Maitra, A., Smith, D.A., Skalak, M., Bu, J., Michaud, M., Trauger, S.A., Barshack, I., Golan, T., Sandbank, J., Flaherty, K.T., Mandinova, A., Garrett, W.S., Thayer, S.P., Ferrone, C.R., Huttenhower, C., Bhatia, S.N., Gevers, D., Wargo, J.A., Golub, T.R., and Straussman, R.: Potential role of intratumor bacteria in mediating tumor resistance to the chemotherapeutic drug gemcitabine. Science 357, 1156-1160 (2017).

- Gibson, K.L., Wu, Y.-C., Barnett, Y., Duggan, O., Vaughan, R., Kondeatis, E., Nilsson, B.O., Wikby, A., Kipling, D., and Dunn-Walters, D.K.: B-cell diversity decreases in old age and is correlated with poor health status. Aging Cell 8, 18-25 (2009).
- Goodwin, A.C., Shields, C.E.D., Wu, S., Huso, D.L., Wu, X., Murray-Stewart, T.R., Hacker-Prietz, A., Rabizadeh, S., Woster, P.M., Sears, C.L., and Casero Jr., R.A.: Polyamine catabolism contributes to enterotoxigenic Bacteroides fragilisinduced colon tumorigenesis. Proc. Natl. Acad. Sci. USA 108, 15354-15359 (2011).
- Gopalakrishnan, V., Spencer, C.N., Nezi, L., Reuben, A., Andrews, M.C., Karpinets, T.V., Prieto, P.A., Vicente, D., Hoffman, K., Wei, S.C., Cogdill, A.P., Zhao, L., Hudgens, C.W., Hutchinson, D.S., Manzo, T., Petaccia de Macedo, M., Cotechini, T., Kumar, T., Chen, W.S., Reddy, S.M., Szczepaniak Sloane, R., Galloway-Pena, J., Jiang, H., Chen, P.L., Shpall, E.J., Rezvani, K., Alousi, A.M., Chemaly, R.F., Shelburne, S., Vence, L.M., Okhuysen, P.C., Jensen, V.B., Swennes, A.G., McAllister, F., Marcelo Riquelme Sanchez, E., Zhang, Y., Le Chatelier, E., Zitvogel, L., Pons, N., Austin-Breneman, J.L., Haydu, L.E., Burton, E.M., Gardner, J.M., Sirmans, E., Hu, J., Lazar, A.J., Tsujikawa, T., Diab, A., Tawbi, H., Glitza, I.C., Hwu, W.J., Patel, S.P., Woodman, S.E., Amaria, R.N., Davies, M.A., Gershenwald, J.E., Hwu, P., Lee, J.E., Zhang, J., Coussens, L.M., Cooper, Z.A., Futreal, P.A., Daniel, C.R., Ajami, N.J., Petrosino, J.F., Tetzlaff, M.T., Sharma, P., Allison, J.P., Jenq, R.R., and

Wargo, J.A.: Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. Science 359, 97-103 (2018).

- Gur, C., Ibrahim, Y., Isaacson, B., Yamin, R., Abed, J., Gamliel, M., Enk, J., Bar-On, Y., Stanietsky-Kaynan, N., Coppenhagen-Glazer, S., Shussman, N., Almogy, G., Cuapio, A., Hofer, E., Mevorach, D., Tabib, A., Ortenberg, R., Markel, G., Miklić, K., Jonjic, S., Brennan, C.A., Garrett, W.S., Bachrach, G., and Mandelboim, O.: Binding of the Fap2 protein of Fusobacterium nucleatum to human inhibitory receptor TIGIT protects tumors from immune cell attack. Immunity 42, 344-355 (2015).
- Harding, C., Pompei, F., and Wilson, R.: Peak and decline in cancer incidence, mortality, and prevalence at old ages. Cancer 118, 1371-1386 (2012).
- Hayashi, A., Sato, T., Kamada, N., Mikami, Y., Matsuoka, K., Hisamatsu, T., Hibi, T., Roers, A., Yagita, H., Ohteki, T., Yoshimura, A., and Kanai, T.: A single strain of Clostridium butyricum induces intestinal IL-10-producing macrophages to suppress acute experimental colitis in mice. Cell Host Microbe 13, 711-722 (2013).
- Herrera, V. and Parsonnet, J.: Helicobacter pylori and gastric adenocarcinoma. Clin. Microbiol. Infect. 15, 971-976 (2009).
- Herrero, C., Marqués, L., Lloberas, J., and Celada, A.: IFN-γ–dependent transcription of MHC class II IA is impaired in macrophages from aged mice. J. Clin. Invest. 107, 485-493 (2001).
- Hooi, J.K.Y., Lai, W.Y., Ng, W.K., Suen, M.M.Y., Underwood, F.E., Tanyingoh, D., Malfertheiner, P., Graham, D.Y., Wong, V.W.S., Wu, J.C.Y., Chan, F.K.L., Sung, J.J.Y., Kaplan, G.G., and Ng, S.C.: Global Prevalence of Helicobacter pylori Infection: Systematic Review and Meta-Analysis. Gastroenterology 153, 420-429 (2017).
- Hsieh, C., Macatonia, S.E., Tripp, C.S., Wolf, S.F., O'Garra, A., and Murphy, K.M.: Development of Th1 CD4+ T Cells Through IL-12 Produced by Listeria-Induced Macrophages. Science 260, 547-

549 (1993)..

- Iida, N., Dzutsev, A., Stewart, C.A., Smith, L., Bouladoux, N., Weingarten, R.A., Molina, D.A., Salcedo, R., Back, T., Cramer, S., Dai, R.M., Kiu, H., Cardone, M., Naik, S., Patri, A.K., Wang, E., Marincola, F.M., Frank, K.M., Belkaid, Y., Trinchieri, G., and Goldszmid, R.S.: Commensal bacteria control cancer response to rherapy by modulating the tumor microenvironment. Science 342, 967-971 (2013).
- Ivanov, I.I., Atarashi, K., Manel, N., Brodie, E.L., Shima, T., Karaoz, U., Wei, D., Goldfarb, K.C., Santee, C.A., Lynch, S.V., Tanoue, T., Imaoka, A., Itoh, K., Takeda, K., Umesaki, Y., Honda, K., and Littman, D.R.: Induction of intestinal Th17 cells by segmented filamentous bacteria. Cell 139, 485-498 (2009).
- Jiang, Y., Chen, B., Duan, C., Sun, B., Yang, J., and Yang, S.: Multigene editing in the Escherichia coli genome via the CRISPR-Cas9 system. Appl. Environm. Microbiol. 81, 2506-2514 (2015).
- Jing, Y., Gravenstein, S., Chaganty, R.N., Chen, N., Lyerly, K.H., Joyce, S., and Deng, Y.: Aging is associated with a rapid decline in frequency, alterations in subset composition, and enhanced Th2 response in CD1d-restricted NKT cells from human peripheral blood. Exp. Gerontol. 42, 719-732 (2007).
- Kostic, A.D., Chun, E., Robertson, L., Glickman, J.N., Gallini, C.A., Michaud, M., Clancy, T.E., Chung, D.C., Lochhead, P., Hold, G.L., El-Omar, E.M., Brenner, D., Fuchs, C.S., Meyerson, M., and Garrett, W.S.: Fusobacterium nucleatum potentiates intestinal tumorigenesis and modulates the tumor immune microenvironment. Cell Host Microbe 14, 207-215 (2013).
- Larkin, J., Chiarion-Sileni, V., Gonzalez, R., Grob, J.J., Cowey, C.L., Lao, C.D., Schadendorf, D., Dummer, R., Smylie, M., Rutkowski, P., Ferrucci, P.F., Hill, A., Wagstaff, J., Carlino, M.S., Haanen, J.B., Maio, M., Marquez-Rodas, I., McArthur, G.A., Ascierto, P.A., Long, G.V., Callahan, M.K., Postow, M.A., Grossmann, K., Sznol,

M., Dreno, B., Bastholt, L., Yang, A., Rollin, L.M., Horak, C., Hodi, F.S., and Wolchok, J.D.: Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. New Engl. J. Med. 373, 23-34 (2015).

- Linton, P.J. and Dorshkind, K.: Age-related changes in lymphocyte development and function. Nat. Immunol. 5, 133-139 (2004).
- Loeb, L.A., Loeb, K.R., and Anderson, J.P.: Multiple mutations and cancer. Proc. Natl. Acad. Sci. USA 100, 776-781 (2003).
- Ma, C., Han, M., Heinrich, B., Fu, Q., Zhang, Q., Sandhu, M., Agdashian, D., Terabe, M., Berzofsky, J.A., Fako, V., Ritz, T., Longerich, T., Theriot, C.M., McCulloch, J.A., Roy, S., Yuan, W., Thovarai, V., Sen, S.K., Ruchirawat, M., Korangy, F., Wang, X.W., Trinchieri, G., and Greten, T.F.: Gut microbiome-mediated bile acid metabolism regulates liver cancer via NKT cells. Science 360, eaan5931 (2018).
- Marullo, R., Werner, E., Degtyareva, N., Moore, B., Altavilla, G., Ramalingam, S. S., and Doetsch, P.W.: Cisplatin induces a mitochondrial-ROS response that contributes to cytotoxicity depending on mitochondrial redox status and bioenergetic functions. PLoS One 8, 1-15 (2013).
- Matson, V., Fessler, J., Bao, R., Chongsuwat, T., Zha, Y., Alegre, M.-L., Luke, J.J., and Gajewski, T.F.: The commensal microbiome is associated with anti–PD-1 efficacy in metastatic melanoma patients. Science 359, 104-108 (2018).
- Mocchegiani, E., Giacconi, R., Cipriano, C., Gasparini, N., Bernardini, G., Malavolta, M., Menegazzi, M., Cavalieri, E., Muzzioli, M., Ciampa, A.R., and Suzuki, H.: The variations during the circadian cycle of liver CD1d-unrestricted NK1.1+TCR gamma/ delta+ cells lead to successful ageing. Role of metallothionein/IL-6/gp130/PARP-1 interplay in very old mice. Exp. Gerontol. 39, 775-788 (2004).
- Narendra, B.L., Reddy, K.E., Shantikumar, S., and Ramakrishna, S.: Immune system: A double-edged sword in cancer. Inflamm. Res. 62, 823-834 (2013).

- Nougayrede, J.P., Homburg, S., Taieb, F., Boury, M., Brzuszkiewicz, E., Gottschalk, G., Buchrieser, C., Hacker, J., Dobrindt, U., and Oswald, E.: Escherichia coli Induces DNA Double-Strand Breaks in Eukaryotic Cells. Science 313, 848-852 (2006).
- Olszak, T., An, D., Zeissig, S., Vera, M. P., Richter, J., Franke, A., Glickman, J.N., Siebert, R., Baron, R.M., Kasper, D.L., and Blumberg, R.S.: Microbial exposure during early life has persistent effects on natural killer T cell function. Science 336, 489-493 (2012).
- Penders, J., Thijs, C., Vink, C., Stelma, F.F., Snijders, B., Kummeling, I., van den Brandt, P.A., and Stobberingh, E.E.: Factors influencing the composition of the intestinal microbiota in early infancy. Pediatrics 118, 511-521 (2006).
- Pickard, J.M., Zeng, M.Y., Caruso, R., and Núñez, G.: Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease. Immunol. Rev. 279, 70-89 (2017).
- Polk, D.B. and Peek, R.M.: Helicobacter pylori: Gastric cancer and beyond. Nat. Rev. Cancer 10, 403-414 (2010).
- Pompei, F., Polkanov, M., and Wilson, R.: Age distribution of cancer in mice: the incidence turnover at old age. Toxicol. Ind. Health 17, 7-16 (2001).
- Ponnappan, S. and Ponnappan, U.: Aging and Immune Function: Molecular Mechanisms to Interventions. Antioxid. Redox Signal. 14, 1551-1585 (2011).
- Prorok-Hamon, M., Friswell, M.K., Alswied, A., Roberts, C.L., Song, F., Flanagan, P.K., Knight, P., Codling, C., Marchesi, J.R., Winstanley, C., Hall, N., Rhodes, J.M., and Campbell, B.J.: Colonic mucosa-associated diffusely adherent afaC+ Escherichia coli expressing lpfA and pks are increased in inflammatory bowel disease and colon cancer. Gut 63, 761-770 (2014).
- Rajilić-Stojanović, M., Heilig, H.G.H.J., Tims, S., Zoetendal, E.G., and de Vos, W.M.: Long-term monitoring of the human intestinal microbiota composition. Environ. Microbiol. 15, 1146-1159 (2013).

- Renshaw, M., Rockwell, J., Engleman, C., Gewirtz, A., Katz, J., and Sambhara, S.: Cutting edge: impaired Toll-like receptor expression and function in aging. J. Immunol. 169, 4697-4701 (2002).
- Rhee, K.J., Wu, S., Wu, X., Huso, D.L., Karim, B., Franco, A.A., Rabizadeh, S., Golub, J.E., Mathews, L.E., Shin, J., Sartor, R.B., Golenbock, D., Hamad, A.R., Gan, C.M., Housseau, F., and Sears, C.L.: Induction of persistent colitis by a human commensal, enterotoxigenic Bacteroides fragilis, in wild-type C57BL/6 mice. Infect. Immun. 77, 1708-1718 (2009).
- Ridlon, J.M., Kang, D.J., and Hylemon, P.B.: Bile salt biotransformations by human intestinal bacteria. J. Lipid Res. 47, 241-259 (2006).
- Rothstein, D.M. and Camirand, G.: New insights into the mechanisms of Treg function. Curr. Opin. Organ Transplant. 20, 376-384 (2015).
- Round, J.L. and Mazmanian, S.K.: The gut microbiota shapes intestinal immune responses during health and disease. Nat. Rev. Immunol. 9, 313-323 (2009).
- Round, J.L. and Mazmanian, S.K.: Inducible Foxp3+ regulatory T-cell development by a commensal bacterium of the intestinal microbiota. Proc. Natl. Acad. Sci. USA 107, 12204-12209 (2010).
- Routy, B., Le Chatelier, E., Derosa, L., Duong, C.P.M., Alou, M.T., Daillère, R., Fluckiger, A., Messaoudene, M., Rauber, C., Roberti, M.P., Fidelle, M., Flament, C., Poirier-Colame, V., Opolon, P., Klein, C., Iribarren, K., Mondragón, L., Jacquelot, N., Qu, B., Ferrere, G., Clémenson, C., Mezquita, L., Masip, J.R., Naltet, C., Brosseau, S., Kaderbhai, C., Richard, C., Rizvi, H., Levenez, F., Galleron, N., Quinquis, B., Pons, N., Ryffel, B., Minard-Colin, V., Gonin, P., Soria, J.C., Deutsch, E., Loriot, Y., Ghiringhelli, F., Zalcman, G., Goldwasser, F., Escudier, B., Hellmann, M.D., Eggermont, A., Raoult, D., Albiges, L., Kroemer, G., and Zitvogel, L.: Gut microbiome influences efficacy of PD-1based immunotherapy against epithelial

tumors. Science 359, 91-97 (2018).

- Rubinstein, R.M., Wang, X., Liu, W., Hao, Y., Cai, G., and Han, Y.W.: Fusobacterium nucleatum promotes colorectal carcinogenesis by modulating E-cadherin/β-catenin signaling via its FadA adhesin. Cell Host Microbe 14, 195-206 (2013).
- Schwabe, R.F. and Jobin, C.: The microbiome and cancer. Nat. Rev. Cancer 13, 800-812 (2013).
- Secher, T., Samba-Louaka, A., Oswald, E., and Nougayrède, J.P.: Escherichia coli producing colibactin triggers premature and transmissible senescence in mammalian cells. PLoS One 8, e77157 (2013).
- Shabbir, M.A.B., Hao, H., Shabbir, M.Z., Wu, Q., Sattar, A., and Yuan, Z.: Bacteria vs. Bacteriophages: Parallel Evolution of Immune Arsenals. Front. Microbiol. 7, 1292 (2016).
- Shields, B.D., Mahmoud, F., Taylor, E.M., Byrum, S.D., Sengupta, D., Koss, B., Baldini, G., Ransom, S., Cline, K., Mackintosh, S.G., Edmondson, R.D., Shalin, S. and Tackett, A.J.: Indicators of responsiveness to immune checkpoint inhibitors. Sci. Rep. 7, 807 (2017).
- Smith, B.D., Smith, G.L., Hurria, A., Hortobagyi, G.N., and Buchholz, T.A.: Future of cancer incidence in the United States: burdens upon an aging, changing nation. J. Clin. Oncol. 27, 2758-2765 (2009).
- Stringer, A.M., Gibson, R.J., Bowen, J.M., Logan, R.M., Ashton, K., Yeoh, A.S. J., Al-Dasooqi, N., and Keefe, D.M.K.: Irinotecan-induced mucositis manifesting as diarrhoea corresponds with an amended intestinal flora and mucin profile. Int. J. Exp. Pathol. 90, 489-499 (2009).
- Sun, M., He, C., Cong, Y., and Liu, Z.: Regulatory immune cells in regulation of intestinal inflammatory response to microbiota. Mucosal Immunol. 8, 969-978 (2015).
- Swift, M.E., Burns, A.L., Gray, K.L., and DiPietro, L.A.: Age-related alterations in the inflammatory response to dermal injury. J. Invest. Dermatol. 117, 1027-1035 (2001).

- Thakkar, J.P., McCarthy, B.J., and Villano, J.L.: Age-specific cancer incidence rates increase through the oldest age groups. Am. J. Medical Sci. 348, 65-70 (2014).
- Thursby, E. and Juge, N.: Introduction to the human gut microbiota. Biochem. J. 474, 1823-1836 (2017).
- Tilg, H., Adolph, T.E., Gerner, R.R., and Moschen, A.R.: The Intestinal Microbiota in Colorectal Cancer. Cancer Cell 33, 1-11 (2018).
- Tomkovich, S., Yang, Y., Winglee, K., Gauthier, J., Mühlbauer, M., Sun, X., Mohamadzadeh, M., Liu, X., Martin, P., Wang, G.P., Oswald, E., Fodor, A.A., and Jobin, C.: Locoregional effects of microbiota in a preclinical model of colon carcinogenesis. Cancer Res. 77, 2620-2632 (2017).
- Vinay, D.S., Ryan, E.P., Pawelec, G., Talib, W.H., Stagg, J., Elkord, E., Lichtor, T., Decker, W.K., Whelan, R.L., Kumara, H.M.C.S., Signori, E., Honoki, Κ., Georgakilas, A.G., Amin, A., Helferich, W.G., Boosani, C.S., Guha, G., Ciriolo, M.R., Chen, S., Mohammed, S.I., Azmi, A.S., Keith, W.N., Bilsland, A., Bhakta, D., Halicka, D., Fujii, H., Aquilano, K., Ashraf, S.S., Nowsheen, S., Yang, X., Choi, B.K., and Kwon, B.S.: Immune evasion in cancer: Mechanistic therapeutic basis and strategies. Semin. Cancer Biol. 35 Suppl., S185-S198 (2015).
- Wenisch, C., Patruta, S., Daxböck, F., Krause, R., and Hörl, W.: Effect of age on human neutrophil function. J. Leukoc. Biol. 67, 40-45 (2000).

- White, M.C., Holman, D.M., Boehm, J.E., Peipins, L.A., Grossman, M., and Henley, S.J.: Age and cancer risk: A potentially modifiable relationship. Am. J. Prev. Med. 46, Suppl. 1, S7-S15 (2014).
- Wu, H.J. and Wu, E.: The role of gut microbiota in immune homeostasis and autoimmunity. Gut Microbes 3, 4-14 (2012).
- Wu, S., Dreyfus, L.A., Tzianabos, A.O., Hayashi, C., and Sears, C.L.: Diversity of the metalloprotease toxin produced by enterotoxigenic Bacteroides fragilis. Infect. Immun. 70, 2463-2471 (2002).
- Wu, S., Rhee, K., Albesiano, E., Rabizadeh, S., Wu, X., Yen, H., Huso, D.L., Brancati, F.L., Wick, E., McAllister, F., Housseau, F., Pardoll, D.M., and Sears, C.L.: A human colonic commensal promotes colon tumorigenesis via activation of T helper type 17 T cell responses. Nat. Med. 15, 1016-1022 (2009).
- Yoshimoto, S., Loo, T. M., Atarashi, K., Kanda, H., Sato, S., Oyadomari, S., Iwakura, Y., Oshima ,K., Morita, H., Hattori, M., Honda, K., Ishikawa, Y., Hara, E., and Ohtani, N.: Obesity-induced gut microbial metabolite promotes liver cancer through senescence secretome. Nature 499, 97-101 (2013).
- Yu, T., Guo, F., Yu, Y., Sun, T., Ma, D., Han, J., Qian, Y., Kryczek, I., Sun, D., Nagarsheth, N., Chen, Y., Chen, H., Hong, J., Zou, W., and Fang, J.Y.: Fusobacterium nucleatum Promotes Chemoresistance to Colorectal Cancer by Modulating Autophagy. Cell 170, 548-563 (2017).

AGEING AND THE MICROBIOME: SUMMARY OF THE 32ND OLD HERBORN UNIVERSITY SEMINAR AND THE DISCUSSIONS

MICHAEL ZASLOFF

MedStar Georgetown Transplant Institute, Georgetown University School of Medicine, Washington, DC 20007, USA

Following a warm introduction, *Volker Rusch* opened the Seminar. The focus of this year's meeting was ageing, with an emphasis on whether the gut microbiome played any role in the process.

Michael Zasloff presented a brief introductory lecture ("Some Insights about Ageing from a Shark and an Oyster"). He pointed out that the two documented longest-lived animals were the "Ming Clam" (400 years) and the Greenland shark (500 years). Their survival obviously required nonsenescent stem cells, and an immune system that could protect against all potential pathogens. He pointed out that the details of the animal "ageing clock" is unknown. Zasloff finished his presentation with a discussion of recently published studies in mice that suggested that hypothalamic inflammation leading to the reduced expression of certain hypothalamic hormones (i.e. GnRH) drives peripheral ageing. These observations suggest that an "elixir of youth" might be more real than we believe.

Robert Pignolo presented a general overview of the physiology of human ageing from the perspective of a research gerontologist ("*Physiology of Human Ageing*"). He made the point that it becomes difficult to separate ageing per se from the onset of disease processes that are associated with ageing. If we could slow down ageing, how would we document the response? One approach is to focus on the loss of function of a system, such as the GI tract, the immune system, the endocrine system and the nervous system cataloguing the known progressive changes that occur in the otherwise healthy person. He highlighted the known progressive changes that occur with respect to specific tissues, such as muscle, bone, skin and adipose tissue. He stressed that a deeper analysis of these processes could be extended mechanistically to dissect the specific pathways leading to the age-related changes. Ageing in humans is frequently associated with a loss of resilience and those factors that increase an individual's fragility, and reduce their probability of recovery. He stressed that much current interest in human ageing is focused on the mechanistic basis of reduced stem cell survival, cellular senescence and the underlying mechanisms responsible for chronic inflammation. In particular, considerable research in human ageing is now focused on drugs that deplete senescent cells.

Brian Kennedy posed the provocative question of whether we could intervene to delay human ageing ("The Age of Ageing: Can We Intervene"). He reviewed the known conditions that have been shown in various systems to extend lifespan, such as caloric restriction. Of particular interest were small molecules that had already been developed and were currently in human use for various medical conditions. In particular he highlighted metformin and rapamycin. While the mechanism underlying the benefit of metformin was still unclear, rapamycin's effects on ageing appear to be attributable to mTOR inhibition. mTOR regulates protein synthesis in response to nutrient and energy availability. Recent studies suggest that mTOR inhibition extends stem cell survival. He presented remarkable studies that supported the effect of mTOR inhibition on age related stem cell loss in the murine trachea.

He then tackled the persistent theme of how best to measure the impact of an intervention that delayed ageing in humans. One promising potential biomarker is the accumulation of methylation across the 450K CpG site assessable in the human genome. As an individual age the number of methylated CpG sites increases providing an "epigenetic clock" that might be of use. Kennedy described another approach his colleagues have explored, that being facial recognition. Facial recognition technology has advanced to such an extent that chronological age can be reliably estimated through this modality, possibly providing a tool to assess the impact of anti-ageing interventions.

Dario Riccardo Valenzano presented his work with the turquoise killifish as a new model animal to study the ageing process ("Life Span Regulation by the Gut Microbiota in the Naturally Short Lived African Turquoise Killi*fish*"). The killifish has a lifespan of between11-26 weeks, bridging the brief lifespan of C. elegans (3 weeks) and Drosophila (9-13 weeks) and the longer lifespan of the mouse (130 weeks). This species is indigenous to Africa. The genomic sequence was published in 2015 as well as techniques to genetically modify the animals. Ageing in killifish is characterized by dramatic skeletal disturbances including a form of scoliosis, reduced capacity to learn, decreased fertility, slower overall movement, sarcopenia, and gut

fibrosis. Valenzano noted that the microbiome of the killifish changes dramatically as the animal ages, and then posed the question of whether the gut microbiome influenced ageing. The most abundant taxa in its microbiome are shared with mammals, and common microbial taxa are shared between animals caught in the wild and those maintained in captivity. Can the course of ageing be changed by altering the Microbial microbiome? transplants from young or old fish were introduced into middle aged fish that had been pretreated with antibiotics. The most dramatic outcome of this study was with young microbial transplants: survival was extended, swimming activity was increased to a level normal for young animals, changes were seen at the transcriptional level in the intestinal tissues. He hypothesized that the effects observed could in part be due to the differences in serum metabolites associated with the young or old gut microbial communities and the interaction with host physiology.

Christoph Kaleta presented a broad overview of ageing ("*Evaluating Potential Contributions of the Microbiome to Ageing Pathologies*"). His overriding thesis was that a major factor in the ageing process is due to the accumulation of dead or damaged cells, which, in turn, provokes chronic inflammation. Might the ageing microbiome impact on the clearance of senescent cells? Perhaps metabolites of microbiome origin play a role. And if so, we might be able to intervene.

Thomas C. G. Bosch presented an expansive perspective on ageing focusing on his studies of Hydra ("Ageing and the Microbiome: Lessons from Non-Senescent Models"). He stressed that both genetics and environment play a role in longevity. He highlighted the remarkable role of FoxO in the biology of Hydra, FoxO being one of the two genes that have been associated with longevity in humans. A three classes of stem cells in *Hydra* express FoxO, and genetic inhibition of FoxO expression slows down the population growth rate in this animal. Remarkably, FoxO also plays a role in managing the microbiome of *Hydra*. It turns out that FoxO is required for expression of antimicrobial peptides, which in turn, regulate the composition and density of the microbiome. So, by virtue of FoxO's integration into the design of *Hydra*, it controls cross-talk between the stem cell and the microbiome. This relationship is dramatically altered in the setting of a *Hydra* strain in which a tumour has arisen. The presence of the tumour, which is of an epithelial nature, dramatically alters the microbiome, increasing the proportion of spirochetes. These spirochetes are not passive bystanders, since their elimination with antibiotics causes regression of the tumour. Re-inoculation with these bacteria causes a recurrence of the tumour. Careful analysis of the microbial interactions involved, demonstrate that a Pseudomonas strain is also required for tumorigenesis, implicating the two species in this process. Bosch stressed that this story underscores the principle that the physiology of an organism cannot be understood without including the microbial elements with which coexists, without viewing the entirety of the holobiont.

Wolfgang Kunze focused on the ageing of the enteric nervous system (ENS) and how gut microbes might influence the activity of the ENS ("Ageing and Gut to Brain Signalling"). He reminded us that gut microbiota change with ageing and that feeding Bifidobacteria to mice has been shown to increase lifespan. He suggested that gut microbes could contribute to ageing through neural communication to the brain, for example, via the vagus.

When the Lactobacillus strain, JB-1 is introduced into the mouse gut, 90% of the signals stimulated by JB-1 are transmitted through vagal afferents. He highlighted that as we age the ENS progressively deteriorates. Peristalsis is reduced. Afferent activity decreases. Ion channel activity, such as the Ik potassium channel, diminishes. The consequences of this deterioration can be observed in the brain. For one, stimulation of the vagus has been shown to activate growth hormone releasing factor in the hypothalamus, demonstrating a clear neural linkage between the hypothalamus and the gut. Introduction of JB-1 or squalamine into the intestine (studied *ex vivo*) leads to increased vagal firing in aged mice, suggesting that the diminished activity of the aged ENS can be corrected therapeutically.

Josef Anrather introduced the concept that inflammation within the brain surrounding stroke could be modulated by the gut microbiome ("From the Gut to the Brain: The Roles of Microbiota in Stroke"). He described several models of stroke as studied in rodents. The basic premise of his hypothesis is that IL-17 T cells are major players in the inflammatory events that follow a stroke, that the gut is the major reservoir of IL-17 T cells, and that the microbiome plays a role in the differentiation of IL-17 T cell population. Following a stroke intestinal IL-17 T cells traffic to sites such as the meninges, and then into the brain. By altering the population of intestinal microbes, it is possible to influence the dendritic cells of the intestinal lamina propria to shift the T cell population towards Tregs. He has begun studies that suggest that a factor extractable from the "good" flora can recapitulate the beneficial effects on the intestinal T cell population of the gut. These studies suggest an entirely novel understanding of the pathophysiology of the events that follow a stroke, and treatments that might be developed.

Janelle C. Arthur focused on the role of the microbiome in colorectal cancer, a disease of ageing ("The Microbiome and Cancer"). She reminded us that the current explanation for the occurrence of colorectal cancer is chronic inflammation, caused either by microbes or by organic compounds of dietary origin present in faeces. From studies with IL-10 knock out mice, an animal that develops colorectal cancer, it is clear that the microbiome plays a role, since germ-free animals of this strain do not exhibit this phenotype. Arthur asked whether specific microbes might be involved. Since inflammation precedes the appearance of cancer in these animals, might inflammation produce a change in the microbiome that encourages the appearance of carcinogenic microbes? To answer these questions, she used germ-free IL-10 knock out mice. The animals were inoculated with SPF microbiota, then treated with azoxymethane, a chemical that provokes both intestinal inflammation and colorectal cancer. The microbiome was then examined and compared with the initial inoculum with respect to its species. Treatment with Azoxymethane led to a 100-fold increase in E. coli. If the same experiment was performed with either a pure culture inoculum of E. coli or E. fae*calis* (rather than the SPF inoculum) only the E. coli inoculated mice developed significant colorectal cancers. It appears that strains of E. coli that harbour the polyketide synthase (pks) pathogenicity island appear to be the culprits. A specific substance produced by these strains of E. coli can induce cancer independently of inflammation. The product of the *pks* gene cluster is a poorly characterized molecule called Colibactin which has been shown to cause chromosomal damage by an unknown mechanism and requires contact between the bacterial cell and the epithelium. These studies suggest that age related changes in the microbiome could be a cause of colorectal cancer, and clearly amenable to intervention.

The formal Academic ceremony followed the completion of the Seminar, with the granting of Honorary Professorships at the Old Herborn University to the speakers.